

This Page Is Inserted by IFW Operations
and is not a part of the Official Record

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images may include (but are not limited to):

- BLACK BORDERS
- TEXT CUT OFF AT TOP, BOTTOM OR SIDES
- FADED TEXT
- ILLEGIBLE TEXT
- SKEWED/SLANTED IMAGES
- COLORED PHOTOS
- BLACK OR VERY BLACK AND WHITE DARK PHOTOS
- GRAY SCALE DOCUMENTS

IMAGES ARE BEST AVAILABLE COPY.

**As rescanning documents *will not* correct images,
please do not report the images to the
Image Problems Mailbox.**

ADVANCES IN
PHYSIOLOGY EDUCATION
See Part 3, S1-S54

Univ. of Minn.
Bio-Medical
Library

1 11 93

DECEMBER 1992

Volume 263/Number 6

Part 1 of Three Parts

ISSN 0002-9513

American Journal of Physiology

PUBLISHED BY
THE AMERICAN PHYSIOLOGICAL SOCIETY

Central hypertensinogenic effects of glycyrrhizic acid and carbenoxolone

ELISE P. GOMEZ-SANCHEZ AND CELSO E. GOMEZ-SANCHEZ

Research Service and Department of Internal Medicine, James A. Haley Veterans Hospital, and University of South Florida Health Science Center, Tampa, Florida 33612

Gomez-Sanchez, Elise P., and Celso E. Gomez-Sanchez. Central hypertensinogenic effects of glycyrrhizic acid and carbenoxolone. *Am. J. Physiol.* 263 (Endocrinol. Metab. 26): E1125-E1130, 1992.—The apparent mineralocorticoid excess syndrome of patients ingesting large amounts of licorice or its derivatives is thought to be caused by the antagonism by these compounds of the enzyme 11β -hydroxysteroid dehydrogenase (11β -HSD). 11β -HSD inactivates cortisol and corticosterone, allowing the more abundantly produced glucocorticoids access to the mineralocorticoid receptor (MR) in the kidney, where they act as mineralocorticoids. We have found that the infusion of both glycyrrhizic acid, an active principle of licorice, and carbenoxolone, a synthetic analogue, into a lateral ventricle of the brain [intracerebroventricular (icv)] of a rat, at a dose less than that which has an effect when infused subcutaneously, produces hypertension. Furthermore, the hypertension produced by the oral administration of carbenoxolone or glycyrrhizic acid is blocked by the icv administration of RU 28318, an MR antagonist, at a dose below that which has an effect on blood pressure when infused subcutaneously. While the oral administration caused saline polydipsia and polyuria typical of chronic systemic mineralocorticoid excess, the icv licorice derivatives produced hypertension without affecting saline appetite. Sensitizing the rats to mineralocorticoid hypertension by renal mass reduction and increasing salt consumption was not necessary for the production of hypertension. These findings provide additional evidence for a central role in blood pressure control by mineralocorticoids that is distinct from their renal effects. They also suggest that more is involved in licorice-induced hypertension than only inhibition of 11β -HSD.

hypertension; licorice; mineralocorticoids; RU 28318; steroid 11β -hydroxysteroid dehydrogenase

ALDOSTERONE acts through type I receptors, or mineralocorticoid receptors (MR), in the kidney to produce sodium retention and potassium and hydrogen ion excretion. The MR is widely distributed and is present in the colon, parotid, vasculature, and, in particular, specific areas of the brain (5, 13). The affinity of isolated MR from various sources, including expressed MR cDNA in COS cells, is similar for aldosterone, corticosterone, and cortisol (3, 4, 16). MR, regardless of the source, are physicochemically identical (16, 32), and appear to be a product of the same cDNA (3). Corticosterone and cortisol normally do not act as mineralocorticoids in the kidney in vivo. Specificity, originally thought to be intrinsic to the receptor, has been shown to be conferred extrinsically by corticosterone/cortisol-binding globulin (CBG), which reduces free circulating glucocorticoid available to the receptor, and by 11β -hydroxysteroid dehydrogenase (11β -HSD). 11β -HSD reversibly converts corticosterone and cortisol to the inactive 11 -dehydrocorticosterone and cortisone (7, 9, 12). The location of the 11β -HSD enzyme has been controversial. It appears that 11β -HSD is expressed in some mineralocorticoid target cells along with the MR, thus

serving as an autocrine control, as well as in cells proximate to MR-containing cells, serving a paracrine function (6, 9, 21, 24).

Under normal conditions, most MR in the rat brain are almost fully occupied by corticosterone, while occupation of the type II receptor, or glucocorticoid receptor (GR), for which corticosterone has less affinity, is less complete and follows the circadian rhythm of glucocorticoid levels (7). It has been suggested that the occupation of the MR in the brain, particularly in the hippocampus, by corticosterone at low, physiological serum levels is possible because CBG does not penetrate the blood-brain barrier (7, 9) and because the activity of 11β -HSD in this organ is negligible (9, 12). However, in situ hybridization techniques have demonstrated the presence of 11β -HSD in the brain (19), as well as the kidney. Whether 11β -HSD is bioactive in any, all, or only specific parts of the brain is controversial (9, 19, 21). There are different tissue-specific forms and regional activity of the 11β -HSD enzyme (20) that may account for the apparent "glucocorticoid-selective" MR in some parts, particularly the hippocampus, of the brain, in contrast to the "aldosterone-preferring" MR in the anterior hypothalamus (7, 18). Seckl et al. (27) have reported that 11β -HSD inhibition by glycyrrhetic acid in vivo in rats increased 2-deoxy- $[^{14}\text{C}]$ glucose use in those areas of the brain where 11β -HSD mRNA expression has been documented. Corticosterone and aldosterone have different actions in some areas of the brain, even though both are thought to be acting with the same affinity and through the same receptor. Aldosterone antagonizes important central nervous system (CNS) effects of corticosterone (7, 26); corticosterone blocks the hypertension induced by the intracerebroventricular (icv) infusion of aldosterone (13, 15).

Apparent mineralocorticoid excess is a rare hypertensive syndrome in which patients have all of the manifestations of excessive production of mineralocorticoids, including hypokalemia, but steroid measurements are normal or low. The defect has been identified as a deficiency in 11β -HSD (11, 28, 30, 31). The pseudohyperaldosteronism, including hypokalemia and low-renin hypertension, produced by excessive licorice consumption and the treatment of peptic ulcers with licorice derivatives or their synthetic analogues has been attributed to the inhibition of this enzyme, allowing the more abundant circulating cortisol/corticosterone access to the MR in the kidney (9). Licorice derivatives and the synthetic analogue carbenoxolone have been used to study the mechanisms responsible for the syndrome of apparent mineralocorticoid excess, as well as the extrinsic factors conferring apparent ligand specificity to the MR (8, 10, 22). We herein describe studies of the central and

NOTICE: THIS MATERIAL MAY BE PROTECTED BY COPYRIGHT LAW (TITLE 17 U.S. CODE)

systemic effects of the icv, subcutaneous (sc), and oral administration of glycyrrhizic acid, a derivative of licorice, and carbenoxolone, a synthetic analogue, on the blood pressure using the specific MR antagonist RU 28318 (14) to inhibit the MR.

METHODS

Cannulas were placed into the right lateral cerebral ventricles of male outbred Sprague-Dawley rats weighing 180–200 g, using aseptic surgical technique under a combination of fentanyl and droperidol (Innovar-Vet, Pitman-Moore), 0.01 ml/100 g body wt sc, as preanesthetic and isoflurane as anesthetic. Rats received standard food (0.3% NaCl) and tap water or 0.9% saline ad libitum to amplify the hypertension as detailed below. Implanted miniosmotic pumps (Alzet 2002, Alza, Palo Alto, CA), which delivered $0.49 \pm 0.02 \mu\text{l/h}$ for 14 days, were used for icv and sc infusions. Pumps were changed on day 14 under isoflurane anesthesia, and pumps of the same lot were used throughout the experiment to ensure consistency. Carbenoxolone, RU 28318, and corticosterone were dissolved in cerebrospinal fluid (CSF) or 0.86% NaCl with 10% propyleneglycol for icv and sc infusion. A potassium gluconate solution that delivered the same amount of K^+ as the RU 28318 solution was used as control for the mineralocorticoid antagonist experiments (14). Reagents were purchased from Sigma, except for the RU 28318, which was a gift from Roussel (Romainville, France). All solutions were made and sterilized by filtration through $0.2\text{-}\mu\text{m}$ filters (Acrodisc 13, Gelman Scientific) immediately before filling and implanting the pumps. Oral carbenoxolone or glycyrrhizic acid was administered individually twice a day as 0.1 or 0.2 ml of a slurry mixed in corn syrup that the rats accepted readily. Indirect systolic blood pressures (IITC, Woodhills, CA) and weights were measured twice a week starting before treatment as described previously (13). Twenty-four- or forty-eight-hour urine volumes were measured once a week in a stainless steel rat metabolism cage.

Effect of icv administration of carbenoxolone: dose response. Carbenoxolone was infused icv at a rate of 0.3, 1.0, and $3.0 \mu\text{g/h}$ and sc at a rate of $3.0 \mu\text{g/h}$ into intact rats provided with 0.9% saline to drink ad libitum.

Effect of icv administration of carbenoxolone and corticosterone. Carbenoxolone was infused icv at a rate of $5.0 \mu\text{g/h}$ and corticosterone at a rate of 20 ng/h , alone and together. Two types of experiments were done. For one, the rats were intact and drank tap water ad libitum. For the other, the right kidneys were removed and the rats drank 0.9% saline ad libitum to be comparable to the classical maneuvers used to amplify mineralocorticoid hypertension.

Effect of oral administration of carbenoxolone with and without RU 28318. Carbenoxolone was administered orally in corn syrup 45 mg/kg twice daily for 10 days and increased to 90 mg/kg twice daily for the next 4 days to ascertain that the hypertensive effect was maximal; the control rats received corn syrup orally. RU 28318 was infused icv at $1.1 \mu\text{g/h}$ in one-half of the animals receiving carbenoxolone; the other animals received a potassium gluconate solution to supply the equivalent amount of K^+ icv. We have previously shown that $1.1 \mu\text{g/h}$ RU 28318 icv has no intrinsic effect on the blood pressure but protects the rat from the hypertension of systemic mineralocorticoid excess, while being well below the dose required to affect on the blood pressure when infused sc (13, 14). The rats were intact and drank tap water ad libitum.

Effect of oral administration of glycyrrhizic acid with and without RU 28318. The effects of both glycyrrhizic acid and carbenoxolone were studied because of evidence that carbenoxolone may have a larger range of effects, including the inhibition of 11- α -oxidoreductase, than does glycyrrhizic acid (29).

Glycyrrhizic acid was administered orally in corn syrup 35 mg/kg twice daily for 14 days. RU 28318 was infused icv and sc at $1.1 \mu\text{g/h}$ in two of three glycyrrhizic acid groups; the other glycyrrhizic acid animals received a potassium gluconate solution icv to supply the equivalent amount of K^+ icv. Another group received corn syrup orally and the potassium gluconate solution icv. The rats were intact and drank tap water ad libitum.

Animals were killed at the end of the studies by CO_2 narcosis and asphyxiation. Autopsies, including dye infusions to check cannula placement, were done at the conclusion of the study, and data from any animal in which there was doubt about the delivery of the solutions or which had evidence of illness causing undue stress were eliminated from the experiment. At the time of the biweekly pump changes, if the catheter was found to be disconnected from the pump or cannula, the data from the preceding two weeks were discarded and the animal eliminated from the study. We started with 8–10 animals per group so that the groups were never reduced to fewer than 7 animals by the end of the experiment. Data were compared by analysis of variance and the Dunnett t and Fisher PLSD tests (StatView 512+, BrainPower, Calabasas, CA).

RESULTS

Carbenoxolone, $3 \mu\text{g/h}$ administered icv to intact rats drinking 0.9% saline ad libitum, increased the blood pressure of rats significantly ($P < 0.01$) within 3 days and was maximal by day 5 (Fig. 1). There was no significant change in the blood pressure of rats receiving $0.3 \mu\text{g/h}$ CSF, or $1 \mu\text{g/h}$ carbenoxolone icv or $3 \mu\text{g/h}$ carbenoxolone sc over 14 days. No significant difference was found in rate of weight gain or 24-h urine volume between any groups in the icv studies. In separate studies it was found that doses of carbenoxolone $>5 \mu\text{g/h}$ resulted in precipitation of the drug in the pump and cannulas.

The icv infusion of corticosterone at 20 ng/h , a dose known to inhibit the hypertension produced by the icv infusion of aldosterone (15) while having no effect in and of itself, did not significantly blunt the increase in blood pressure produced by icv carbenoxolone, nor did it have any effect on the blood pressure by itself (Fig. 2). There was no difference in urine volume or weight gain between groups in the same experiments. Removing one kidney and giving saline to drink did not alter the hypertension




Fig. 1. Effect on indirect systolic blood pressure (BP) at day 5 of intracerebroventricular (icv) and subcutaneous (sc) infusion of carbenoxolone (Cx) at $3.0 \mu\text{g/h}$ in intact rats drinking 0.9% saline ad libitum. CSF, cerebrospinal fluid.

HYPERTENSION, CARBENOXOLONE, AND GLYCYRRHIZIC ACID

E1127




Fig. 2. Effect on indirect systolic blood pressure of icv infusion of carbenoxolone at 5.0 µg/h and corticosterone at 20 ng/h, alone and together, in nonsensitized rats (A; intact and drinking tap water ad libitum) compared with sensitized rats (B; one kidney removed and drinking 0.9% saline ad libitum). β , 11 β -hydroxysteroid dehydrogenase. * $P < 0.05$. ** $P < 0.01$.




Fig. 3. Effect on indirect systolic blood pressure of oral administration of carbenoxolone in corn syrup at 45 mg/kg twice daily for 10 days, then 90 mg/kg twice daily for the next 4 days, while receiving an icv infusion of either RU 28318 at 1.1 ng/h or vehicle, in intact rats drinking tap water ad libitum.

produced by icv carbenoxolone or the effect of icv corticosterone. At day 11 of the sensitization study there was a 41 and 39% difference in blood pressure between the controls and the icv carbenoxolone and icv carbenoxolone plus corticosterone, respectively, compared with 41 and 31% increases for the nonsensitized rats.

The blood pressure of intact rats drinking water and receiving oral carbenoxolone at 45 mg/kg twice daily increased significantly within 6 days from 105 mmHg to a plateau of 127 mmHg (Fig. 3). Doubling the dose to 90

mg/kg twice daily did not further increase the blood pressure. The icv infusion of 1.1 µg/h RU 28318 completely prevented the increase in blood pressure. We have shown in multiple studies, including those described below using glycyrrhizic acid instead of carbenoxolone, that the sc infusion of 1.1 µg/h RU 28318 is too low to affect the blood pressure. We have also reported that the icv infusion of the antagonist at three times this dose has no effect on the blood pressure of normal animals (14). The blood pressure in the animals receiving the icv control solution returned to normal within 3 days of discontinuing the oral administration of carbenoxolone. Orally administered carbenoxolone doubled the urine volume; this increase in urine volume was not prevented by the icv administration of the mineralocorticoid antagonist, which abolished the hypertension (Fig. 4). There was no difference in weight gain between groups.

The oral administration of glycyrrhizic acid at 35 mg/kg twice daily also significantly increased the blood pressure of intact rats drinking tap water. The icv infusion of 1.1 µg/h RU 28318 prevented the rise in blood pressure (Fig. 5). There was no difference in weight gain between groups.

DISCUSSION

The importance of the CNS in the development of mineralocorticoid hypertension has been well documented (5, 13). MR are found in the hippocampus, amygdala, lateral septum, and hypothalamus, particularly in the periventricular regions, areas known to be or suspected of being important in the regulation of adrenocorticotrophic hormone (ACTH) release, arousal, fluid and fluid osmolality equilibrium, and the maintenance of normal blood pressure. The chronic icv infusion of aldosterone at a dose two orders of magnitude less than that necessary to produce hypertension when infused sc has been reported to produce hypertension in rats and dogs (21). The icv infusion of the mineralocorticoid antagonist RU 28318, at doses that have no effect on the blood pressure when given icv alone and that are ineffective as




Fig. 4. Effect on 24-h urine volume of oral administration of carbenoxolone in corn syrup at 45 mg/kg twice daily for 10 days, then 90 mg/kg twice daily for the next 4 days, while receiving an icv infusion of either RU 28318 at 1.1 ng/h or vehicle, in intact rats drinking tap water ad libitum.




Fig. 5. Effect on indirect systolic blood pressure of oral administration of glycyrrhizic acid (GI) in corn syrup at 35 mg/kg twice daily, while receiving an icv infusion of either RU 28318 at 1.1 ng/h or vehicle, at day 13 in intact rats drinking tap water ad libitum.

an antagonist when administered sc, blocks the hypertension of both the icv and systemic administration of aldosterone and the sc infusion of deoxycorticosterone acetate. The systemic, but not icv, aldosterone hypertension is associated with a chronically increased urine volume indicative of saline polydipsia/polyuria. The icv infusion of the antagonist prevents the rise in pressure produced by the systemic administration of aldosterone without preventing the associated polydipsia/polyuria (13, 14). These findings suggest distinct mineralocorticoid effects in the brain and kidney.

In the studies reported herein, the icv, but not sc, infusion of 3 µg/h carbenoxolone produced hypertension, implying that the site of action is in the brain. The hypertension produced by the oral and icv administration of carbenoxolone or glycyrrhizic acid resembles that of chronic systemic and icv aldosterone infusion in the amplitude of the increase in blood pressure and the effectiveness of mineralocorticoid receptor blockade by icv RU 28318 (13, 14). In addition, as with aldosterone, an increase in urine volume occurred only with the systemic, and not icv, administration of hypertensinogenic amounts of both licorice compounds. Blocking the hypertension of animals receiving oral carbenoxolone with the icv infusion of RU 28318 at doses too low to be effective when infused sc did not reduce their increase in urine volume. Classically mineralocorticoid-salt hypertension is associated with an initial retention of sodium and water followed by an "escape" from further retention and the establishment of a new equilibrium at a higher overall fluid volume. Polydipsia/polyuria may persist after reaching a balance with no additional net gain in water (13). Assuming that the carbenoxolone when given orally is causing a mineralocorticoid excess syndrome as far as the kidneys are concerned, one would expect initial sodium and water retention, followed by escape. The rats in these studies apparently were placed in metabolism cages after the water retention phase, assuming it occurred, after an equilibrium had been reached, because their urine output was consistently greater, not less, than that of controls. Weight gains were "real," not water gains, as evidenced by the fact that the weights of the oral carbenoxolone rats did not fall after the drug was withdrawn.

There was a consistent difference in the time of onset of the hypertension. Icv aldosterone hypertension takes

from 7 to 11 days to become significant (13), while icv carbenoxolone hypertension was evident in 3-6 days. Considering the relatively long delay of onset, that of days rather than minutes or hours, it seems unlikely that this difference is due to a more rapid passage of the licorice compounds across the blood-brain barrier; it probably reflects a more basic difference in the mechanism of action. Removing one kidney and giving saline to drink did not exacerbate the hypertension produced by icv carbenoxolone. This was surprising because the classical way to amplify mineralocorticoid hypertension is to reduce renal mass and increase sodium consumption and because in the model of central mineralocorticoid hypertension, equihypertensinogenic doses of icv aldosterone in non-sensitized rats were nine times that of sensitized rats (13).

Glycyrrhizic acid and carbenoxolone are not thought to act as agonists at the receptor level because their affinity for the MR is negligible (2). They are presumed to work by inhibiting 11β-HSD, thereby removing the protection of the MR from corticosterone and allowing it to act as a mineralocorticoid (12). However, if 11β-HSD were active in the brain, and if it were inhibited by carbenoxolone, previous studies from our laboratory suggest that the resulting accumulation of corticosterone would not be expected to increase blood pressure. An additional difference between the icv aldosterone and icv carbenoxolone models is that the icv infusion of corticosterone, at a dose that would have been expected from our previous work to antagonize the icv aldosterone model, had no effect on the blood pressure of rats receiving icv carbenoxolone. It is assumed that the inhibitory action of icv corticosterone on icv aldosterone hypertension is mediated by the MR because RU 26988, a selective GR agonist, had no effect when infused alone or in combination with aldosterone (15).

While most reported studies indicate that carbenoxolone does not affect the mineralocorticoid activity of aldosterone (25), others suggest that it enhances the sodium retention produced by aldosterone and 11-deoxycorticosterone (23). Glycyrrhetic acid has been found to inhibit the hepatic 5β-reductase and 3β-HSD but not the 5α-reductase or 3α-HSD. Another proposed mechanism for the enhancement of mineralocorticoid activity by licorice derivatives is the accumulation of aldosterone, deoxycorticosterone, and 11-deoxycorticosterone and their biologically active 5α-dehydro derivatives due to the inhibition of the 5β-reductase and 3β-HSD enzymes, as well as of glucocorticoids due to 11β-HSD inhibition (17).

Patients with apparent mineralocorticoid excess appear to be deficient in 11β-dehydrogenase but not 11-oxoreductase enzyme activity (30). While it has been assumed that 11β-HSD is an enzyme complex consisting of an 11β-dehydrogenase and a distinct 11-oxoreductase (9, 22, 30), a rat cDNA has been cloned and expressed as a single enzyme that interconverts cortisol/corticosterone to cortisone/11-dehydrocorticosterone (1). It has been reported that glycyrrhizic acid and carbenoxolone are not identical in their clinical activities and that glycyrrhizic acid inhibits the conversion of cortisol/corticosteron

to cortisone/11-dehydrocorticosterone unidirectionally, while carbenoxolone inhibits both the dehydrogenase and reductase directions (29). In our studies, the activity of glycyrrhizic acid and carbenoxolone were similar.

There is evidence for yet another mechanism of action of carbenoxolone. The MR is either missing or defective in patients with pseudohypoaldosteronism. Funder (10) has reported that the administration of carbenoxolone with a selective GR agonist in patients with pseudohypoaldosteronism and in adrenalectomized rats alters the function of the glucocorticoid, causing it to produce the same renal effects, Na^+ retention and K^+ excretion, as a mineralocorticoid would, presumably by causing GR-ligand complexes to act as activated MR. The animals in our experiments had intact adrenals; in fact, the mineralocorticoid effects of licorice depend on intact adrenal glands or replacement corticosteroids (9). Normally, most of the MR and many of the GR of the brain, depending on the area, are tonically bound by corticosterone, even in the unstressed rat (7, 26). While the concomitant icv infusion of corticosterone blocks icv aldosterone hypertension, the icv infusion of a selective glucocorticoid, presumably to the GR only, does not antagonize icv aldosterone hypertension. If there are two classes of MR in the brain, as has been postulated by De Kloet (7), carbenoxolone and glycyrrhizic acid may be altering the "corticosterone-preferring" MR to functionally "aldosterone-preferring" MR. If carbenoxolone were producing hypertension by "recruiting" GR and/or corticosterone-preferring MR bound to endogenous corticosterone to the pool of functionally activated MR, not only might the same cellular response be elicited as by activated MR in a mineralocorticoid-sensitive central blood pressure control area, but, more important, it might also remove the receptors that mediate the inhibition of icv aldosterone hypertension. This might explain why icv corticosterone, when given with carbenoxolone, neither increased the blood pressure, because the receptors were already surfeited, nor decreased it, because they were being diverted from their usual role of buffering the hypertensinogenic effect of aldosterone. The more rapid induction of hypertension by licorice compounds compared with aldosterone may be due more to the removal of local inhibitory effects than to the recruitment of more functional MR. The yin-yang relationship of the two classes of corticosteroids has been described elsewhere, including in the brain (7).

These data provide additional evidence for a central role in blood pressure control by mineralocorticoids that is distinct from their renal effects and that involves a complex homeostatic relationship between the two classes of corticosteroids in their central effects on blood pressure. They suggest that our understanding of functional specificity of the corticosteroid receptor-ligand complex, particularly in the brain, is incomplete. Finally, these studies indicate that more is involved in licorice-induced hypertension than the inhibition of 11β -HSD.

The expert technical help of Chris Fort is gratefully acknowledged. These studies were supported by Medical Research Funds from the Department of Veterans Affairs and a by National Heart, Lung, and Blood Institute Grant HL-27737.

Address for reprint requests: E. P. Gomez-Sanchez, Research Service

(151R), James A. Haley Veterans Hospital, 13000 Bruce B. Downs Blvd., Tampa, FL 33612.

Received 24 February 1992; accepted in final form 24 July 1992.

REFERENCES

1. Agarwal, A. K., C. Monder, B. Eckstein, and P. C. White. Cloning and expression of rat cDNA encoding corticosteroid 11β -dehydrogenase. *J. Biol. Chem.* 264: 18939-18943, 1989.
2. Armanini, D., I. Karbowiak, and J. W. Funder. Affinity of liquorice derivatives for mineralocorticoid and glucocorticoid receptors. *Clin. Endocrinol.* 19: 609-612, 1983.
3. Arriza, J. L., R. B. Simerly, L. W. Swanson, and R. M. Evans. The neuronal mineralocorticoid receptor as a mediator of glucocorticoid response. *Neuron* 1: 887-900, 1988.
4. Beaumont, K., and D. D. Fanestil. Characterization of rat brain aldosterone receptors reveals high affinity for corticosterone. *Endocrinology* 113: 2043-2051, 1983.
5. Bohr, D. F. What makes the pressure go up? A hypothesis. *Hypertension Dallas* 3; Suppl. II: 11-160-II-165, 1981.
6. Castello, R., R. Schwarting, C. Muller, K. Hierholzer, and I. Lichtenstein. Immunohistochemical localization of 11β -hydroxysteroid dehydrogenase in rat kidney with a monoclonal antibody. *Renal Physiol. Biochem.* 12: 320-327, 1989.
7. De Kloet, R. E. Brain corticosteroid receptor balance and homeostatic control. *Front. Neuroendocrinol.* 12: 95-184, 1991.
8. Doyle, D., R. Smith, S. Krozowski, and J. W. Funder. Mineralocorticoid specificity of renal type I receptors: binding and metabolism of corticosterone. *J. Steroid Biochem.* 33: 165-170, 1989.
9. Edwards, C. R. W., and P. M. Stewart. The cortisol-cortisone shuttle and the apparent specificity of glucocorticoid and mineralocorticoid receptors. *J. Steroid Biochem. Mol. Biol.* 39: 859-865, 1991.
10. Funder, J. W. How can aldosterone act as a mineralocorticoid? *Endocrinol. Res.* 15: 227-238, 1989.
11. Funder, J. W., P. T. Pearce, K. Myles, and L. P. Roy. Apparent mineralocorticoid excess, pseudohypoaldosteronism, and urinary electrolyte excretion: toward a redefinition of mineralocorticoid action. *FASEB J.* 4: 3234-3238, 1990.
12. Funder, J. W., P. T. Pearce, R. Smith, and A. I. Smith. Mineralocorticoid action: target tissue specificity is enzyme, not receptor, mediated. *Science Wash. DC* 242: 583-585, 1988.
13. Gomez-Sanchez, E. P. What is the role of the central nervous system in mineralocorticoid hypertension? *Am. J. Hypertens.* 4: 374-381, 1991.
14. Gomez-Sanchez, E. P., C. M. Fort, and C. E. Gomez-Sanchez. Intracerebroventricular infusions of RU 28318 blocks aldosterone-salt hypertension. *Am. J. Physiol.* 258 (Endocrinol. Metab. 21): E482-E484, 1990.
15. Gomez-Sanchez, E. P., M. T. Venkataraman, and D. Thwaites. Icv infusion of corticosterone antagonizes icv-aldosterone hypertension. *Am. J. Physiol.* 258 (Endocrinol. Metab. 21): E649-E653, 1990.
16. Krozowski, Z. S., and J. W. Funder. Renal mineralocorticoid receptors and hippocampal corticosterone binding species have identical intrinsic steroid specificity. *Proc. Natl. Acad. Sci. USA* 80: 6056-6060, 1983.
17. Latif, S. A., T. J. Conca, and D. J. Morris. The effects of the licorice derivative, glycyrrhetic acid, on hepatic 3α - and 3β -hydroxysteroid dehydrogenases and 5α - and 5β -reductase pathways of metabolism of aldosterone in male rats. *Steroids* 55: 52-58, 1990.
18. McEwen, B. S., L. T. Lambdin, T. C. Rainbow, and A. F. De Nicola. Aldosterone effects on salt appetite in adrenalectomized rats. *Neuroendocrinology* 43: 38-43, 1986.
19. Moisan, M. P., J. R. Seckl, and C. R. W. Edwards. 11β -Hydroxysteroid dehydrogenase bioactivity and messenger RNA expression in rat forebrain: localization in hypothalamus, hippocampus, and cortex. *Endocrinology* 127: 1450-1455, 1990.
20. Monder, C., and V. Lakshmi. Evidence for kinetically distinct forms of corticosteroid 11β -dehydrogenase in rat liver microsomes. *J. Steroid Biochem.* 32: 77-83, 1989.
21. Monder, C., and V. Lakshmi. Corticosteroid 11β -dehydrogenase of rat tissues: immunological studies. *Endocrinology* 126:

- 2435-2443, 1990.
22. Monder, C., P. M. Stewart, V. Lakshmi, R. Valentino, D. Burt, and C. R. W. Edwards. Licorice inhibits corticosteroid 11 β -dehydrogenase of rat kidney and liver: in vivo and in vitro studies. *Endocrinology* 89: 1046-1252, 1989.
 23. Morris, D. J., and G. W. Souness. The 11 β -OHSD inhibitor, carbenoxolone, enhances Na retention by aldosterone and 11-deoxycorticosterone. *Am. J. Physiol.* 258 (Renal Fluid Electrolyte Physiol. 27): F756-F759, 1990.
 24. Naray-Fejes-Toth, A., C. O. Watlington, and G. Fejes-Toth. 11 β -Hydroxysteroid dehydrogenase activity in the renal target cells of aldosterone. *Endocrinology* 129: 17-21, 1991.
 25. Porter, G. A., C. Rhodes, and P. Sacra. Comparative studies on the mineralo-corticoid action of aldosterone and carbenoxolone sodium in the adrenalectomized rat. *Pharmacology* 12: 224-229, 1974.
 26. Reul, J. M. H. M., F. R. Van Den Bosch, and E. R. De Kloet. Relative occupation of type-I and type-II corticosteroid receptors in rat brain following stress and dexamethasone treatment: functional implications. *J. Endocrinol.* 115: 459-467, 1987.
 27. Seckl, J. R., P. A. T. Kelly, and J. Sharkey. Glycyrrhetic acid, an inhibitor of 11 β -hydroxysteroid dehydrogenase, alters local cerebral glucose utilization in vivo. *J. Steroid Biochem. Mol. Biol.* 39: 777-779, 1991.
 28. Stewart, P. M., J. E. T. Corrie, C. H. L. Shackleton, and C. R. W. Edwards. Syndrome of apparent mineralocorticoid excess. *J. Clin. Invest.* 82: 340-349, 1988.
 29. Stewart, P. M., A. M. Wallace, S. M. Atherdon, C. H. Shearing, and C. R. Edwards. Mineralocorticoid activity of carbenoxolone: contrasting effects of carbenoxolone and liquorice on 11 β -hydroxysteroid dehydrogenase activity in man. *Clin. Sci. Lond.* 78: 49-54, 1990.
 30. Ulick, S., L. S. Levine, and P. Gunzler. A syndrome of apparent mineralocorticoid excess associated with defects in the peripheral metabolism of cortisol. *J. Clin. Endocrinol. Metab.* 49: 757-64, 1979.
 31. Ulick, S., R. Tedde, and F. Mantero. Pathogenesis of the type 2 variant of the syndrome of apparent mineralocorticoid excess. *J. Clin. Endocrinol. Metab.* 70: 200-206, 1990.
 32. Wrangé, O., and Z.-Y. Yu. Mineralocorticoid receptor in rat kidney and hippocampus: characterization and quantitation by isoelectric focusing. *Endocrinology* 113: 243-250, 1983.



d
F
ir
E
(
ir
st
tl
v
ti
it
p
tl
tr
tl
a
c
r
t
w
c
A
d
ti
tl
ir
p
di

P
li
se
ir
(
di
p
lo
af
p
a
is
di
fc

p
p
te
tl
w

Journal of

PHARMACY &

PHARMACOLOGY

Univ. of Min
Bio-Medica
Library

05 05 94

Volume 46 · Number 4 · April 1994

ISSN 0022-3573



J. Pharm. Pharmacol. 1994, 46: 305-309
Received August 27, 1993
Accepted November 15, 1993

© 1994 J. Pharm. Pharmacol.

A Novel 11β -Hydroxysteroid Dehydrogenase Inhibitor Contained in Saiboku-To, a Herbal Remedy for Steroid-dependent Bronchial Asthma

MASATO HOMMA, KITARO OKA, TOMOYUKI NIITSUMA* AND HISAO ITOH*

Department of Clinical Pharmacology, Tokyo College of Pharmacy, Horinouchi, Hachioji, Tokyo 192-03, Japan, and *Third Internal Medicine, Tokyo Medical College Hospital, Shinjuku-ku, Tokyo 160, Japan

Abstract—To identify the inhibitor of prednisolone metabolism contained in Saiboku-To, we conducted in-vitro experiments of 11β -hydroxysteroid dehydrogenase (11β -HSD), using rat liver homogenate and cortisol as a typical substrate. We studied the effects of ten herbal constituents on 11β -HSD. Five herbal extracts showed inhibitory activity with *Glycyrrhiza glabra* > *Perillae frutescens* > *Zizyphus vulgaris* > *Magnolia officinalis* > *Scutellaria baicalensis*. This suggests that unknown 11β -HSD inhibitors are contained in four herbs other than *G. glabra* which contains a known inhibitor, glycyrrhizin (and glycyrrhetic acid). Seven chemical constituents which have been identified as the major urinary products of Saiboku-To in healthy and asthmatic subjects were studied; magnolol derived from *M. officinalis* showed the most potent inhibition of the enzyme (IC_{50} , 1.8×10^{-4} M). Although this activity was less than that of glycyrrhizin, the inhibition mechanism (non-competitive) was different from a known competitive mechanism. These results suggest that magnolol might contribute to the inhibitory effects of Saiboku-To on prednisolone metabolism through inhibition of 11β -HSD.

Saiboku-To is the most popular anti-asthmatic Chinese herbal medicine (Kampo medicine in Japan) and has been used for corticosteroid-dependent asthma to obtain a steroid-sparing effect in prednisolone therapy (Nagano et al 1988). On the basis of animal experiments, the mechanism of action of Saiboku-To has been attributed to hormonal stimulation of the adrenal cortex (Hiai et al 1981; Shimizu et al 1984) and synergistic adjuvant effects on autacid secretions (Toda et al 1988) or allergic reactions (type I and IV) (Nishiyori et al 1983, 1985).

Recently, we proposed another mechanism which involves suppression of the systemic elimination of prednisolone (Taniguchi et al 1992). This pharmacokinetic effect seemed to result from 11β -hydroxysteroid dehydrogenase (11β -HSD) metabolic enzyme inhibition, because plasma prednisolone/prednisone ratios following Saiboku-To administration increased significantly (Taniguchi et al 1992). Since other Kampo-preparations containing *Glycyrrhiza glabra* did not show an effect on prednisolone pharmacokinetics (unpublished data), the effect of Saiboku-To could not be explained by known enzyme inhibitors such as glycyrrhizin and its aglycone glycyrrhetic acid, which are contained in *G. glabra*. These observations suggested that Saiboku-To must contain as yet uncharacterized 11β -HSD inhibitors.

In the present study, we carried out in-vitro experiments of 11β -HSD inhibition using cortisol and rat liver homogenate.

Materials and Methods

Materials

Saiboku-To (TJ-96, Tsumura Co., Tokyo, Japan) consists of fine brownish granules containing ten different herbal extracts (Table 1). Original herbs used for the assay were

purchased from Uchida Wakanyaku Co. (Tokyo, Japan). The extracts of Saiboku-To and of original herbs were prepared as follows. One gram Saiboku-To or the crushed herb in 15 mL 35% ethanol was gently refluxed for 1 h on a steam bath. After cooling to room temperature, water was added to make a total volume of 10 mL before centrifugation at 1600 g for 10 min. The resulting supernatant was used for the assay.

Glycyrrhizin, glycyrrhetic acid, wogonin, and baicalin were purchased from Wako Pure Chemicals (Osaka, Japan). Magnolol and honokiol were donated by Professor Y. Sashida of Tokyo College of Pharmacy (Fujita et al 1973). Medicarpin and oroxylin A were kindly contributed by Professor T. Nomura of Toho University School of Pharmacy (Tokyo, Japan) and Tsumura Co., respectively. 8,9-Dihydroxydihydromagnolol was prepared by us from magnolol by osmic acid oxidation (Homma et al 1992). Liquiritigenin was isolated from *G. glabra* according to Shibata & Saitoh (1978). Chemical structures of these compounds are given in Fig. 1. Cortisol and cortisone were purchased from Sigma Chemical Co. (St Louis, MO, USA). Other organic and inorganic reagents were of analytical grade.

Rat liver homogenates were prepared in the usual manner: fresh liver was isolated from a male Wistar rat (freely fed, body weight 250 g) and was cut into small pieces. The pieces were homogenized in 10 vol 0.25 M sucrose in a glass-homogenizer with a Teflon piston. The homogenates were frozen at -80°C and stored until incubation.

Instruments

Our HPLC system for determination of glucocorticoids in incubation mixtures consisted of a solvent delivery pump (VIP-I, Jasco, Tokyo), a UV-detector (Uvidex-100-III, Jasco), a single pen recorder (Pantos U-228, Nippon Denshi, Tokyo), a sample injector with a loop volume of 100 μL .

Correspondence: M. Homma, Department of Clinical Pharmacology, Tokyo College of Pharmacy, Horinouchi, Hachioji, Tokyo 192-03, Japan.

Table 1. Herbal composition of Saiboku-To.

| Constituent herb | Family | Composition (% w/w) |
|---|---------------|---------------------|
| <i>Bupleurum falcatum</i> L. | Umbelliferae | 20.6 |
| <i>Pinellia ternata</i> Beitenbach | Araceae | 14.7 |
| <i>Portia cocos</i> Woll. | Polyporaceae | 14.7 |
| <i>Scutellaria baicalensis</i> Georgi | Labiatae | 8.8 |
| <i>Zizyphus vulgaris</i> Lam. | Rhamnaceae | 8.8 |
| <i>Panax ginseng</i> C. A. Meyer | Araliaceae | 8.8 |
| <i>Magnolia officinalis</i> | Magnoliaceae | 8.8 |
| <i>Glycyrrhiza glabra</i> L. | Leguminosae | 5.9 |
| <i>Perillae frutescens</i> Britton var. <i>acuta</i> Kudo | Labiatae | 5.9 |
| <i>Zingiber officinale</i> Roscoe | Zingiberaceae | 3.0 |

(Model 7125, Rheodyne, CA, USA), and a silica gel column (LiChrosorb Si-60, 5 μ m, i.d. 4 mm \times 250 mm, Merck, Darmstadt, Germany). The mobile phase was a mixture of water/methanol/dichloromethane/n-hexane (0.1/8.0/30.0/61.9 v/v) with a flow rate of 1.5 mL min⁻¹. Detector sensitivity was set at 0.005–0.01 a.u. at 245 nm. We used a disposable syringe minicolumn (Extrasht, Kusano Sci. Co., Tokyo) to perform sample injections (Homma et al 1989; Kouno et al 1990).

Determination of 11 β -HSD inhibition activity

We measured 11 β -HSD activity in rat liver homogenate incubation mixtures, detecting chemical transformation of cortisol to cortisone in the presence of 11 β -HSD inhibitors. Oxidation at the C-11 position of the steroid nucleus was kinetically characterized by measuring the conversion rate of cortisol to cortisone in the presence of NADP⁺ in rat liver homogenate according to the procedure of Monder et al (1989) with minor modification. The incubation mixtures

consisted of 620 μ L 0.1 M Tris-HCl buffer (pH 8.5) containing 0.014% Triton-X, 50 μ L 5 mM NADP⁺, 100 μ L rat liver homogenate, and 200 μ L aqueous solution for Saiboku-To and original herbal extracts or 200 μ L buffer solution for each chemical such as the known inhibitors (glycyrrhizin and glycyrrhetic acid) and our candidates isolated from urine of subjects receiving the preparation. These chemicals were dissolved in a buffer solution directly or after pre-solubilization in a small amount of ethanol with a final concentration in incubation mixtures of less than 2%. After 10 min pre-incubation at 37°C, 200 μ L 0.3 mM cortisol was added and the resulting mixtures were further incubated for 10 min. The enzyme reaction was terminated by an addition of 100 μ L 5% sulphuric acid. Cortisol and cortisone in the mixtures were determined by HPLC using Extrasht as described in our previous papers (Homma et al 1989; Kouno et al 1990). Briefly, 5 μ L incubation mixture and 2 μ L sodium hydroxide solution were loaded onto Extrasht which was then attached to the sample-loading injector of the HPLC system.




FIG. 1. Chemical structures of test compounds. * These compounds have not been detected in urine following Saiboku-To administration.

Ethanol into the syringe. in the i The rec mixture less than control of the u

Effects. Effects: cortisol of Saib reaction and five the inhi (87.5% vulgaris (19.1%

Effects Seven

Ta hei in

Sa B. P. S. Z. P. M. G. P. Z.

exi trr

Table 3. nary me

Inhibito Urinary Magr Dihy Wogr Baica Orox Liqui Medi

Known Glycy Glyc

Ethanol/dichloromethane (2/98 v/v, 130 μ L) was injected into the system through Extrasbot using a tuberculin glass syringe. Thus, extraction and injection of the glucocorticoids in the incubation mixtures were achieved simultaneously. The recovery rates of glucocorticoids from the incubation mixture were more than 95% with coefficient of variations less than 5%. Direct peak-height calibration of the test and control mixtures afforded inhibitory activity (% inhibition) of the test materials against 11β -HSD.

Results

Effects of herbal extracts

Effects of original herbal extracts on conversion of cortisol to cortisone by rat liver homogenate were compared with that of Saiboku-To (Table 2). Cortisone production in the reaction mixture was significantly inhibited by Saiboku-To and five original herbal extracts ($P < 0.05$). The magnitude of the inhibition (% inhibition) was in the order Saiboku-To (87.5%) > *G. glabra* (80.8%) > *P. frutescens* (30.9%) > *Z. vulgaris* (27.6%) > *M. officinalis* (19.8%) > *S. baicalensis* (19.1%).

Effects of urinary metabolites of Saiboku-To

Seven candidates (Fig. 1) were tested with respect to the

Table 2. Effects of Saiboku-To and its constituent herbal extracts on 11β -hydroxysteroid dehydrogenase in rat liver homogenate.

| | % inhibition* | % activity of Saiboku-To |
|-----------------------|------------------|--------------------------|
| Saiboku-To | 87.5 \pm 3.4** | 100.0 |
| <i>B. falcatum</i> | 7.7 \pm 5.7 | 8.8 |
| <i>P. ternata</i> | 5.8 \pm 4.2 | 6.6 |
| <i>P. cocos</i> | — | — |
| <i>S. baicalensis</i> | 19.1 \pm 11.5* | 21.8 |
| <i>Z. vulgaris</i> | 27.6 \pm 4.0** | 31.5 |
| <i>P. ginseng</i> | 10.9 \pm 6.9 | 12.5 |
| <i>M. officinalis</i> | 19.8 \pm 3.7** | 22.6 |
| <i>G. glabra</i> | 80.8 \pm 1.0** | 92.3 |
| <i>P. frutescens</i> | 30.9 \pm 9.6** | 35.3 |
| <i>Z. officinale</i> | 12.8 \pm 8.7 | 14.6 |

* Data are presented as mean \pm s.d. of triplicate experiments. ** $P < 0.05$, *** $P < 0.01$ compared with control.

Table 3. Inhibition of 11β -hydroxysteroid dehydrogenase by urinary metabolites of Saiboku-To and known inhibitors.

| Inhibitor | Inhibition (%) | |
|-----------------------------------|----------------|----------------|
| | 10 μ M | 100 μ M |
| Urinary metabolites of Saiboku-To | | |
| Magnolol | 15.1 \pm 4.4 | 43.9 \pm 3.0 |
| Dihydroxydihydromagnolol | — | — |
| Wogonin | — | 7.4 \pm 0.8 |
| Baicalin | 6.8 \pm 1.6 | 14.8 \pm 1.6 |
| Oroxylin A | — | 5.1 \pm 5.5 |
| Liquiritigenin | — | — |
| Medicarpin | — | 12.2 \pm 3.3 |
| Known inhibitors | | |
| Glycyrrhizin | 81.1 \pm 5.4 | 97.3 \pm 1.1 |
| Glycyrrhetic acid | 100.0 | — |




Fig. 2. Chromatographic comparison of the effect of magnolol (100 μ M) on transformation of cortisol (peak 1) to cortisone (peak 2) by 11β -hydroxysteroid dehydrogenase. A. Before incubation with magnolol; B. after incubation without magnolol; C. after incubation with magnolol.




Fig. 3. Dose-dependent inhibitory effects of magnolol (●), honokiol (○), glycyrrhizin (▲), and glycyrrhetic acid (Δ) on 11β -hydroxysteroid dehydrogenase. Data are presented as mean \pm s.d. of triplicate experiments.

effects on rat liver 11β -HSD at concentrations of 10 and 100 μ M. The results were compared with those of the known inhibitors, glycyrrhizin and glycyrrhetic acid (Table 3). Five of seven candidates showed inhibitory activity at 100 μ M, although their activities were weaker than those of the known inhibitors. Dihydroxydihydromagnolol in *M. officinalis* and liquiritigenin in *G. glabra* did not show any activity at the test concentrations. Wogonin, baicalin, and oroxylin A (flavonoids derived from *S. baicalensis*), and medicarpin (a




Fig. 4. Lineweaver-Burk double reciprocal plots of initial enzyme velocity and concentration of cortisol in the presence of magnolol at concentrations of 0 (O), 0.1 (●), 0.15 (Δ), and 0.2 (▲) mM.

homoisoflavonoid in *G. glabra*) showed weak activity. However, considerable inhibition was observed with magnolol, a neolignan derived from *M. officinalis*. A typical chromatogram for determination of the inhibitory activity of magnolol is shown in Fig. 2, where the chemical transformation from cortisol to cortisone was clearly suppressed. The dose-dependent inhibitory effect of magnolol is compared with those of glycyrrhizin and glycyrrhetic acid in Fig. 3. The IC₅₀ values of magnolol, glycyrrhizin, and glycyrrhetic acid were 1.8×10^{-4} , 2.6×10^{-6} , and 9.0×10^{-3} M, respectively. Since *M. officinalis* contains another congener of magnolol, honokiol (not a urinary metabolite), we also examined the effect of honokiol on 11 β -HSD and found a dose-dependent inhibitory effect with IC₅₀ of 7.0×10^{-3} M (Fig. 3).

Mechanism of magnolol in 11 β -HSD inhibition

Fig. 4 shows the inhibitory effects of magnolol on rat liver 11 β -HSD. The data were plotted according to the Lineweaver-Burk linear transformation of the Michaelis-Menten equation. The double reciprocal plots on Fig. 4 suggested magnolol has a unique non-competitive inhibitory mechanism. We were unable to estimate an inhibition constant (K_i) of magnolol by the Dixon plot because of this non-competitive inhibition.

Discussion

This paper suggests the presence of several novel inhibitors of 11 β -HSD in five constituent herbs. *G. glabra*, *P. frutescens*, *Z. vulgaris*, *M. officinalis* and *S. baicalensis*. Although these inhibitors seem to contribute to in-vitro activity of Saiboku-To, their contributions to prednisolone metabolism during clinical Saiboku-To treatment has been unclear. However, we emphasize the importance of this possibility, since our biologically active compounds in herbal medicine are found in biofluids following administration (Homma et al 1992, 1993a).

In our previous study, we found seven phenolic compounds in urine after oral administration of Saiboku-To (Homma et al 1992, 1993a, b). These compounds seemed to be possible candidates which explain in-vivo effects of Saiboku-To. Five of these compounds showed inhibitory

activity against 11 β -HSD in-vitro (Table 3). The intensities of those activities were almost equal to those of the corresponding herbal extracts, except that *G. glabra*, containing glycyrrhizin, concealed the effects of liquiritigenin and medicarpin. Magnolol exhibited activity at concentrations higher than 1×10^{-3} M (Fig. 3). Similar activity was also observed in honokiol, a hydroxylated derivative of magnolol isolated from *M. officinalis* but not found as a urinary metabolite of Saiboku-To.

The novel 11 β -HSD inhibitors found in this study belong to a class of phenolic compounds, lignans and flavonoids, whose chemical structures are completely different from those of the previously described inhibitors. Unexpectedly, the inhibition mechanism of magnolol seems to be different from those of the known inhibitors, the latter exhibiting competitive inhibition (Monder et al 1989). Although 11 β -HSD inhibitors have been considered so far to belong to a limited class of liquorice triterpenoids, the present results suggested that the naturally occurring lignans and flavonoids also possess inhibitory activity through a different mechanism.

Urinary non-conjugated magnolol in responders to Saiboku-To is significantly higher than that in the non-responders (Homma et al 1993a, b). This suggests that magnolol is an important chemical constituent for the clinical effects of Saiboku-To, playing an important role for alteration of prednisolone pharmacokinetics.

The inhibitory effects of liquorice glycosides on 11 β -HSD are so marked in animal experiments in-vivo and in-vitro (Monder et al 1989; Mackenzie et al 1990), that Saiboku-To could inhibit 11 β -HSD even though the glycyrrhizin content is relatively small. However, the effect of Saiboku-To cannot be explained by glycyrrhizin alone, because another Kampo preparation, Sho-Saiko-To which contains *G. glabra* but not *P. cocos*, *M. officinalis* or *P. frutescens*, did not affect prednisolone pharmacokinetics in healthy subjects (unpublished data). Animal experiments using pure compounds will be needed to clarify the role of lignans and flavonoids on prednisolone metabolism.

Acknowledgements

This work was supported by the Ministry of Education in Japan (Grant-in-Aid for Scientific Research 03857345). Dr Y. Sashida and T. Nomura are gratefully acknowledged for kindly providing magnolol and honokiol and medicarpin. We thank Miss E. Yoshida and Mr H. Tamura for their technical assistance.

References

- Fujita, M., Itokawa, H., Sashida, Y. (1973) Study on the components of *Magnolia obovata* Thunb. II. On the components of the methanol extract of the bark. *Yakugaku Zasshi* 93: 422-428 (in Japanese)
- Hiai, S., Yokoyama, H., Nagasawa, T., Oura, H. (1981) Stimulation of the pituitary-adrenocortical axis by saikosaponin of *Bupleurum* radix. *Chem. Pharm. Bull. (Tokyo)* 29: 495-499
- Homma, M., Oka, K., Takahashi, N. (1989) Liquid chromatographic determination of theophylline concentration with syringe-type microcolumns for direct plasma injection. *Anal. Chem.* 61: 784-787
- Homma, M., Oka, K., Yamada, T., Niitsuma, T., Itoh, H., Takahashi, N. (1992) A strategy for discovering biologically active compounds with high probability in traditional Chinese

herb n
Anal.
Homma.
netic e
1995
Homma.
Itoh, J
tions I
Chine
Kouno,
(1990)
chron
togr. I
Macke
Benra
glycy
young
Monder.
Edwa
dehyd
Endo
Nagano.
study

- herb remedies: an application of Saiboku-To in bronchial asthma. *Anal. Biochem.* 202: 179-187
- Homma, M., Oka, K., Niitsuma, T., Itoh, H. (1993a) Pharmacokinetic evaluation of traditional Chinese herb remedies. *Lancet* 341: 1595
- Homma, M., Oka, K., Kobayashi, H., Niitsuma, T., Yamamoto, S., Itoh, H., Takahashi, N. (1993b) Impact of free magnolol excretions in asthmatic patients who responded well to Saiboku-To, a Chinese herbal medicine. *J. Pharm. Pharmacol.* 45: 844-846
- Kouno, Y., Ishikura, C., Takahashi, N., Homma, M., Oka, K. (1990) Direct sample injection into the high-performance liquid chromatographic column in theophylline monitoring. *J. Chromatogr.* 515: 321-327
- Mackenzie, M. A., Hoetnagels, W. H. L., Jansen, R. W. M. M., Benrad, T. J., Kloppenborg, P. W. C. (1990) The influence of glycyrrhetic acid on plasma cortisol and cortisone in healthy young volunteers. *J. Clin. Endocrinol. Metab.* 70: 1637-1643
- Monder, C., Stewart, P. M., Lakshmi, V., Valentino, R., Burt, D., Edwards, C. R. W. (1989) Licorice inhibits corticosteroid 11β -dehydrogenase of rat kidney and liver: in vivo and in vitro studies. *Endocrinology* 125: 1046-1053
- Naganó, J., Kobayashi, S., Nakajima, S., Egashira, Y. (1988) A study of the long-term effects of Saiboku-To in bronchial asthma—evaluation in the field of pediatrics by multi-center open trial. *Respiration* 7: 76-87 (in Japanese)
- Nishiyori, T., Nakatomi, L., Matsuura, N., Nagai, H., Koda, A. (1983) Effect of Chinese blended medicine, Saiboku-To, on Type IV allergic reaction. *Jpn. J. Allergol.* 32: 317-323
- Nishiyori, T., Tsuchiya, H., Inagaki, N., Koda, A. (1985) Effects of Chinese blended preparation, Saiboku-To, in type I allergic reaction and experimental atopic asthma in particular. *Folia Pharmacol. Japon.* 85: 7-16 (in Japanese)
- Shibata, S., Saitoh, T. (1978) Flavonoid compounds in licorice root. *J. Ind. Chem. Soc.* 55: 1184-1191
- Shimizu, K., Amagaya, S., Ogihara, Y. (1984) Combination effects of Sho-Saiko-To (Chinese traditional medicine) and prednisolone on the anti-inflammatory action. *J. Pharmacobiodyn.* 7: 891-899
- Taniguchi, C., Homma, M., Oka, K., Kobayashi, H., Takahashi, N., Yamamoto, S., Itoh, H. (1992) Effects of Saiboku-To on prednisolone metabolism. *Jpn. J. Ther. Drug Monit.* 9: 18-24 (in Japanese)
- Toda, S., Kimura, M., Ohnishi, M., Nakanishi, K. (1988) Effects of the Chinese herbal medicine "Saiboku-To" on histamine release from and the degranulation of mouse peritoneal mast cells induced by compound 48/80. *J. Ethnopharmacol.* 24: 303-309

intensities
of the
ra, con-
itigenin
centra-
was also
agnolol
urinary

belong
onoids,
at from
ectedly,
different
hibiting
gh 11β -
ong to a
t results
onoids
nechan-

to Sai-
re non-
sts that
for the
role for

11β -HSD
in-vitro
oku-To
content
cannot
Kampo
but not
ot affect
(unpub-
nds will
oids on

ation in
345). Dr
dged for
licarpin.
for their

e compo-
nts of the
2-428 (in

imulation
Bupreul

hromato-
h syringe-
chem. 61:

Itoh, H.,
ologically
Chinese

CURRENT
AWARENESS
TITLE

JANUARY 1996

CLINICAL PHARMACOLOGY & THERAPEUTICS

OFFICIAL PUBLICATION OF THE
American Society for Clinical Pharmacology and Therapeutics
American Society for Pharmacology and Experimental Therapeutics

VOLUME 59 NUMBER 1

MARCUS M. REIDENBERG, MD *Editor*

APPEARING IN THIS ISSUE

■ COMMENTARY

Hematopoietic growth factors with cytotoxic drugs

■ PHARMACOKINETICS

Rifampin-midazolam interaction

Zidovudine-atovaquone interaction

Netivudine metabolite inhibits dihydropyrimidine dehydrogenase

Vinorelbine in patients with liver metastases

Intestinal CYP3A in celiac disease

Dapsone or erythromycin as probes for CYP3A

Oxycodone disposition and effects

■ PHARMACODYNAMICS

Grapefruit juice inhibits 11 β -hydroxysteroid dehydrogenase

Losartan and enalapril effects on angiotensin responses

■ PHARMACOEPIDEMIOLOGY

Sleep symptoms and sedative hypnotic use in the elderly

ASCPT meeting program

Univ. of Minn.
Bio-Medical
Library

M Mosby
ISSN 0009-9236

2 5 96

PHARMACODYNAMICS AND DRUG ACTION

Grapefruit juice and its flavonoids inhibit 11 β -hydroxysteroid dehydrogenase

Introduction: The enzyme 11 β -hydroxysteroid dehydrogenase (11 β -OHSD) oxidizes cortisol to inactive cortisone. Its congenital absence or inhibition by licorice increases cortisol levels at the mineralocorticoid receptor, causing mineralocorticoid effects. We tested the hypothesis that flavonoids found in grapefruit juice inhibit this enzyme in vitro and that grapefruit juice itself inhibits it in vivo.

Methods: Microsomes from guinea pig kidney cortex were incubated with cortisol and nicotinamide adenine dinucleotide (NAD) or nicotinamide adenine dinucleotide phosphate (NADP) and different flavonoids and the oxidation to cortisone measured with use of HPLC analysis. In addition, healthy human volunteers drank grapefruit juice, and the ratio of cortisone to cortisol in their urine was measured by HPLC and used as an index of endogenous enzyme activity.

Results: Both forms of 11 β -OHSD requiring either NAD or NADP were inhibited in a concentration-dependent manner by the flavonoids in grapefruit juice. Normal men who drank grapefruit juice had a fall in their urinary cortisone/cortisol ratio, suggesting in vivo inhibition of the enzyme.

Conclusion: Dietary flavonoids can inhibit this enzyme and, at high doses, may cause an apparent mineralocorticoid effect. (CLIN PHARMACOL THER 1996;59:62-71.)

Yil Seob Lee, MD,* Beverly J. Lorenzo, BS, Theo Koufis, MS, and
Marcus M. Reidenberg, MD New York, N.Y.

The enzyme 11 β -hydroxysteroid dehydrogenase (11 β -OHSD) oxidizes cortisol to inactive cortisone. This enzyme in the kidney regulates the amount of mineralocorticoid activity there, because cortisol binds as avidly to the mineralocor-

ticoid receptor as aldosterone does. Deficiency of this enzyme in children, first described by Ulick et al.¹ in 1977, causes high cortisol levels in the kidney that result in hypertension and hypokalemia. Licorice-induced hypermineralocorticoidism is probably due to the inhibition of 11 β -OHSD by glycyrrhizic acid, the active principle of licorice.^{2,4} Much research has been done since 1977 on syndromes of apparent mineralocorticoid excess.^{5,6}


Gossypol, a polyphenolic constituent of cotton seed, has been studied in China as a potential male oral contraceptive, but hypokalemia developed in some Chinese men while they were taking it.⁷ We found that gossypol inhibited 11 β -OHSD activity in guinea pig⁸ and human renal cortical microsomes.⁹ Because there are structural similarities between gossypol and some flavonoids, we tested some of these and some other compounds, such as diuretics, that cause hypokalemia⁹⁻¹⁰ and discovered that some inhibit this enzyme. Narin-

From the Departments of Pharmacology and Medicine, Division of Clinical Pharmacology, Cornell University Medical College. Supported by grant RR47 from the National Institutes of Health (Bethesda, Md.) and by grants from Hoffmann-La Roche Inc. (Nutley, N.J.), Sandoz Pharmaceuticals Inc. (East Hanover, N.J.), The Rockefeller Foundation (New York, N.Y.), and Han-Dok Remedia (Seoul, Korea).

Received for publication May 11, 1995; accepted Aug. 17, 1995. Reprint requests: Marcus M. Reidenberg, MD, Department of Pharmacology, Cornell University Medical College, 1300 York Ave., New York, NY 10021.

*Present address: Han-Dok Remedia Ind. Co., Ltd., 735 Yoksam-I-Dong, Kangnan-Ku, SL Young Dong, PO Box 1560, Seoul, Korea.

Copyright © 1996 by Mosby-Year Book, Inc.
0009-9236/96/35.00 + 0 13/1/68639



genin, the aglycone of naringin, is a major flavonoid in grapefruit juice and inhibits this enzyme.¹⁰ Recent work suggests that there are two isoforms of this enzyme, nicotinamide adenine dinucleotide (NAD)-dependent 11 β -OHSD and nicotinamide adenine dinucleotide phosphate (NADP)-dependent 11 β -OHSD with specific tissue distributions.¹¹⁻¹⁴ The effects of these flavonoids are worth study because about 25 mg of flavonoids has been recently estimated to be ingested daily in the diet,¹⁵ whereas older studies cite as much as 1 gm per day.¹⁶

The objective of this study was to learn which other flavonoids in grapefruit juice inhibit 11 β -OHSD in vitro and whether grapefruit juice inhibits the enzyme in vivo.

MATERIAL AND METHODS

In vitro study

Chemicals and solutions. All flavonoids (see Structures), cortisone, cortisol, corticosterone, NAD, NADP, 99.9% dimethyl sulfoxide (DMSO), and Sigma Diagnostic Total Protein Kit (cat. No. 690-A) were purchased from Sigma Chemical Co., St. Louis, Mo. All flavonoids were dissolved in DMSO. Cortisone, cortisol, and corticosterone were dis-

solved in methanol (J.T. Baker HPLC grade purchased from VWR Scientific, Piscataway, N.J.) (1.4 mmol/L) and kept at -4° C. NAD and NADP (5 mmol/L) were dissolved in Tris hydrochloric acid buffer (0.1 mol/L, pH 8.0).

Microsomal preparation and assay of 11 β -OHSD activity. Guinea pig kidney cortex was obtained from long-haired male Hartley guinea pigs (Hilltop, Pa.). Tissue was homogenized by a Tekmar Tissuemizer (Cincinnati, Ohio). Microsomes were prepared, diluted to a concentration of 1.25 mg protein/ml as measured by the Sigma Diagnostics Total Protein Kit, and stored at -70° C. The enzyme activity in the microsomes was determined by measuring the rate of conversion of cortisol to cortisone in the presence of NAD or NADP as described previously.⁶⁻¹⁰ Each flavonoid was studied with use of NAD and NADP as the cofactor. The conversion rates from cortisol to cortisone were determined, and the extent of inhibition was calculated. The drug concentrations that inhibited the enzyme by 50% (IC₅₀) were estimated from duplicate incubations at each concentration of at least three different concentrations of each flavonoid by use of the dose-response program of Chou and Chou (*Dose-effect Analysis with Microcomputers*, Elsevier-Biosoft, Cambridge, En-

gland, 1989). For each flavonoid studied, at least one concentration was above and one below the IC_{50} .

Analytical method for urinary cortisone and cortisol

We modified our HPLC method for measurement of these compounds from microsomal incubation mixtures.⁹ The equipment consisted of a Waters Automated Gradient Controller with two Waters 6000A pumps (Waters Chromatography, Milford, Mass.). The injector was a Waters U6K and the detector was a Waters 486 Tunable Absorbance Detector set at a wavelength of 246 nm and 0.15 absorbance units full scale. The separation was performed with a Waters Nova-Pak C₁₈ 3.9 × 150 mm stainless steel column (4 μm spherical particle size, pore size 60 Å, 7% carbon load, end-capped) or with a Waters μBondapak C₁₈ 3.9 × 300 mm column (10 μm irregular particle size, pore size 125 Å, 10% carbon load, end-capped). The peak areas were recorded on a SE120 plotter purchased through Waters Chromatography.

The mobile phase was methanol/water, initially at 70% water:30% methanol. Conditions were changed over the first 6 minutes to 56% water:44% methanol in a linear gradient that was then held isocratically for 14 minutes. The gradient was then reversed linearly to 70:30 over 3 minutes and the column equilibrated for 5 minutes before the next injection. The flow rate was 1 ml/min.

Procedure

To each 10 ml aliquot of every standard and sample (performed in duplicate) was added 40 μl of the 25 μg/ml corticosterone* as the internal standard. The samples were briefly vortexed to mix. One milliliter of 0.1 mol/L of sodium hydroxide was added to each test tube and again briefly vortexed to mix. Three milliliters of methylene chloride were added to each sample, capped with Teflon-lined screw tops, and rotated for 45 minutes on a mechanical rotator at approximately 20 rpm. The samples were centrifuged at 3000 rpm (1000g) for 15 minutes. The aqueous layer (top) was aspirated to waste. Again the samples were centrifuged for 10 minutes at 3000 rpm and the remainder of the aque-

ous phase was aspirated. A small spatula full of sodium chloride (~150 mg) was added to each sample, and any emulsion was broken up with a Pasteur pipet. The samples were then again centrifuged for 10 minutes. The organic layer was carefully transferred to clean test tubes and evaporated to dryness in a warm water bath (~45° C) under a stream of nitrogen. The residue was redissolved in 150 μl of HPLC grade methanol and injected into the HPLC.

The retention times were 16.5, 19.0, and 23.5 minutes for cortisone, cortisol, and corticosterone, respectively, on a Waters 10 micron, 300 × 3.9 mm stainless steel μBondapak C₁₈ column. On a Waters 4 micron, 150 × 3.9 mm Nova-Pak, the retention times for cortisone, cortisol, and corticosterone were 12.8, 13.6, and 17.8 minutes. Levels measured in about 60 human urine samples ranged from 7.1 to 215.4 ng/ml for cortisone and 4.5 to 230.1 ng/ml for cortisol. The ratio of cortisone to cortisol was 0.2 to 5.7.

The absolute recovery was 70% for cortisol and 69% for cortisone. The interday coefficient of variation for cortisone was 6.5% for 25 ng/ml and 1.5% for the 100 ng/ml standard. For cortisol, the values were 6.3% for 25 ng/ml and 1.1% for 100 ng/ml. Cortisone dissolved in methanol was chromatographed and the peak was collected. The putative cortisone peak from extracted urine was also collected, and both fractions were scanned with a Varian Cary 219 spectrophotometer. The peaks had identical absorption spectra, with absorption maximums at 239 nm. (The *CRC Handbook of Chemistry and Physics* states that the absorption maximum of cortisone in alcohol is 237 nm).

All samples were assayed twice in duplicate. Standard curves for cortisone and cortisol were determined and plotted as in the *in vitro* study. Concentrations of these steroids in unknown samples were extrapolated from these standard curves.¹⁰

In vivo preliminary study

Six male volunteers aged from 35 to 65 years (two investigators and four other members of the Department of Pharmacology) who were living at home gave daily morning urine samples for 4 days. They then drank grapefruit juice, requested to be at a dose of a quart a day, for 7 days, and gave daily morning urine samples on the last 4 days of this period. After a 3-day washout period, the subjects again gave daily morning urine samples for 4 days.

*Corticosterone is excreted by humans at a rate that averages 6 μg/24 hours¹⁷ or less than of 1% of 1.5 to 4.0 mg/24 hour production rate.¹⁸ Thus the concentration from endogenous sources is less than 10% of that added, a negligible amount for this assay.


**EFFECT OF GRAPEFRUIT JUICE ON
URINARY CORTISONE TO CORTISOL RATIO**

Fig. 1. Preliminary study results in six subjects living at home. Subjects 1 and 6 were two of the authors, who are known to have consumed the full amount of grapefruit juice.

The cortisone and cortisol concentrations were measured in each urine sample. The two investigators (subjects 1 and 6) had a decrease in the ratio of urinary cortisone to cortisol during the grapefruit juice period compared with the control periods before and after grapefruit juice (mean \pm SD for

subject 1 was 11.4 ± 3.1 , 5.7 ± 0.9 , and 10.2 ± 2.1 ; mean \pm SD for subject 6 was 4.8 ± 0.4 , 3.6 ± 0.6 , and 4.7 ± 0.8). The other four subjects had no significant change. All data are shown in Fig. 1. Subjects 1 and 6 then volunteered for the dose-response study.




Fig. 2. Concentration-response relationships for the inhibition of 11 β -hydroxysteroid dehydrogenase by different flavonoids with use of nicotinamide adenine dinucleotide (NAD; solid bars) or nicotinamide adenine dinucleotide phosphate (NADP; open bars) as a cofactor. * $p < 0.05$.

In vivo dose-response study

These two volunteers (subjects 1 and 6) gave urine samples for the last 4 days of four 7-day study periods. (1) First control period: Each subject collected a 10-hour (7 AM to 5 PM) urine sample daily for 4 days (one subject missed 1 day of sample collection). (2) Low-dose period: Each subject drank 950 to 1060 ml grapefruit juice a day for 7 days and gave 10-hour urine samples for the last 4 days of the 7-day period. (3) High-dose period: Each subject drank 1900 to 2100 ml (double volume of low-dose period) grape-

fruit juice for 7 days and gave daily 10-hour urine samples for the last 4 days. (4) Second control period: Each subject gave daily 10-hour urine samples for 4 days after 3 days of a washout period.

In vivo metabolic balance study

Protocol. Two different healthy male volunteers (aged 26 and 31 years), not previously screened for responsiveness to grapefruit juice, gave informed written consent and were admitted to the clinical research center for 3 weeks. An evaluation before the study


**EFFECT OF DIFFERENT DOSES OF GRAPEFRUIT
JUICE ON URINARY CORTISONE TO CORTISOL RATIO**

Fig. 3. Urinary cortisone/cortisol ratios in subjects in dose-response study. Each period include four daily urine collections. Each urine sample was assayed twice, each assay in duplicate. Each point represents a single assay (four points per daily urine).

showed normal physical findings, serum chemistry, hematology, and ECG for both subjects. They ate a diet that had a constant amount of sodium and potassium during the study (potatoes, bananas, and lemonade during control periods to balance the grapefruit juice during the experimental period). Their blood pressures and body weights were measured daily. Twenty-four-hour urine was collected for free cortisone, cortisol, Na^+ , and K^+ for the last 4 days of three 7-day study periods. Blood samples were drawn for Na^+ and K^+ for the same periods. Plasma renin activity, aldosterone and cortisol, and urinary aldosterone excretion were measured at the end of each period. The first and third weeks were the control periods. The second week was the experimental period in which 1500 ml grapefruit juice (100% from concentrate, Ocean Spray Cranberries Inc., Lakeville, Mass.) was consumed daily.

Statistics. The Bonferroni *t* test after a one-way ANOVA was used to assess statistically significant dif-

Table I. Inhibition of 11 β -OHSD in microsomes of guinea pig kidney by various flavonoids in the presence of NAD or NADP

| Flavonoids | IC ₅₀ ($\mu\text{mol/L}$) | |
|-------------|--|--------------------|
| | NAD | NADP |
| Quercetin* | 192 \pm 18 | 355 \pm 82 |
| Apigenin* | 284 \pm 25 | 125 \pm 16 |
| Kaempferol | 322 \pm 13 | 293 \pm 62 |
| Naringenin* | 496 \pm 77 | 264 \pm 63 |
| Hesperetin* | 769 \pm 69 | 509 \pm 45 |
| Naringin* | 21,191 \pm 4,949 | 10,550 \pm 1,136 |
| Hesperidin | >55,000 | >50,000 |

Data are mean values \pm SD.
11 β -OHSD, 11 β -Hydroxysteroid dehydrogenase; NAD, nicotinamide adenine dinucleotide; NADP, nicotinamide adenine dinucleotide phosphate; IC₅₀, drug concentration that inhibited the enzyme by 50%.
**p* < 0.05; NAD compared with NADP.

ferences. Statistical significance was assumed when the corresponding *p* values were lower than $\alpha = 0.05$.

Approval. All human studies were approved by the Cornell Institutional Review Board.

SUBJECT A




Fig. 4. Values for subject A in metabolic balance study. Open bars, Daily urinary sodium excretion; solid bars, potassium excretion. The cortisone/cortisol ratios in the grapefruit juice period are significantly different from each normal diet period. The urinary free cortisol during grapefruit juice differs significantly from the first but not the second control period.

RESULTS

In vitro study

The renal cortex homogenate obtained from male guinea pigs readily converted cortisol to cortisone after 1 hour of incubation at 37°C with NAD or NADP as a cofactor. There was no difference in the conversion rate between NAD and NADP (mean \pm SD, 75.1% \pm 7.53% with NAD versus 71.0% \pm 6.85% with NADP; $p > 0.05$). However, the Michaelis-Menten constant (K_m) values for NAD and NADP calculated from the double reciprocal plots were significantly different (36.4 \pm 7.02 μ mol/L with NAD versus 57.6 \pm 13.1 μ mol/L with NADP; $p < 0.05$).

Each flavonoid inhibited the enzyme in a concentration-dependent manner. The inhibition rates for most flavonoids with use of NAD differed from that with use of NADP (Fig. 2). The IC_{50} values of the flavonoids to inhibit the NAD- or NADP-utilizing form of 11 β -OHSD are given in Table I. Quercetin was the most potent inhibitor

with NAD; apigenin, kaempferol, and naringenin had similar potencies. Apigenin was found to be the most potent inhibitor with NADP, whereas the potency of naringenin, kaempferol, and quercetin were similar. Naringin and hesperidin were poor inhibitors, and their IC_{50} values were much less than that of their aglycons, naringenin and hesperetin. The IC_{50} values of each flavonoid with use of NAD as a cofactor differed from the IC_{50} values with NADP as the cofactor, except for kaempferol.

In vivo dose-response study

The two subjects who drank grapefruit juice showed a dose-dependent decrease in their urinary cortisone/cortisol ratios, indicating inhibition of 11 β -OHSD by grapefruit juice (Fig. 3). Each 4-day period was statistically significantly different from the control periods, and the low- and high-dose periods differed in subject 1 statistically and in subject 6 numerically but not statistically.

SUBJECT B




Fig. 5. Values for subject B in metabolic balance study. Open bars, Daily urinary sodium excretion; solid bars, potassium excretion. The cortisone/cortisol ratios and the urinary free cortisol during the grapefruit juice period are significantly different from both control periods.

In vivo metabolic balance study

The actual values for each subject are presented in Figs. 4 and 5. The mean ratios of the urinary cortisone to cortisol fell with grapefruit juice, and recovered during the second control period (mean \pm SD, 3.27 ± 0.48 during the first control period, 1.88 ± 0.28 during the grapefruit juice period, and 3.52 ± 0.46 during the second control period). Urinary free cortisol levels also were increased during the grapefruit juice period and returned to the control level after subjects ceased to drink grapefruit juice (mean \pm SD, 34.3 ± 19.0 for the first control period, 58.2 ± 9.2 for the grapefruit juice period, and 26.3 ± 12.9 for the second control period; $p < 0.05$ for each control period compared with grapefruit juice period). There was a little change in the body weight during the study (67.3 ± 0.4 , 67.8 ± 0.3 , and 67.5 ± 0.1 kg for subject A and 73.5 ± 0.8 , 74.4 ± 0.2 , and 74.4 ± 0.1 kg for subject B, in the first control, grapefruit juice, and second

control periods, respectively). The urinary sodium and potassium values were variable during the study. There were no significant changes in plasma potassium levels and blood pressure values during the study.

DISCUSSION

Grapefruit juice is known to inhibit the first-pass oxidation of felodipine and nitrendipine,^{19,20} presumably because of compounds in the juice that inhibit cytochrome P450 3A. We did this study to learn if it also inhibited another in vivo oxidation, that of 11β -OHSD. We tested several flavonoids present in grapefruit juice for their ability to inhibit 11β -OHSD from guinea pig renal cortex microsomes. The two different isoforms of the enzyme, NAD-dependent and NADP-dependent 11β -OHSD, had different K_m values for cortisol, and the flavonoids had different IC_{50} values for the two forms. We confirmed

the finding of Walker et al.¹¹ of similar conversion rates for the two forms.

There are a number of flavonoids in grapefruit juice. Naringin is the most abundant flavonoid, present in concentrations of up to 1 mmol/L.²¹ It is thought to be converted to the aglycone naringenin in the intestine after oral administration. Because the flavonoids in grapefruit juice inhibited 11 β -OHSD in vitro, we evaluated the ability of grapefruit juice to inhibit the enzyme in vivo. Drinking grapefruit juice lowered the urinary cortisone/cortisol ratios in the two investigators and both inpatient subjects, indicating in vivo inhibition of the enzyme. At the doses consumed, it did not change renal electrolyte clearance. Natural licorice in very high doses causes mineralocorticoid effects by inhibition of this enzyme.^{4,22,23} We think that grapefruit juice inhibited 11 β -OHSD, but the effect was too mild to cause electrolyte changes in these subjects because their urinary free cortisol did not exceed the normal range. A possible alternative explanation is that ring A reduction of cortisol and not 11 β -OHSD inhibition is the major cause of the syndrome of apparent mineralocorticoid excess.²³⁻²⁶

If the conventional view that 11 β -OHSD inhibition is the cause of the syndrome, and if there are differences in different people's enzyme sensitivity to these inhibitors, as we found with different strains of guinea pigs for gossypol inhibition,⁸ some people may increase their potassium clearance if they drink large amounts of grapefruit juice. Furthermore, flavonoids are sold in tablet form in health food stores and drug stores. If people take large quantities of flavonoids as dietary supplements, it is possible that the flavonoids may cause sufficient 11 β -OHSD inhibition to produce the syndrome of apparent mineralocorticoid excess.

We thank Patricia Danton for her help.

References

1. Ulick S, Ramirez LC, New MI. An abnormality in steroid reductive metabolism in a hypertensive syndrome. *J Clin Endocrinol Metab* 1977;44:799-802.
2. Card WL, Mitchell W, Strong JA, Taylor NRW, Tompsett SL, Wilson JMG. Effects of liquorice and its derivatives on salt and water metabolism. *Lancet* 1953;1:663-8.
3. Borst JCG, ten Holt SP, de Vries LA, Molhuysen JA. Synergistic action of liquorice and cortisone in Addison's and Simmond's disease. *Lancet* 1953;1:657-63.
4. Farese RV, Biglieri EG, Shackleton CHL, Irony I, Gomez-Fontes R. Licorice-induced hypermineralocorticoidism. *N Engl J Med* 1991;325:1223-7.
5. Monder C, White PC. 11 β hydroxysteroid dehydrogenase. *Vitam Horm* 1993;47:187-271.
6. Edwards CRW, Walker BR, Benediktsson R, Seckl JR. Congenital and acquired syndromes of apparent mineralocorticoid excess. *J Steroid Biochem Mol Biol* 1993;45:1-5.
7. Qian SZ, Wang ZG. Gossypol: a potential antifertility agent for males. *Annu Rev Pharmacol Toxicol* 1984;24:329-60.
8. Sang GW, Lorenzo BJ, Reidenberg MM. Inhibitory effects of gossypol on corticosteroid 11 β -hydroxysteroid dehydrogenase from guinea pig kidney: a possible mechanism for causing hypokalemia. *J Steroid Biochem Mol Biol* 1991;39:169-76.
9. Song D, Lorenzo BJ, Reidenberg MM. Inhibition of 11 β hydroxysteroid dehydrogenase by gossypol and bioflavonoids. *J Lab Clin Med* 1992;120:792-7.
10. Zhang YD, Lorenzo BJ, Reidenberg MM. Inhibition of 11 β hydroxysteroid dehydrogenase obtained from guinea pig kidney by furosemide, naringenin and some other compounds. *J Steroid Biochem Mol Biol* 1994;49:81-5.
11. Walker BR, Campbell JC, Williams BC, Edwards CRW. Tissue-specific distribution of the NAD⁺-dependent isoform of 11 β -hydroxysteroid dehydrogenase. *Endocrinology* 1992;131:970-2.
12. Mercer WR, Krozowski ZS. Localization of an 11 β -hydroxysteroid dehydrogenase activity to the distal nephron. Evidence for the existence of two species of dehydrogenase in the rat kidney. *Endocrinology* 1992;130:540-543.
13. Monder C. The forms and functions of 11 β -hydroxysteroid dehydrogenase. *J Steroid Biochem Mol Biol* 1993;45:161-5.
14. Lakshmi V, Nath N, Muneyyirch-Delale O. Characterization of 11 β -hydroxysteroid dehydrogenase of human placenta: evidence for the existence of two species of 11 β -hydroxysteroid dehydrogenase. *J Steroid Biochem Mol Biol* 1993;45:391-7.
15. Hertog MGL, Feskens EJM, Hollman PCH, Katan MB, Kromhout D. Dietary antioxidant flavonoids and risk of coronary heart disease: the Zutphen elderly study. *Lancet* 1993;342:1007-11.
16. Brown JP. A review of the genetic effects of naturally occurring flavonoids, anthraquinones, and related compounds. *Mutat Res* 1980;75:243-77.
17. Ayres PJ, Garrod O, Simpson SA, Tait JF. A method for the determination of aldosterone, cortisol, and corticosterone in biological extracts, particularly applied to human urine. *Biochem J* 1957;65:639-46.
18. Peterson RE, Pierce CE. The metabolism of corticosterone in man. *J Clin Invest* 1960;30:741-57.
19. Bailey DG, Spence JD, Munoz C, Arnold JMO. In-

- teraction of citrus juices with felodipine and nifedipine. *Lancet* 1991;337:268-9.
20. Soons P, Vogels B, Roosemalen N, et al. Grapefruit juice and cimetidine inhibit stereoselective metabolism of nitrendipine in humans. *CLIN PHARMACOL THER* 1991;50:394-403.
 21. Bailly DG, Arnold MO, Spence JD. Grapefruit juice and drugs. *Clin Pharmacokinet* 1994;26:91-8.
 22. Stewart PM, Wallace AM, Valentino R, Burt D, Shackleton CH, Edwards CR. Mineralocorticoid activity of liquorice: 11 β -hydroxysteroid dehydrogenase deficiency comes of age. *Lancet* 1987;2:821-4.
 23. Epstein MI, Espiner EA, Donald RA, Hughes H, Cowles RJ, Lun S. Licorice raises urinary cortisol in man. *J Clin Endocrinol Metab* 1978;47:397-400.
 24. Ulick S, Tedde R, Wang JZ. Defective ring A reduction of cortisol as the major metabolic error in the syndrome of apparent mineralocorticoid excess. *J Clin Endocrinol Metab* 1992;74:593-9.
 25. Ulick S, Wang JZ, Hanukoglu A, Rösler A. The effect of carbenoxolone on the peripheral metabolism of cortisol in human patients. *J Lab Clin Med* 1993;22:673-6.
 26. Morris DJ. The role of steroid metabolism in protective and specificity conferring mechanisms of mineralocorticoid action. *Vitam Horm* 1995;50:461-85.

Availability of JOURNAL Back Issues

As a service to our subscribers, copies of back issues of *CLINICAL PHARMACOLOGY & THERAPEUTICS* for the preceding 5 years are maintained and are available for purchase from the publisher, Mosby-Year Book, Inc., at a cost of \$12.50 per issue. The following quantity discounts are available: 25% off quantities of 12 to 23, and 33% off quantities of 24 or more. Please write to Mosby-Year Book, Inc., Subscription Services, 11830 Westline Industrial Dr., St. Louis, MO 63146-3318, or call (800)453-4351 or (314)453-4351 for information on availability of particular issues. If unavailable from the publisher, photocopies of complete issues are available from UMI, 300 N. Zeeb Rd., Ann Arbor, MI 48106 (313)761-4700.

- C. N., Wirth, P., Colligen, J. E., Albrechtsen, R., Mudry, M., and Sobel, M. E. (1986). Altered levels of laminin receptor mRNA in various human carcinoma cells that have different abilities to bind laminin. *Proc. Natl. Acad. Sci. U. S. A.* 83, 7137-7141.
- Wille, M. S., and Skubitz, A. P. N. (1991). Human keratinocytes adhere to multiple distinct peptide sequences of laminin. *J. Invest. Dermatol.* 97, 141-146.
- Woo, H.-J., Lutz, M. M., Jung, J. V., and Mercier, A. M. (1991). Carbohydrate-binding protein 35 (huc-2), a laminin-binding lectin, forms functional dimers using cysteine 186. *J. Biol. Chem.* 266, 18419-18422.
- Yamada, K. M. (1991). Adhesive recognition sequences. *J. Biol. Chem.* 266, 12809-12812.
- Yow, H., Wong, J. M., Chen, H. S., Lee, C., Sletta, O. D., and Chen, L. B. (1988). Increased mRNA expression of a laminin binding protein in human colon carcinoma. Complete sequence of a full-length cDNA encoding the protein. *Proc. Natl. Acad. Sci. U. S. A.* 85, 6394-6398.
- Yurchenco, P. D., Tailhery, E. C., Chroma, A. S., and Porthmann, H. (1985). Laminin polymerization in vitro. *J. Biol. Chem.* 260, 7638-7644.

11 β -Hydroxysteroid Dehydrogenase

CARL MONDER*¹ AND PERRIN C. WHITE¹

^{*}The Population Council
Center for Biomedical Research
New York, New York 10021

¹Department of Pediatrics
Cornell University Medical School
New York, New York 10021

- I. Historical Origins
- II. Distribution, Properties, and Behavior of 11-11SD
 - A. Tissue Distribution
 - B. Physiological Functions
 - C. Enzymatic Properties
 - D. Effects of Hormones
- III. Developmental Biology and 11-11SD
 - A. Fetal Development
 - B. Postnatal Development
- IV. Are 11-Dehydrocorticosteroids Biologically Active?
- V. 11-11SD in Lower Vertebrates
- VI. The Forms of 11-11SD Expression: Uniqueness or Multiplicity?
 - A. On the Question of Reversibility
 - B. Characteristics of Microsomal 11-11SD
- VIII. Clinical Studies
 - A. 11 β -Dehydrogenase Deficiency
 - B. 11-Oxoreductase Deficiency
- VIII. Enzymology and Molecular Biology
 - A. The Uniqueness of 11-11SD
 - B. Preparation and Properties of Homogeneous 11-11SD
 - C. Molecular Analysis
- IX. 11-11SD Function in Specific Organs
 - A. Kidney
 - B. The Vascular Bed
 - C. The Skin
 - D. The Nervous System
 - E. Leydig Cells, Stress, and 11-11SD
 - F. Mammary Gland
- X. Epilogue
- References

I. HISTORICAL ORIGINS

It was experimentally established in 1856 (Brown-Séquard, 1856) that adrenomedullary in animals is faint, but not until 1927 was it shown

by Huggoff and Stewart (1927) that adrenal extracts could maintain adrenalecuminized animals. By 1937, Reichstein in Basel, Switzerland, and Kendall in Rochester, Minnesota, had demonstrated that the active substances in adrenal cortical extracts were steroids. In that year Reichstein and his colleagues synthesized 11-deoxycorticosterone (Steiger and Reichstein, 1937), which, although it was not a quantitatively important secretory product of the adrenal cortex, was used for many years to treat Addisonian patients with some, but by no means total, success. A consensus soon emerged that the therapeutically active adrenal steroid contained oxygen at C-11 (Kendall, 1941; Ingle, 1940; Thorn, 1944; Olson *et al.*, 1944). What complicated discussion of the identity of the hormone of the adrenal cortex was the large number of steroids extracted from alouphorhous tissue (Fieser and Fieser, 1959). Many of these were 11-oxygenated and were thus candidates for the active steroid. Studies with isolated, perfused adrenals (Reichstein and Shoppee, 1943), analysis of urine from normal individuals (Conn *et al.*, 1951) or patients with Cushing's disease (Mason, 1950; Sprague *et al.*, 1951), and adrenal vein blood (Reich *et al.*, 1950; Nelson *et al.*, 1951; Bush, 1953) led to the conclusion that cortisol (F) and corticosterone (B) are the primary secretory products of the adrenal gland. There was evidence that adrenal enzymes were capable of oxidizing the 11-hydroxy group of corticosterone and cortisol to an 11-keto group (Hechter *et al.*, 1951; Burrell *et al.*, 1953; Fackas *et al.*, 1970), and that patients treated with cortisone (E) excreted some F (Mason, 1950; Sprague *et al.*, 1951; Burton *et al.*, 1953; Miller and Axelrod, 1953). Although the interconversion of the 11-oxygenated pairs, F and E, and B and 11-dehydrocorticosterone (A) could be readily demonstrated, their biological relationships to each other were not obvious. It was suggested that both cortisol and cortisone were converted to the true active steroid (Hechter *et al.*, 1953). However, cortisone was ineffective when injected into arthritic joints, in contrast with the pronounced antiarthritic effects of the orally administered steroid (Hollander *et al.*, 1951; Zaccò *et al.*, 1954). The conclusion that cortisone is biologically inactive and must be converted to its physiologically active form, cortisol, by reduction of the 11-oxo group was supported by other clinical observations (Boland, 1952; Dixon and Bywaters, 1953; Cope and Hurlock, 1954), and by studies with experimental animals (Eisenstein, 1952; Fish *et al.*, 1953; Burton *et al.*, 1953; Campi *et al.*, 1953; Amelung *et al.*, 1954; Dabrtner, 1951; Savard *et al.*, 1953). In 1953, sufficient quantities of cortisol became available to permit its metabolism in the studied Fieser and Fieser, 1959). Oral administration of cortisol acetate to human subjects resulted in the

excretion of 11-oxo (C_{21} and C_{19} steroids (Burstein *et al.*, 1953). An enzyme responsible for catalyzing the oxidation of cortisol to cortisone was found in rat liver (Amelung *et al.*, 1953a,b) and named "11 β -hydroxy steroid dehydrogenase" (Hubener *et al.*, 1956). It is now known as 11 β -hydroxysteroid dehydrogenase (11-HSD). Figure 1 illustrates the transformations catalyzed by this enzyme.

II. DISTRIBUTION, PROPERTIES, AND BEHAVIOR OF 11-HSD

A. Tissue Distribution

Catalysis of 11-oxidation and 11-oreduction is not uniformly distributed among tissues. In liver, 11-oreduction is the dominant activity, in most other tissues, it is 11 β -hydroxy oxidation. Whether this behavior is due to the expression of separate enzymes or to the tissue-specific behavior of a unique 11 β -hydroxysteroid dehydrogenase was a question first posed 35 years ago (Bush, 1956, 1959; Bush and Mahesh, 1959a). Most investigators have interpreted the results of their studies on steroid metabolism at position 11 in terms of a single enzyme, designated by the Nomenclature Committee of the International Union of Biochemistry as EC 1.1.1.146 (11 β -hydroxysteroid:NADP+ 11-oreductase) (Webb, 1964). Within this context, there have been suggestions of multiple enzyme forms, based on the fact that the char-




FIG. 1. Transformations catalyzed by 11 β -hydroxysteroid dehydrogenase.

activities of the enzyme in different tissues varied over a range that was too wide to be accounted for by error or interlaboratory variability. Table I summarizes our current knowledge of the tissue distribution of 11- β -HSD. For convenience, activity reported to occur in the oxidative direction is called corticosteroid 11 β -dehydrogenase (11-DH), and reductive activity is called corticosteroid 11-oxoreductase (11-OR). From the available literature, it is difficult to evaluate whether 11-DH and 11-OR activities are both present in a particular organ (Fish *et al.*,

TABLE I
DISTRIBUTION OF 11- β -HSD IN MAMMALIAN TISSUES*

| Tissue | 11 β -Dehydrogenase | 11-Oxoreductase |
|-------------------|---------------------------|-----------------|
| Liver | II, D, M, Rb, G, Rt | M, Rb, G, Rt, C |
| Kidney | II, D, M, Rb, G, Rt | — |
| Lung | D, M, Rb, G, Rt | II, M, Rb, Rt |
| Testis | II, Rb, Rt, M | — |
| Brain | D, M, Rt, B | — |
| Spleen | D, M, G | R |
| Adrenal cortex | II, D, Rb, Rt, Sh, C, MV | M |
| Diaphragm | D, Rt | MV |
| Skeletal muscle | M, Rt, C | — |
| Blood vessels | Rb, Rt | — |
| Heart | — | — |
| Lymphocytes | — | — |
| Thymocytes | M, Rt | — |
| Small intestine | M, Rt | — |
| Colon | II, D, Rt | — |
| Placenta | Rt | — |
| Ovary | II, Rt, B | — |
| Uterus | — | — |
| Myometrium | — | — |
| Amniotic membrane | — | — |
| Uterine | — | — |
| Chorion | — | — |
| Adipose tissue | — | — |
| Salivary gland | — | — |
| Membrane gland | — | — |
| Skin | — | — |
| Gingival tissue | — | — |

*II, human; D, dog; M, mouse; Rb, rabbit; G, guinea pig; Rt, rat; C, cattle; N, baboon; MV, meadow vole; Sh, sheep. The table cites positive identification of 11- β -HSD in the oxidative direction (11 β -dehydrogenase) or in the reductive direction (11-oxoreductase) in the investigated species. Absence of measurable activity or no reported activity is indicated by a dash.

1953; Bush *et al.*, 1968; Koerner, 1969; Monder and Lakshmi, 1989a). Where reversibility has been reported, the results have not generally been consistent. The 11- β -HSD in human adipose tissue has been reported to catalyze only oxidation. Results with intestinal mucosa and skin (Murphy, 1981; Hsieh and Hsiao, 1986; Hammami and Sileri, 1990; Burton and Anderson, 1983) are in conflict. The reasons for the great differences between laboratories are not clear, but the following sources of variation are probably important: (a) the pH of measurement; (b) the relative stabilities of the dehydrogenase and oxoreductase activities (Lakshmi and Monder, 1985a); (c) incompletely expressed or "latent" enzyme (Lakshmi and Monder, 1985b); (d) the age of the animal, its sex and diet, and the possible presence of endogenous inhibitors or activators; (e) the developmental stage of the animal (Murphy, 1981); (f) substrate specificity (Koerner, 1969).

B. PHYSIOLOGICAL FUNCTIONS

It has been suggested that 11- β -HSD protects cells against the toxic effects of excess corticosteroid (Dougherty *et al.*, 1961; Berliner, 1965). 11 β -Dehydrogenase may be a component of a degradation pathway, strategically placed to inactivate corticosteroids prior to their exposure to receptor or to prevent the return of the steroid to receptor. The enzyme also serves a conservation function, since the oxidized form of the steroid can be reduced by 11-oxoreductase to its active reduced form thus contributing to the circulating cortisol, and providing a buffer against the changes in blood level caused by paroxysmal secretion of the adrenal. The enzyme can thus integrate the availability of corticosteroids to target organs and their metabolism. An overview of these proposed interrelationships is presented in Fig. 2.

C. ENZYMATIC PROPERTIES

1. Substrate Specificity

In Tables IIa and IIb are listed all steroids for which oxidation or reduction catalyzed by 11 β -hydroxysteroid dehydrogenase have been reported. Table IIc lists the steroids that were found to be neither oxidized nor reduced at C-11. Based on the data in Table II the qualitative effects of substituents on oxidation, i.e., whether steroids carrying the indicated functional group are substrates for 11- β -HSD, are summarized in Table III.

From the data summarized in Tables II and III it is possible to deduce




FIG. 2. An overview of the proposed role of 11 β -hydroxysteroid dehydrogenase in modulating corticosteroid function. In this model, 11 β -hydroxysteroid function and metabolism are illustrated with cortisol (F), and its 11-oxo derivative, cortisone (E). F, synthesized in the adrenal cortex, is transported through the circulation (1) to the target cell (2). The steroid binds to the cytosolic receptor (3), which is transported to the nucleus (4), or within the nucleus to the steroid receptor (5). At an as yet undefined point in the cell response process the steroid released (6) from the R_h is oxidized to E by nuclear 11- β -HSD (6), or it leaves the nucleus (7) and is oxidized by microsomal 11-HSD (8). The F, thus formed, is transported in the circulation (9) to the liver (10) where it is metabolized to inactive end product, or converted to cortisone by 11- α -reductase (11) to the liver corticoid pool, as well (13). The kidney is a major contributor to cortisone oxidation (14). Cortisone may be converted to active steroid by peripheral tissues (15) as well as liver; however, evidence for this regenerative pathway is scanty.

how structural changes in steroids bring about corresponding changes in bioactivity. There is considerable evidence that any substituent that inhibits oxidation of the 11 β -hydroxy group can enhance corticosteroid potency even if the structural modification does not have an enhancing effect on some other parameter, such as receptor affinity. Bush *et al.* (1968) and Koerner (1969) have analyzed the effects of substituents on oxidation or reduction of 11-oxygenated steroids. From their data, the following conclusions may be drawn: (a) A flat A/B junction (5a) is essential for binding to the active site, whereas a buckled A/B junction (5b) prevents binding; (b) bulky groups that obstruct the α -surface (2a-

TABLE II
Steroid Substrates of 11 β -Hydroxysteroid Dehydrogenase

| (a) 11-OH \rightarrow 11-oxo | |
|--|--|
| Cortisol | Osinaki (1960); Koerner (1969) |
| Corticosterone | Osinski (1960); Koerner (1969) |
| 11 β ,17,20 β ,21-Tetrahydroxy-pregn-4-en-3-one | Bradlow <i>et al.</i> (1968) |
| 11 β -Hydroxypregn-4-en-20-one | Koerner (1969); Bush <i>et al.</i> (1968) |
| 3 α ,11 β ,17,21-Tetrahydroxy- Δ^5 -pregnan-20-one | Bush and Mahesh (1959a); Koerner (1969) |
| 11 β ,17 α ,20 β ,21-Tetrahydroxy-pregn-4-en-3-one | Koerner (1969) |
| 11 β ,17 α ,21-Trihydroxy- Δ^5 -pregnan-3,20-dione | Koerner (1969) |
| 11 β ,17 α ,21-Trihydroxy- Δ^5 -pregnan-3,20-dione | Koerner (1969) |
| 3 β ,11 β ,17,21-Tetrahydroxy- Δ^5 -pregnan-20-one | Koerner (1969) |
| 3 β ,11 β ,17,21-Tetrahydroxy-pregn-5-en-20-one | Koerner (1969) |
| 11 β ,17,21-Trihydroxy-pregn-1,4-diene-3,20-dione | Koerner (1969) |
| 11 β ,17,20 α ,21-Tetrahydroxy-pregn-4-en-3-one | Koerner (1969) |
| 11 β ,17-Dihydroxy-pregn-4-en-3,20-dione | Koerner (1969) |
| 16 α -Methyl-cortisol | Bush <i>et al.</i> (1968) |
| 16 α -Methyl-cortisol | Bush <i>et al.</i> (1968) |
| 3 α ,11 β ,17,21-Tetrahydroxy- Δ^5 -pregnan-20-one-3-acetate | Koerner (1969) |
| (b) 11-oxo \rightarrow 11 β -OH | |
| Cortisone | Fish <i>et al.</i> (1953); Burton <i>et al.</i> (1953) |
| 21-Hydroxypregn-4-en-3,11,20-trione | Mason (1950) |
| Androst-4-ene-3,11,17-trione | Hübener <i>et al.</i> (1956); Bush <i>et al.</i> (1968) |
| Pregn-4-ene-3,11,20-trione | Hübener <i>et al.</i> (1956); Bush <i>et al.</i> (1968) |
| 17 α ,20 β ,21-Trihydroxy-pregn-4-en-3,11-dione | Koerner (1969); Hübener <i>et al.</i> (1956) |
| 17,21-Dihydroxy-pregn-1,4-diene-3,11,20-trione | Bush <i>et al.</i> (1968) |
| 17,21-Dihydroxy- Δ^5 -pregnan-3,11,20-trione | Bush <i>et al.</i> (1968) |
| 8 α -Fluorocortisone | Bush <i>et al.</i> (1968) |
| 9 α -Fluoro-11-oxo-pregsterone | Bush <i>et al.</i> (1968) |
| 12 α -Fluoro-11-oxo-pregsterone | Bush <i>et al.</i> (1968) |
| 3 α -Hydroxy- Δ^5 -androstane-11,17-dione | Bush <i>et al.</i> (1968) |
| 9 α -Fluoro-androst-4-en-3,11,17-trione | Bush <i>et al.</i> (1968) |
| 9 α -Chlorocortisone | Bush <i>et al.</i> (1968) |
| 9 α -Chloro-androst-4-en-3,11,17-trione | Bush <i>et al.</i> (1968) |
| 12 α -Bromo-11-dehydrocorticosterons | Bush <i>et al.</i> (1968) |
| (c) Unreactive steroids | |
| Tetrahydrocortisol | Hübener <i>et al.</i> (1956); Bradlow <i>et al.</i> (1968) |

(continued)

TABLE II (Continued)

| | |
|--|---|
| Tetrahydrocortisone | Huebner <i>et al.</i> (1956) |
| 3 α -Hydroxy-5 β -androstan-11,17-dione | Huebner <i>et al.</i> (1956); Bush and Mahesh (1959b) |
| 17 α ,20 β ,21-Trihydroxypregn-4-ene-3,11-dione | Bradlow <i>et al.</i> (1966) |
| 11 β ,17 α ,21-Trihydroxy-5 β -pregnan-3-one | Bush <i>et al.</i> (1956b); Koerner (1969) |
| 2 α -Methylcortisone | Bush and Mahesh (1959b) |
| 2 α -Methylcortisol | Huebner <i>et al.</i> (1956) |
| 2 α ,17 α ,20 α ,21-Tetrahydroxy-5 β -pregnan-11-one | Huebner <i>et al.</i> (1956) |
| 3 α ,17 α ,20 β ,21-Tetrahydroxy-5 β -pregnan-11-one | Koerner (1969) |
| 11 β ,17-dihydroxyestrone | Bush <i>et al.</i> (1956) |
| 12 α -Fluoro-11-dehydrocorticosterone | Bush <i>et al.</i> (1956) |
| 12 α -Methyl-11-oxoprogesterone | Bush <i>et al.</i> (1956) |
| 9 α -Fluorocortisol | Bush <i>et al.</i> (1956) |
| 12 α -Fluoro-11 β -hydroxypregnenolone | Bush <i>et al.</i> (1956) |
| 12 α -Fluorocorticosterone | Bush <i>et al.</i> (1956) |
| 12 α -Fluorocorticosterone | Bush <i>et al.</i> (1956) |
| 2 α -Methyl-9 α -fluorocortisol | Bush <i>et al.</i> (1956) |
| 2 α -Methyl-9 α -fluoro-11 β -hydroxypregnenolone | Bush <i>et al.</i> (1956) |
| 11 β ,17,21-Trihydroxy-16 α -methyl-pregn-1,4-diene-3,20-dione | Bush <i>et al.</i> (1956) |
| 9 α -Fluoro-11 β ,16 α ,17 α ,21-tetrahydroxypregn-1,4-diene-3,20-dione | Bush <i>et al.</i> (1956) |
| 2 α -Methyl-androst-4-ene-3,11,17-trione | Bush <i>et al.</i> (1956) |
| Cortisol-21-sulfate | Koerner (1969) |
| Cortisol-21-phosphate | Koerner (1969) |
| Cortisol-21-acetate | Koerner (1969) |
| Cortisol-21-hemiacetate | Koerner (1969) |
| 3 α -Hydroxy-5 β -androstane-11,17-dione | Bush <i>et al.</i> (1956) |

methyl) inhibit binding; (c) aromatic A ring is forbidden; (d) effects of halogens are more likely to be the consequence of their inductive effects than their steric effects; (e) steroids with bulky substituents (acetyl, phosphate) at C-21 are not substrates. Structural studies lead to the conclusion that the steroid α -surface binds to the enzyme, and that hydrogen transfer occurs from the 11 α -position. Consequently, variations in the velocities of nonhalogenated steroids can be attributed to steric factors.

The inability of 2 α -methyl steroids to be oxidized or reduced at C-11 played an important historical role in reinforcing the conclusion that cortisol, and not cortisone, was the active steroid hormone (Bush and Mahesh, 1959b). The metabolically active 2 α -methyl-11 β -hydroxysteroid could not be oxidized by 11-HSD and the 2 α -methyl-11-oxosteroid was inactive as a glucocorticoid, supporting the importance

TABLE III
EFFECTS OF FUNCTIONAL GROUPS
ON SUBSTRATE SPECIFICITY OF 11-HSD

| Functional group | Oxidation* | Reduction* |
|-------------------------------|------------|------------|
| 1-one | + | + |
| 2 α -Methyl | - | - |
| 3 α -Hydroxy | + | NR |
| 3 β -Hydroxy | + | NR |
| 4 α -3-Oxo | + | + |
| 5 β | - | - |
| 6 α | + | + |
| 9 α -Fluoro | - | + |
| 12 α -Fluoro | - | + |
| 16 α -Methyl | + | NR |
| 17 α -Hydroxy | + | + |
| 20 α -Hydroxy (a or b) | + | + |
| 21-Methyl | + | + |
| 21-Hydroxy | + | + |

* +, steroid with indicated functional group is a substrate; -, steroid with indicated functional group is not a substrate; NR, not reported. In evaluating the effects of multiple substituents on substrate specificity, "—" precedes "1". Substituents for which only single examples exist are omitted here, and are listed in Table II.

of the 11 β -hydroxy group in glucocorticoid function. These results also helped to disprove the hypothesis that steroids affect metabolism by directly participating as cofactors in transhydrogenation reactions (Williams-Ashman and Liao, 1964).

2. Steroid Inhibitors

The catalysis of 11-oxidation is inhibited by a number of structurally diverse steroids, including representatives of the C₂₁ and C₁₉ series. Inhibitors of reduction have also been shown to include C₂₁ and C₁₉ steroids, though fewer studies have been performed in this direction. Some C₁₉, C₂₀, and C₂₁ steroids inhibit neither oxidation nor reduction. The steroids that have been investigated for their ability to inhibit 11-HSD are listed in Table IV. On the basis of the data, we conclude that inhibition of 11 β -dehydrogenase is not caused by the following: 2 α -Cl, 6 β -H, 6 α -OH, 6 β -OH, 12 α -OH, 16 α -OH, 20 α -OH, 11-oxo, 16(17)-ene. Steroids devoid of oxygen at C-11 are generally not inhibitors, or inhibit oxidation poorly. The 11 α -

TABLE IV
Steroid Inhibitors of 11 β -Hydroxysteroid Dehydrogenase

| (a) Oxidation (11-OH \rightarrow 11-oxo) | |
|--|---|
| C ₂₁ and C ₂₅ steroids | |
| 11 α ,17,21-Trihydroxy-pregn-4-en-3-one (11-epicortisol) | Burton (1966) |
| 11 α ,17,21-Trihydroxy-pregn-1,4-dione-3-one | Burton (1966) |
| 11 α -Hydroxypregn-4-en-3-one (11 α -hydroxyprogesterone) | Burton (1966), Murphy and Vedady (1982) |
| 17,21-Dihydroxypregn-4-ene-3,11-dione (cortisone) | Bernal <i>et al.</i> (1980), Murphy (1979b) |
| Cortisol 21-acetate | Bernal <i>et al.</i> (1980), Murphy and Vedady (1982) |
| Pregesterone | Bernal <i>et al.</i> (1980), Murphy and Vedady (1982) |
| 11-Hydroxy-11 α -methyl-11 β -fluorohydrocortisone (10 α -amelinone) | Bernal <i>et al.</i> (1980) |
| 11 α ,17,21-Trihydroxypregn-1,4-dione-3-one (prednisolone) | Bernal <i>et al.</i> (1980), Murphy and Vedady (1982) |
| 9 α -Fluorocortisol | Bush <i>et al.</i> (1966) |
| 3 α ,11 β ,17,21-Tetrahydroxy-6 α -pregnane-20-one (alobendrolol) | Dicks and DeMoore (1966) |
| 3 α ,11 β ,17,20 β ,21-Penta-hydroxy-6 α -pregnane (alocortol) | Dicks and DeMoore (1966) |
| 11 α ,17 α ,21-Trihydroxy-5 α -pregnane-3,20-dione (alodehydrocortisol) | Dicks and DeMoore (1966) |
| 11 β -Hydroxycortisone | Monder and Lakshmi (1989a) |
| 11 β -Hydroxymethyl-4-ene-3,17-dione | Dicks and DeMoore (1966), Monder and Lakshmi (1989a) |
| 3 α ,11 β ,17 β -Trihydroxyandrosterone | Monder and Lakshmi (1989a) |
| 3 β -Hydroxyandrost-5-en-17-one | Monder and Lakshmi (1989a) |
| 11 β ,17 β -Dihydroxy-6 β -androst-3-one | Monder and Lakshmi (1989a) |
| 11 β ,17 β -Dihydroxy-6 α -androst-3-one | Monder and Lakshmi (1989a) |
| (b) Reduction (11-oxo \rightarrow 11-OH) | |
| C ₂₁ and C ₂₅ steroids | |
| 11-Dehydrocortisone | Burton <i>et al.</i> (1976) |
| 3 α ,17,21-Trihydroxy-6 β -pregnan-3,20-dione (tetrahydrocortisol) | Bernal <i>et al.</i> (1980) |
| 21-Hydroxypregn-4-ene-3,11,20-trione | Bernal <i>et al.</i> (1980) |
| Androst-4-ene-3,11,20-trione | Dicks and DeMoore (1966) |
| 3 β -Hydroxyandrost-5-en-17-one | Dicks and DeMoore (1966) |

(continued)

TABLE IV (Continued)

| (c) Do not inhibit (11 β -OH \rightarrow 11-oxo) | |
|---|---|
| C ₂₁ steroids | |
| 21-Hydroxypregn-4-ene-3,20-dione | Murphy and Vedady (1982) |
| 17 α ,21-Dihydroxypregn-4-ene-3,11-dione | Murphy and Vedady (1982) |
| 11 β -Hydroxypregn-4-ene-3,20-dione-21-sulfate | Murphy and Vedady (1982) |
| 11 β ,17 α -Dihydroxypregn-4-ene-3,20-dione-21-sulfate | Murphy and Vedady (1982) |
| 6 α -Hydroxypregn-4-ene-3,20-dione | Murphy and Vedady (1982) |
| 12 α -Hydroxypregn-4-ene-3,20-dione | Murphy and Vedady (1982) |
| 6 β ,11 β ,17 α ,21-Tetrahydroxypregn-4-ene-3,20-dione | Murphy and Vedady (1982) |
| 11 β ,21-Dihydroxy-18-oxo-pregn-4-ene-3,20-dione | Murphy and Vedady (1982) |
| 16 α -Hydroxypregn-4-en-3-one | Murphy and Vedady (1982) |
| 16 α -Hydroxypregn-4-en-3-one | Murphy and Vedady (1982) |
| 3,20-Dioxo-pregn-4,16-dione | Murphy and Vedady (1982) |
| 3 α ,20 α -Dihydroxy-6 β -pregnane | Murphy and Vedady (1982) |
| 6 α -Fluor-11 β ,17 α ,21-trihydroxy-11 β -methylpregn-1,4-dione-3,20-dione | Murphy and Vedady (1982) |
| Tetrahydrocortisol | Bernal <i>et al.</i> (1980), Dicks and DeMoore (1966) |
| 3 α ,11 β ,17,21 α -Penta-hydroxy-6 β -pregnane (cortisol) | Dicks and DeMoore (1966) |
| 3 α ,11 β ,17,20 β ,21-Penta-hydroxy-6 β -pregnane (p-cortisol) | Dicks and DeMoore (1966) |
| Tetrahydrocortisone | Bernal <i>et al.</i> (1980), Dicks and DeMoore (1966) |
| 2 α -Methylcortisol | Bush <i>et al.</i> (1966) |
| C ₂₅ steroids | |
| 3 β ,11 β -Dihydroxy-5 α -androst-17-one | Murphy and Vedady (1982) |
| 3 α ,11 β -Dihydroxy-5 α -androst-17-one | Murphy and Vedady (1982) |
| 3 β ,11 β -Dihydroxy-5 β -androst-17-one | Murphy and Vedady (1982) |
| 3 β ,11 β ,16 α -Trihydroxyandrost-5-en-17-one | Monder and Lakshmi (1989a), Murphy and Vedady (1982) |
| 3 α ,11 β -Dihydroxy-5 β -androst-17-one | Vedady (1982) |
| 3 β -Hydroxyandrost-5-en-17-one-3-anilide | Murphy and Vedady (1982) |
| Testosterone | Bernal <i>et al.</i> (1980) |
| 6 α -Hydroxycortisone | Bernal <i>et al.</i> (1980) |
| 3 α -Hydroxy-5 β -androst-17-one | Dicks and DeMoore (1966) |
| 3 α -Hydroxy-6 α -androst-17-one | Dicks and DeMoore (1966) |
| Androst-4-ene-3,11,17-trione | Dicks and DeMoore (1966) |
| Dihydroepiandrosterone | Dicks and DeMoore (1966) |
| 11 β -Hydroxy-5 β -androstane | Pope and Albrecht (1984) |
| 3 α ,11 β -Dihydroxyandrost-17-one | Monder and Lakshmi (1989a) |
| | Monder and Lakshmi (1989a) |

(continued)

TABLE IV (Continued)

| | | |
|---|--|---|
| C ₁₉ | | Bernal <i>et al.</i> (1980); Abramovitz <i>et al.</i> (1984) |
| Estriol | | Bernal <i>et al.</i> (1980); Abramovitz <i>et al.</i> (1984) |
| Estrene | | Abramovitz <i>et al.</i> (1984) |
| (d) Do not inhibit (11-oxo → 11-OH) | | |
| C ₂₁ | | |
| 2 α -Methylcortisone | | Bush <i>et al.</i> (1968) |
| Cortisol | | Bush <i>et al.</i> (1968) |
| 20 α -Cortol | | Bush <i>et al.</i> (1968) |
| 20 α -Cortol | | Bush <i>et al.</i> (1968) |
| 3 α ,11 β ,17,20 α ,21-Pentahydroxy-5 α -pregnen-20-one (allo-cortol) | | Bush <i>et al.</i> (1968) |
| Cortolone | | Bush <i>et al.</i> (1968) |
| C ₂₇ | | |
| Androst-4-ene-3,17-dione | | Bush <i>et al.</i> (1968) |
| 3 α -Hydroxy-5 α -androstane-17-one | | Bush <i>et al.</i> (1968) |
| 3 α -Hydroxy-5 β -androstane-17-one | | Bush <i>et al.</i> (1968) |

hydroxysteroids are probably structural analogs and are competitive inhibitors (Burton, 1965; Bernal *et al.*, 1980; Murphy and Vedady, 1981). Although this may also be true for 11 β -hydroxysteroids, it is not yet determined which are site-specific structural analogs and which are competitive substrates.

A second important binding site may be the side chain, which requires small bulk at C-21 (—CH₃, —CH₂, OH). Introduction of a bulky or charged group at C-21 may diminish the ability of the steroid to act as a substrate, but not as an inhibitor (compare 11 β -hydroxypregn-4-ene-3,20-dione and 11 β -hydroxypregn-4-ene-3,20-dione-21-sulfate) (Murphy, 1982). Cortisol-21-acetate is a potent inhibitor of decidual dehydrogenase (Bernal *et al.*, 1980), yet is not a substrate (Koerner, 1969).

Since 11 β -hydroxy-9 α -fluoro compounds are not oxidized by 11 β -hydroxysteroid dehydrogenase, inhibition of cortisol oxidation by 9 α -fluorocortisol and dexamethasone is due to displacement of the substrate at the active site by homologs in which a negative inductive effect of the halogen stabilizes the 11 β -hydroxy group (Bush *et al.*, 1968; Bush and Mahesh, 1959a). Prednisolone is oxidized by 11-HSD (Koerner, 1969) and the reported inhibitory effect (Bernal *et al.*, 1980) is probably due to substrate competition. A similar explanation is ap-

licable to inhibition of cortisol oxidation by corticosterone (Bernal *et al.*, 1980), which is a better substrate for 11-HSD (Engel *et al.*, 1955; Osinski, 1960; Koerner, 1969) than the former.

There are few studies on steroid inhibition of 11-oxoreductase. Most steroids tested (Table IVd) did not affect 11-oxoreductase. Of several that did, none could be shown to have functional groups especially associated with obligatory inhibition. Reduction of an inhibiting steroid at C-20 (tetrahydrocortisone → cortolone) eliminated its inhibitory effect, suggesting a possible orienting role of the side chain. However, since androgens were inhibitors of 11-oxoreductase, the side chain is not essential for binding to the reductase.

The magnitude of the inhibitory effects of steroid analogs differs between tissues. This is illustrated by the data of Bernal *et al.* (1980), who compared the effects of a variety of steroids on placental and decidual microsomes in the oxidative direction. They observed that testosterone, 5 α -dihydrotestosterone, and tetrahydrocortisol inhibited the decidual enzyme, but not the placental enzyme. Perhaps the two organs contain distinct species of 11-HSD.

3. Subcellular Localization

11 β -Hydroxysteroid dehydrogenase has been found in the microsomal fraction of liver (Ghrat *et al.*, 1975a; Hurlock and Talalay, 1959; Koerner, 1969; Bush *et al.*, 1968), kidney (Mahesh and Ulrich, 1960; Ghrat *et al.*, 1975a; Kobayashi *et al.*, 1987), gonads (Ghrat *et al.*, 1975a), placenta (Bernal *et al.*, 1980), spleen (Deckx and DeMoer, 1966), and lung (Nicholas and Lugg, 1982). Cytosol (post-100,000g supernatant) and mitochondria were devoid of activity, or when activity was present in these fractions, it was due to contamination with microsomal or nuclear debris. Several investigators have presented evidence for 11-HSD activity in nuclei. Ilferholzer and co-workers (Kobayashi *et al.*, 1987; Schulz *et al.*, 1987; Ilferholzer *et al.*, 1990a) confirmed the observation of Mahesh and Ulrich (1960) that kidney nuclei contain significant levels of enzyme activity. The K_m values in the experiments of Kobayashi *et al.* (1987) were 2.2×10^{-7} M (microsomes) and 2.7×10^{-7} M (nuclei), suggesting that these were identical or similar enzymes. Peterson *et al.* (1965) found 11-HSD in the microsomal and nuclear fractions of rat brain. Sakai *et al.* (1992), however, found enzyme activity exclusively in brain and pituitary nuclei. Thus, the location of 11-HSD is not limited to the endoplasmic reticulum and may be distributed between subcellular organelles in a tissue- or cell-specific manner.

4. Nucleotide Specificity

The oxidoreductases that catalyze the transformations of steroid molecules are dependent on pyridine nucleotide coenzymes. These enzymes fall into three categories. The 3 β -4-ene and 3 α -4-ene hydroxysteroid dehydrogenases (Grosso and Unger, 1964) and 21-hydroxysteroid dehydrogenase (Monder and White, 1963, 1965) have dual nucleotide specificity, utilizing either NAD or NADP as cofactors; guinea pig liver and kidney 17 β -hydroxysteroid dehydrogenases exist as particulate NAD-dependent and soluble NADP-dependent forms (Endahl *et al.*, 1960; Endahl and Kochakian, 1962); rat ovarian 20 α -hydroxysteroid dehydrogenase (Wiest and Wilcox, 1961) and liver 5 β -4-ene reductase (Tomkins and Isselbacher, 1964) are strictly NADP-dependent.

A survey of nucleotide specificity of 11- β -ISD in various tissues is presented in Table V. Rat liver enzyme has been reported to have dual nucleotide specificity, with NADP more effective than NAD (Hurluck and Talalay, 1959; Bush *et al.*, 1968; Koerner, 1966, 1969). In other tissues, including lung, kidney, placenta, intestinal mucosa, adipose tissue, striated muscle, and spleen, NADP was more effective than NAD, or NAD was not a cofactor. Two groups found that NAD and NADP were equally effective with enzyme from rat kidney (Mahesh and Ulrich, 1960) and human placenta (Meigs and Engel, 1961). Morit and Kozawaki (1962) have proposed that rat kidney contains distinct NADP- and NAD-dependent forms of 11- β -ISD. NAD was reported to be a better cofactor than NADP with 11 β -dehydrogenase from rat mandibular gland. In one study with rat liver, NADPH was a better cofactor than NADH (Bush *et al.*, 1968). The data are consistent with a heterogeneous distribution of NAD- and NADP-responsive forms of 11- β -ISD.

5. Kinetic Constants

The K_m values summarized in Table VI are taken from a variety of sources and extend over a 1000-fold range in the oxidative direction, extending from 0.1 μ M for mouse spleen microsomes (Decker and DeMoer, 1966) to 172 μ M for mouse liver microsomes (Burton, 1965). Direct comparison of the various values cannot be readily made, because of the large number of variables that must be taken into consideration: steroid substrate, pH, temperature of incubation, tissue preparation, tissue fraction, cofactor concentration. The broad variability persists even if only microsomes measured at 37°C and pH 7.4 are compared (Burton, 1965; Bernal *et al.*, 1960). The K_m values for cortisone reduction also extend over a wide range.

TABLE V
COENZYME SPECIFICITY OF 11- β -ISD

| Tissue | F \rightarrow E* | E \rightarrow F* | Citation* |
|-------------------------|---|---|------------------------------|
| Rat liver | NADP (NAD not tried) NADP > NAD NADP > NAD NADP (NADH not tried) | NADPH > NADH — — NADPH (NAD not tried) | (1) (2) (3-5) (6) |
| Rat lung | NADP (NAD inactive) | — | (4) |
| Rat kidney | NADP = NAD NADP > NAD NADP = NAD NADP > NAD | — — — — | (7) (4,8) (9) (10) |
| Human placenta | NADP (NAD inactive) NADP (NAD inactive) NADP (NAD inactive) NADP (NAD not tried) | — — — — | (11) (11) (11) (12) |
| Human intestinal mucosa | NADP > NAD | — | (13) |
| Human adipose | NAD > NADP NADP > NAD | Not tried NADPH (NADH not tried) | (14,15) (16) |
| Rat submandibular gland | NAD > NADP | — | (17) |
| Rat spleen | NADP > NAD | — | (18) |

* F, cortisole; E, cortisone.

(1) Koerner (1969); (2) Bush *et al.* (1968); (3) Hurluck and Talalay (1959); (4) Koerner (1966); (5) Koerner and Jellman (1964); (6) Nicholas and Lugg (1962); (7) Mahesh and Ulrich (1960); (8) Kobayashi *et al.* (1967); (9) Meigs and Engel (1961); (10) Olnak (1960); (11) Sweet and Byson (1960); (12) Burton and Anderson (1963); (13) Weidenfeld *et al.* (1962); (14) Hoyer and Moller (1977); (15) Ferguson and MacPhee (1976); (16) Decker and DeMoer (1966).

* NADPH had little or no effect in the reductive direction.

6. pH Optimum

The recorded values for the pH optimum of 11 β -hydroxysteroid dehydrogenase, like the kinetic constants, vary broadly when measured in the oxidative or reductive directions. Oxidation of cortisole to cortisone by microsomes of mature rat liver was optimal at about pH 10 (Bush *et al.*, 1968; Koerner, 1969). Fetal mouse liver had a reported pH optimum of 8 (Michaud and Burton, 1977). The value for salivary gland homogenate was pH 7.6 (Ferguson and MacPhee, 1976). Human placenta homogenate was reported by one laboratory to optimally oxidize cortisole in the pH range 8 to 9 (Olnak, 1960). Another laboratory reported a maximum above pH 10, with a plateau between pH 7 and 8

TABLE VI
 MICHAELIS CONSTANTS REPORTED FOR 11- β ISD IN VARIOUS TISSUES

| Tissue | Fraction* | Variable substrate† | pH | K_m (μ M) | Citation* |
|-------------------|-----------|---------------------|-----|------------------|-----------|
| Rat liver | mc | Cortisol | 7.4 | 16.4 | (1) |
| Guinea pig liver | mc | Cortisol | 7.4 | 27.1 | (1) |
| Rat liver | mc | Cortisol | 7.4 | 30 | (2) |
| Rat liver | mc | Cortisol | 8.5 | 17.6 | (3) |
| Rat liver | mc | Corticosterone | 8.5 | 0.2 | (3) |
| Rat liver | mc | Corticosterone | 8.5 | 0.22 | (4) |
| Mouse liver | mc | Cortisol | 7.4 | 172 | (5) |
| Mouse fetal liver | mt | Cortisol | 8.0 | 10 | (6) |
| Rat lung | mt | Cortisol | 7.4 | 4.1 | (7) |
| Mouse spleen | mc | Cortisol | 7.4 | 1.7 | (7) |
| Mouse spleen | mc | Corticosterone | 10 | 0.11 | (8) |
| Human adipose | hom | Cortisol | 7.2 | 0.5 | (9) |
| Human placenta | mine | Cortisol | 7.4 | 3.0 | (10) |
| Human placenta | mc | Cortisol | 7.4 | 0.3 | (11) |
| Human decidua | mc | Cortisol | 7.4 | 3.2 | (11) |

* mc, microsomal fraction; mt, nuclear fraction; mt, mitochondrial fraction; hom, homogenate.

† Constant co-substrate was NADP.

* (1) Bush *et al.* (1968); (2) Koerner and Hellman (1961); (3) Moulter and Lakshmi (1969a); (4) Murphy (1979b); (5) Burton (1966); (6) Michaud and Burton (1977); (7) Nicholae and Lugg (1982); (8) Deckx and DeMoer (1966); (9) Weidenfeld *et al.* (1982); (10) Kobayashi *et al.* (1987); (11) Bernal *et al.* (1980).

(Bernal *et al.*, 1980). Spleen microsomes were maximally effective at about pH 10 (Deckx and DeMoer, 1966). In the reverse direction, few values were available. The range was nevertheless broad, embracing values from pH 5.5 to 7.0 (Michaud and Burton, 1977; Deckx and DeMoer, 1966; Bush *et al.*, 1968).

The cause of such a wide range of values is not immediately apparent. That the method of preparation of the tissue may have played a role is suggested by data reported by Monder and Lakshmi (1989a). Freshly prepared rat liver microsomes generated a pH-activity profile with a maximum at pH 10 similar to what was reported by most investigators (Bush *et al.*, 1968; Koerner, 1969; Koerner and Hellman, 1966; Deckx and DeMoer, 1966). When briefly exposed to detergent, a profile resembling that obtained by Bernal *et al.* (1980) was obtained, with a plateau between pH 7 and 8, and a maximum at more alkaline values.

Varying conditions yielded distinctive pH-activity curves that were less a reflection of the intrinsic property of the enzyme than a composite reflection of the environment of the enzyme and its prior treatment.

D. EFFECTS OF HORMONES

1. Androgens and Estrogens

In rats, there is a sex-dependent difference in 11- β ISD of liver (Lax *et al.*, 1978, 1979) and kidney that favors males (Smith and Punder, 1991). Consistent with this observation is the strong evidence that 11- β ISD is affected by the administration or withdrawal of sex steroids. The reduction of cortisone to cortisol by male rat lung is diminished after castration (Nicholae and Lugg, 1982). Gonadectomy may decrease liver 11- β ISD in male rats as well, but appears to have no effect on female rats (Lax *et al.*, 1979). The introduction of testosterone to gonadectomized males is reported to bring the liver enzyme up to normal, whereas estradiol almost completely suppresses liver activity in male and female rats. Testosterone can increase the activity of female liver to about the level of normal male liver (Lax *et al.*, 1979). The response of different organs may not, however, be uniform. It has been reported that 11- β ISD in genital skin fibroblasts of squirrel monkey is inhibited by testosterone (Hammami and Sileri, 1990). These studies suggest that male and female steroids have opposite effects on 11- β ISD expression.

The effects of estradiol and testosterone on liver 11- β ISD of hypophysectomized rats are different from their effects on gonadectomized animals. Inhibition of activity by estradiol is suppressed, whereas testosterone raises the level of activity somewhat above normal. Hypophysectomy appears to release an endogenous suppression in females, raising the activity above that of comparable male rats. The effects of hypophysectomy are complicated, since this process eliminates numerous peptide and steroid hormones. In general, ablation of the pituitary results in loss of sex steroid dependence of liver enzymes of steroid metabolism (Gustafson and Stenberg, 1976). The effects have been attributed to growth hormone. However, no studies have yet been performed on the growth hormone dependence of 11- β ISD.

The response of neonatal rat testis to androgen and estrogen administration was similar to that seen in livers of hypophysectomized rats. Estradiol lowered 11- β ISD activity and testosterone had no effect (Thruv *et al.*, 1975b). It would therefore be expected that differences in

the level of 11- β -HSD would be seen in the two sexes in tissues that are responsive to sex steroids. The administration of the antiestrogen MER-25 to pregnant baboons prevented the increase in the capability of the placenta to oxidize cortisol to cortisone with advancing gestation, indicating that estrogen regulates the activity or synthesis of placental 11- β -HSD (Pepe and Albrecht, 1987). This observation was tested directly by increasing placental estrogen production and slowing a similar increase in the extent of oxidation of cortisol to cortisone (Pepe *et al.*, 1988; Baggia *et al.*, 1990).

In the rat kidney, the effects of gonadectomy are unclear. In one study (Ghiral *et al.*, 1975b), it was found that female animals respond to ovariectomy by developing normal male 11- β -HSD levels, whereas 11 β -HSD levels in males subjected to castration retained the activity unchanged. Female animals, therefore, hypophysectomy is equivalent to gonadectomy. In another study (Smith and Funder, 1991), the opposite was found. Gonadectomy decreased renal 11- β -HSD in males and had no effect on females.

Differences in 11- β -HSD activity in the two sexes are increased after puberty in normal rats. In liver and kidney, the female values are lower, because of the suppressive effect of estradiol. Values for the gonads, where concentrations of the sex steroids are predictably high, are consistent with those for other tissues (Hoff *et al.*, 1973).

In perfused male rat lung, castration decreased reduction of cortisone to cortisol (Nicholas and Lugg, 1982). Therefore the effects of sex steroids on 11- β -HSD are broad ranging and affect the enzyme in several organs. Adrenal 11- β -HSD measured in the oxidative direction is highest in the meadow vole during the winter, and is depressed during the breeding season, which is associated with an increase in corticosterone and increased adrenal size. It has been suggested that the circannual effect is caused by seasonal ingestion of phytoestrogens (Unger *et al.*, 1978).

2. Corticosteroids

Glucocorticoids may intervene in their own metabolism by influencing the activity of 11- β -HSD. Some indirect suggestion that glucocorticoids affect lung 11- β -HSD is based on the observation that stress increases the activity of rat lung 11- β -HSD in the reductive direction (Nicholas and Lugg, 1982). Cortisol, which regulates the mitosis, maturation, and mortality of lymphocytes, also affects the level and direction of their 11- β -HSD activity. Thymic cells of mice pretreated with cortisol for 9 days showed increased activity in the oxidative direction

and no change in the reductive direction (Dougherty *et al.*, 1960). Placental 11- β -HSD of the baboon may be resistant to corticosteroid hormones. Serum cortisone did not alter the level of placental 11- β -HSD activity and possibly decreased 11 β -dehydrogenase (Pepe and Albrecht, 1985a). Although it is unlikely that the effect is due to direct inhibition by cortisone of enzyme activity, the addition of pregnenolone (250 nM), progesterone (25 nM) or cortisone (250 nM) to human or baboon placental homogenates inhibited oxidation of cortisol to cortisone (Pepe and Albrecht, 1984a). Bernal *et al.* (1982) found no changes in human placental 11- β -HSD taken after elective cesarean section, or after spontaneous and induced labor. However, injections of dexamethasone into pregnant rhesus monkeys increased cortisol-to-cortisone conversion by the placenta (Althaus *et al.*, 1982).

Isolated perfused fetal rabbit lung oxidized cortisol, but the reverse reaction was minimal (Torday *et al.*, 1976). Lugg and Nicholas (1978) found that the development of 11- β -HSD in the lung of the fetal rabbit *in vivo* is not affected by dexamethasone after direct injection into the fetus. Fetal rat lung, however, responded to betamethasone, another glucocorticoid analog, with an increase in 11- β -HSD reduction. *In vitro*, cortisol stimulated growth of fetal human lung cells. This corresponded with increased net conversion of cortisone to cortisol (Smith *et al.*, 1973). Abramovitz *et al.* (1982) showed that fetal lung cells diverged during growth in tissue culture into populations of epithelial cells and fibroblast-like cells which could be cultured separately. The former preferentially oxidized cortisol to cortisone; the latter, which was the dominant surviving cell type, reduced cortisone to cortisol. These findings may explain an apparent contradiction between findings of Smith *et al.* (1973) described above and those of Pasqualini *et al.* (1970a) and Murphy (1978), who found that fetal lung primarily catalyzed cortisol oxidation, as did all other fetal tissues. Reduction of cortisone increased with lung maturity. This developmental pattern also applies for the fetal rat (Smith, 1978). The reduction of cortisone by fetal rabbit lung was reported by Ginnopoulos (1974). Mature rat lung rapidly reduced cortisone to cortisol (Nicholas and Kim, 1976). The suggestion that the direction of 11- β -HSD metabolism in lung is determined under physiological conditions by the nature of the cell population remains to be tested.

3. Thyroid

There have been several studies published on the effects of thyroid hormone on 11- β -HSD. Species specifically has been observed on the effects of thyroxine on the oxidation of cortisol to cortisone by liver.

Thyroxine administration causes a decrease in male rat liver (Koerner and Hellman, 1964; Lax *et al.*, 1979) that is only apparent after 7 days of continuous exposure. In contrast, hyperthyroid humans respond with increased hepatic activity (Zunoff *et al.*, 1983; Hellman *et al.*, 1961; Gordon and Southren, 1977). Thyroidectomy or hypothyroidism reverses the response, resulting in increased activity in the rat (Koerner and Hellman, 1964) and decreased activity in humans (Zunoff *et al.*, 1983; Hellman *et al.*, 1961; Gordon and Southren, 1977). The effects of thyroid hormones are tissue specific. No changes occur in the kidney (Koerner and Hellman, 1964; Lax *et al.*, 1979; Smith and Funder, 1991) or in the reticuloendothelial system (Dougherty *et al.*, 1960). The proposal that thyroid hormones act by controlling the availability of pyridine nucleotides (Dougherty *et al.*, 1960) appears not to be borne out by experiment (Koerner and Hellman, 1964). Enzyme inhibitors are not formed (Koerner and Hellman, 1964). Whether thyroid hormones directly affect the level of enzyme is not known. It has been suggested that thyroid hormone controls the level of available testosterone, and thus indirectly influences 11- β ISD.

4. Other Hormones

The activity of placental 11- β ISD, which shows activity almost entirely in the oxidative direction, is not affected by prolactin, hCG, or ACTH *in vitro*. The cortisol and cortisone content of amniotic fluid of diabetic and nondiabetic women are identical. Therefore insulin, glucagon, and the various diabetogenic factors do not influence 11- β ISD (Baird and Bush, 1960).

III. Developmental Biology and 11- β ISD

A. Fetal Development

1. Placental 11- β ISD

The level of active corticosteroid to which the fetus is exposed is crucial to its development and maturation. Too high exposure can lead to developmental disturbances. The placenta catalyzes the oxidation of the 11 β -hydroxy groups of corticosteroids, both natural (Dutton and Jeyes, 1968; Bernal *et al.*, 1981; Giannopoulos *et al.*, 1982; Pasqualini *et al.*, 1970a; Waddell *et al.*, 1988) and synthetic (Loytitz *et al.*, 1978), and thus provides a barrier to the transfer of active glucocorticoid to the fetus by converting the steroids to the biologically inactive

11-oxo form. In keeping with this role, reduction of 11-oxosteroids by placental 11- β ISD is low or not detectable (Osinicki, 1960; Bernal *et al.*, 1980; Murphy *et al.*, 1974; Murphy, 1978b; Kittinger, 1974). As a consequence of this overwhelming oxidative activity, relatively high proportions of 11-oxocorticosteroid metabolites appear in normal amniotic fluid and fetal cord serum (Osinicki, 1960; Baird and Bush, 1960; Brothmann *et al.*, 1962). When the placental barrier is bypassed, intra-amniotically administered cortisol is absorbed by the human fetus and is oxidized in individual organs slowly, resulting in long-term retention of active steroid (Murphy and Vedady, 1982). The proportion of cortisol relative to cortisone is greater in the amniotic fluid than the cord fluid and rises with gestation in humans (Murphy, 1977a). Chorionic membrane catalyzes a reduction of cortisone to cortisol (Murphy, 1977b; Bernal *et al.*, 1980) and may contribute to the rise in active steroid. Although the direction of metabolism of the 11-oxygen by placenta and chorion does not change during gestation, it is not known whether the levels of 11- β ISD activity in these organs change. The few studies that have been performed with human placenta have yielded conflicting results. Giannopoulos *et al.* (1982) have reported quantitative changes in the capacity of placenta and decidua to interconvert cortisol and cortisone; Bernal *et al.* (1982) find no changes in placental corticosteroid metabolism during the terminal stages of pregnancy in the human, but find changes over the longer term. These results may not in fact be contradictory since the span of gestation studied in the two investigations were quite different, the former extending through the major part of pregnancy, and the latter considering only the terminal stages. Tanawell *et al.* (1977) have suggested that reductive capacity of the chorion is valuable for the fetus, because it represents a mechanism for regenerating cortisol for the fetus, possibly by acting as an accessory adrenal gland.

2. The Feto-Placental Unit

The behavior of the placenta *in vitro* confirms that a highly effective barrier exists against the transfer of 11 β -hydroxysteroids from mother to fetus. The ability of the human and primate feto-placental unit to efficiently oxidize cortisol to cortisone results in the transfer of little or no cortisol into the fetus (Althaus *et al.*, 1982) who is thus protected against the teratogenic actions of cortisol (Murphy *et al.*, 1974; Munk and Leung, 1977; Slikker *et al.*, 1982). The existence of this barrier also permits the fetus to retain autonomy over its own cortisol production (Murphy and Branchaud, 1983; Bellina *et al.*, 1972; Mitchell *et al.*, 1981, 1982). The timing of the increase in active corticosteroid level in

the maturing fetus is essential for creating an internal environment compatible with independent life (Murphy, 1977a). Synthetic steroids that are poor substrates for 11-11SD, such as dexamethasone, are transferred to the fetus largely unoxidized (Althaus *et al.*, 1982; Furkenhouser *et al.*, 1978; Anderson *et al.*, 1979). This process may have important pharmacological implications. Pope and co-workers have studied the transplacental regulation of cortisol metabolism in pregnant old world monkeys. They have suggested that transuteroplacental corticosteroid metabolism may play an important role in the maturation of the pituitary-adrenocortical axis (Pope and Albrecht, 1985b). Although transfer of cortisone (from maternal cortisol) to the fetus is extensive, little cortisone is converted to the active hormone (Mitchell *et al.*, 1982). Most of the cortisol available to the late gestation fetus, as illustrated with baboon and rhesus, is endogenous (Mitchell *et al.*, 1981; Althaus *et al.*, 1982; Pepe and Albrecht, 1984b), resulting from secretion of the maturing fetal adrenal.

3. Fetal 11-11SD

Fetal tissues contribute to the net oxidation of corticosteroids in the developing organism. Brain, gut, liver, and lung in the fetal mouse are all strongly oxidizing at 14 days of gestation. By 19 days, with birth approaching, the tissues show increasing capacity for reduction (Tye and Burton, 1980), in some cases evolving from net oxidation to net reduction. The capacity of the mouse liver to catalyze net reduction continues to increase after birth (Burton and Jeyes, 1968).

Other organs change their relative preference of direction of 11-oxoreduction during development. The nonpregnant human uterus preferentially oxidizes cortisol to cortisone, but catalyzes the reverse process during early pregnancy (Murphy, 1977b). The net effect of the metabolic events catalyzed by 11-11SD in the fetus is the oxidation of cortisol to cortisone or corticosterone to 11-dehydrocorticosterone (Murphy, 1981; Pasqualini *et al.*, 1970a,b). The proportion of oxidized to reduced form decreases during gestation as 11-11SD in the liver (Michaud and Burton, 1977; Smith *et al.*, 1982) and lung (Micheline and Lugg, 1982; Smith *et al.*, 1982; Smith, 1978) plays an increasingly important role in reducing the 11-oxo group of the steroid.

The changes in steroid oxidoreduction in the individual fetal organs are intimately connected with the maturational events that prepare the organism for birth and permit its subsequent independent existence (Lugg, 1976). Fetal lung has been the subject of intense study. Pulmonary differentiation is dependent on and accelerated by 11 β -hydroxylated corticoids. Glucocorticoids induce surfactant and enhance

of surfactant and the differentiation of alveolar cells (Avery, 1975). Steroid effects on lung maturation have important clinical applications in the prevention or reversal of hyaline membrane disease (Liggins and Howie, 1972). Since the dominant metabolic transformation of glucocorticoids in the fetus is 11-oxidation, the ability of the lung to catalyze 11-oxoreduction is of particular significance. Conversion of cortisone or 11-dehydrocorticosterone to their respective 11-reduced forms is essential for lung differentiation (Torday, 1980; Gianopoulos, 1974; Murphy, 1981; Torday *et al.*, 1976; Drafa *et al.*, 1976) based on the criteria of lung size (Drafa *et al.*, 1975), cell growth (Torday, 1980; Smith *et al.*, 1973), phosphatidylcholine production (Drafa *et al.*, 1975; Torday *et al.*, 1975), and glycogen content (Smith *et al.*, 1982). The ability of fetal lung to reduce 11-dehydrocorticosteroids increases during gestation in rabbit (Dougherty *et al.*, 1980; Torday *et al.*, 1976; but see Hummelink and Ballard, 1986), human (Smith *et al.*, 1973), mouse (Burton and Turnell, 1968), and rat (Smith *et al.*, 1982). It is possible that in human lung the increase in reductive ability during gestation may in part be due to a large decrease in the 11 β -dehydrogenase as well as an absolute increase in 11-oxoreductase. It has also been suggested that the driving force in the increase in 11-oxoreduction is an increase in NADPH (Torday *et al.*, 1976).

During the second trimester and early third trimester of pregnancy, 11-oxo steroids exceed 11-hydroxysteroids in the fetal circulation (Waddell *et al.*, 1988; Murphy and DiezAñux, 1972; Burton and Jeyes, 1968; Sowell *et al.*, 1971). Murphy *et al.* have documented the extensive catabolism of cortisol to cortisone in the human mid-term fetus (Murphy, 1977b, 1981; Murphy and Branchaud, 1983). The magnitude of oxidation of cortisol and corticosterone is dependent on the combined metabolic actions of the placenta, its associated membranes, and the fetal tissues. The relative oxidative and reductive activities in many tissues change with time. In most tissues, irrespective of species, the oxidation of steroids dominates at mid-gestation. In late gestation, reductive activity is expressed in some tissues. It is not yet known whether the shifts in dehydrogenase-oxoreductase capabilities of some tissues are species specific, nor is it known for most organs when, during development, the expression of 11-11SD activity first appears. Perinatal reduction is dominated by the lung and liver. In mouse liver net 11-reduction continues to increase after birth. The relative capabilities of each tissue to catalyze 11-oxidation or 11-reduction relate well with the proportion of 11-oxo- to 11-hydroxysteroids in these tissues (Smith, 1978; Smith *et al.*, 1982). The ratio of reduced to oxidized steroid in tissues at critical stages of development may pro-

vide important clues to determining key molecular events necessary to complete development. There is no doubt that the presence of the physiologically appropriate level of active corticosteroid at the correct stage of development of fetal organs is essential for their proper development and function. In view of this and the known teratogenic effects of corticosteroids when administered in excess to experimental animals, there is ample reason to suspect that 11-HSD performs a crucial mediating role in fetal development.

B. POSTNATAL DEVELOPMENT

1. *In Vivo* Metabolism of Corticosteroids

After birth, overall corticosteroid metabolism at C-11 is reductive. In some organs, such as uterus, parotid gland, colon, and kidney, metabolism continues to be predominantly oxidative. One consequence of the concurrent selective exposure of steroids to oxidative or reductive conditions in the various organs is the excretion of a mix of 11-oxo- and 11 β -hydroxysteroid metabolites. In humans and primates, who excrete corticosteroid metabolites mainly by way of the kidney, measurement of urinary steroid metabolites provides an accurate reflection of the oxidoreductive balance. In other organisms, such as rats and mice, that utilize the gastrointestinal tract as the dominant excretory pathway for steroids, establishing the net balance of oxidation and reduction is far more difficult, and has not yet been successfully accomplished.

One approach to the study of murine steroid metabolism utilizes biliary steroids. In rats, about 90% of corticosterone metabolites are recovered from bile (Gustafsson and Gustafsson, 1974), reflecting primarily hepatic metabolism (Eriksson and Gustafsson, 1971). Most identified metabolites contain the 11 β -hydroxy group, suggesting that liver metabolism at C-11 is primarily reductive *in vivo*.

In humans, the metabolites of endogenously produced cortisol are excreted into the urine as a mixture of products at different levels of reduction and oxidation (Peterson *et al.*, 1955). These include metabolites reduced in ring A (tetrahydrocortisol, tetrahydrocortisone) and ring A-reduced metabolites in which reduction at C-20 (cortisol, cortisone) of those metabolites in which oxidation dominates, the major examples are the cortic acids, C₂₁ steroids containing a carboxylic acid group at C-21 (Monder and Bradlow, 1980). There are additionally significant amounts of metabolites resulting from the loss of the ketol side chain, and a number of minor metabolites.

2. Corticosteroid Metabolites in Health and Disease

Cortisol in normal humans, male and female, has a biological half-life of 95 to 130 min (Peterson *et al.*, 1955). Cortisone has an average plasma biological half-life of 28 min. This is due in part to its rapid conversion to cortisol, and in part to the greater susceptibility of cortisone to catabolism. Its reduction to cortisol, mediated by 11-HSD, is the basis for its pharmacological action, the leaking away of cortisone by competing catabolic reactions is the reason that cortisone is a less potent pharmacological agent than cortisol. That cortisol is oxidized at C-11, i.e., that the oxidation-reduction process is physiologically irreversible, is revealed by the profile of urinary metabolites. The ratio of the major metabolic products tetrahydrocortisol (THF), al-tetrahydrocortisol (5 α -THF, ATHF) and tetrahydrocortisone (THE), expressed as (THF + ATHF)/(THE) = *R*, has been used as a measure of the physiological oxidoreductase activity at C-11. Numerous studies have confirmed that alterations in physiological status cause significant changes in this ratio. Table VII presents a qualitative assessment of the effects of a variety of conditions on the value of *R* relative to that of normal subjects, whose values range from 0.5 to 2. Irrespective of the nature of the illness, the proportion of 11-reduced metabolites increased relative to control (presumably normal) populations, with few exceptions. The changes were not large, rarely exceeding 50%, and are insufficiently great in magnitude to distinguish whether changes in the level of enzyme or pyridine nucleotide are rate limiting, or whether the difference can be attributed to a selective redistribution of 11-oxosteroids between tetrahydro and penta-hydro metabolites (Bradlow *et al.*, 1968; Zumoff *et al.*, 1968b). Zumoff *et al.* have shown that while (THF + ATHF)/(THE) increased in cirrhosis (Zumoff *et al.*, 1967) the total value of C-11 hydroxy/C-11-oxo, including all urinary steroids, was unchanged from normal (Zumoff *et al.*, 1968a), suggesting increased conversion of THE to cortisone. Where the secretion of cortisol from the adrenal results in peripheral accumulation great enough to exceed the ability of the organism to dehydrogenate at C-11, the urinary cortisol/cortisone and THF/THE ratios increase (Baulieu and Joly, 1957).

During postnatal development, the *R* values change from ca. 0.1 at birth to approximately unity, as Fig. 3 shows. The early low values of this ratio are the consequence of the fact that in the recently born infant, 11-dehydrogenation is highly active, resulting in the excretion of THE, but little THF. This pattern also occurs in primates other than human (Pope and Townsley, 1976). The proportion of THE and THF

TABLE VII
EFFECT OF DISEASE ON THE PROPORTION OF URINARY 11 β -HYDROXY
TO 11-OXO METABOLITES

| Condition | Effect* | Citation |
|--------------------------------|-------------|----------|
| Cushing's disease or ACTH | $R_E > R_N$ | (1, 5) |
| Infection, nonspecific illness | $R_E > R_N$ | (6, 7) |
| Rheumatic disorders | $R_E > R_N$ | (7, 8) |
| Cirrhosis | $R_E > R_N$ | (9) |
| Essential hypertension | $R_E > R_N$ | (10, 11) |
| Chronic myelogenous leukemia | $R_E = R_N$ | (12) |
| Adrenal carcinoma | $R_E > R_N$ | (13) |
| Schizophrenia | $R_E = R_N$ | (14) |
| Hypothyroid | $R_E > R_N$ | (15) |
| Endogenous depression | $R_E < R_N$ | (16) |
| Chronic renal failure | $R_E > R_N$ | (17) |
| Anorexia nervosa | $R_E < R_N$ | (18, 19) |

* $R = (TTIF + ATIIH)/TIIH$; tetrahydrocortisol + allopregnanolone/tetrahydrocortisol; R_E = subjects with designated condition; R_N = normal or control subjects.

(1) Gray *et al.* (1962); (2) Bailey and West (1969); (3) Peterson and Pierce (1960); (4) Bush and Willoughby (1957); (5) Kornel (1970); (6) Zimmelf *et al.* (1974); (7) Ichikawa (1966); (8) Pol (1967); (9) Zimmelf *et al.* (1967); (10) Kornel *et al.* (1968); (11) Walker *et al.* (1991); (12) Gahlagher *et al.* (1965); (13) Fukushima *et al.* (1960); (14) Lommo *et al.* (1957); (15) Hellman *et al.* (1981); (16) Murphy (1991); (17) Walker and Schwartz (1991); (18) Verhulst *et al.* (1990); (19) Verhulst *et al.* (1979).

shifts to the dominant postnatal ratio of 1-2 during the first year of life (Panilascu-Goldinberg and Giroud, 1974; Savage *et al.*, 1975; Blumck, 1968; Kraan *et al.*, 1980). C. H. L. Shuckleton, personal communication. The relationships between F and E in serum and amniotic fluid during the last trimester of pregnancy are similar to those of TIIH and TIII (Nouin *et al.*, 1991). So strong is the oxidation pressure in infants, that the blood F/E ratio will remain < 1 even after intravenous administration of high concentrations of cortisol (maternal F/E = 11) (Buus *et al.*, 1966). The change in the 11-hydroxysteroid/11-oxosteroid ratio during early development is in accord with the changes in the increasing ability of 11-HSD to catalyze 11-reduction relative to 11-oxidation.

No data are available for the prenatal metabolism of corticosterone in humans. At the earliest known age examined, 1 year, the value for (TIIH + ATIIH)/TIIA indicated a strong preference for the reduced




FIG. 3 The effect of age on the value of $R = (TIIH + 6n TIIH)/TIIH$. Modified from Mondri and Shuckleton (1984).

forms, TIIH and alloTIIH (Peterson and Pierce, 1960; Savage *et al.*, 1975; Kornel *et al.*, 1969; Blumck, 1968). At every age, R_E exceeded R_N by two- to eightfold. The values for R_E fluctuated over a wide range between laboratories, and so from the limited data available, it is not possible to draw conclusions about age-related trends.

3. C-11 Metabolism in Specific Organs

Continuing the prenatal trend, 11-HSD increases in several organs during early postnatal development in the mouse (de Moor and Deck, 1966) and rat (Hoff *et al.*, 1973), then decreases. In the mouse, 11-HSD, measured as oxidation of corticosterone at p11 10.5, rises from birth to 10 weeks of age in spleen, kidney, and liver, suggesting that it is due to some coordinated process, then declines to intermediate values. Development in the rat liver is qualitatively similar. Maximum value of cortisol oxidation occurs at 30 days of age, followed by a decline. Thus, in all species, alterations in the interconversion of corticosteroids at C-11 initiated in the fetus continue after birth, each organ following a unique pattern (Mitchell *et al.*, 1981; Pepe, 1979; Krozowski *et al.*, 1990; Moisin *et al.*, 1992).

IV. Are 11-Dehydrocorticosteroids Biologically Active?

Cortisone is converted faster than cortisol to inactive metabolites: it binds poorly to glucocorticoid receptors (G. G. Rousseau *et al.*, 1972) under optimal experimental conditions and probably not at all under physiological conditions. On the basis of these observations it would not be predicted that 11-dehydrocorticosteroids have significant bio-

logical activity. There are, however, several observations that suggest otherwise. Barzeghan *et al.* (1982) reported that cortisone, but not cortisol, strongly inhibited glucose- and arginine-induced insulin secretion. Berliner and Ruhmann (1966) found that though cortisol inhibited fibroblast growth in tissue culture, cortisone increased growth. In guinea pig isolated ileum, cortisone antagonized GABA_A receptor-mediated contractile responses to applied GABA, with the enhancing effect of cortisol (Ong *et al.*, 1990). Souness and Morris (1990) have found that 11-dehydrocorticosterone caused significant kaliuresis in male adrenalectomized rats and blocked the sodium ion retention caused by aldosterone. Souness and Morris (1991) have also found that, though 11-dehydrocorticosterone was not reduced to corticosterone by rod bladder preparations, the steroid nevertheless blocked aldosterone-stimulated short-circuit current.

There is one report, as yet unconfirmed, that 11-dehydrocorticosteroids have intracellular activity, modulating salt metabolism in the nasal gland of the domestic duck (Sandor *et al.*, 1983). The molecule basis for this observation is obscure. Membrane-associated events may explain the effects of cortisone on ileum contractile responses (Ong *et al.*, 1990), and possibly on other cortisone-mediated processes. The number of reports of 11-dehydrocorticosteroid activity are few, and none have been independently confirmed. Nevertheless, the examples cited support the possibility that oxidation of corticosteroid at C-11 may not be exclusively inactivating, and may generate physiologically significant metabolites.

V. 11-HSD IN LOWER VERTEBRATES

The presence of 11 β -hydroxy- and 11-oxosteroids in animals as diverse as fish (Chan and Yeung, 1989; Gottfried, 1964), birds (Holmes *et al.*, 1974), and the platypus (McDonald *et al.*, 1988) indicates that 11-HSD serves an important function in nonmammalian vertebrates. Both cortisol and cortisone are found in salmon blood (Idler *et al.*, 1959a,b) and though oxidation may occur, cortisone does not appear to be effectively reduced to cortisol (Idler and Truscott, 1963). These conversions are probably extrahepatic, possibly occurring in the anterior kidney (Columbo and Bern, 1970), since in a wide range of bony fish there was no evidence of liver 11-HSD (Columbo *et al.*, 1972; Monier and Lakkhimi, 1990a). However, generalizations about the role a particular organ in fish may play in 11-oxidoreduction must be qualified. For example, in the ganoid fish, *Amia calva*, the anterior kidney was deficient in 11-HSD, unlike that of another ganoid, *Lepisosteus osseus*

(Columbo *et al.*, 1972). The appearance in bile of cortisone and tetrahydrocorticosterone after injection of cortisol into trout (Truscott, 1979) or salmon (Donaldson and Fagerlund, 1972) is consistent with an active hepatic 11-HSD (Kime, 1978) in some species of teleost. The resolution of the source of 11-oxidoreduction in fish is of additional importance because C-11 steroids may be the source of the teleost androgens 11 β -hydroxy- and 11-oxotestosterone (Idler and MacNab, 1967; Lelitz and Reinbold, 1987; Rosenblum *et al.*, 1985). The somewhat more advanced African lung fish *Protopterus*, in contrast, is incapable of oxidizing the C-11 hydroxy group (Idler *et al.*, 1972). Early work on the occurrence of 11-oxygenated steroids in lower vertebrates have been summarized by Gottfried (1964).

Direct measurement of 11-HSD in livers of vertebrates has been made by Mondur and Lakkhimi (1989a). No 11 β -dehydrogenase was detected in the liver microsomes of the frog, toad, mud puppy, shark, and several birds. In contrast, all mammals had activity. In the redutive direction, activity was present only in the livers of dogfish, birds, and mammals. Amphibians and teleosts had no detectable enzyme.

The duck nasal gland provides an interesting example of a system in which 11-oxidation may activate a steroid. Marina birds have a specialized organ, the nasal salt-gland, which protects them against the high salinity of ingested sea water. These glands concentrate and excrete the excess salt by a mechanism that is corticosteroid dependent. The endogenous corticosteroid, corticosterone, is rapidly oxidized to 11-dehydrocorticosterone by the nasal gland *in vivo*, *in vitro*, or by cell-free homogenates of the gland (Takemoto *et al.*, 1976; Sandor *et al.*, 1977). The glucocorticoid receptor, or an enzyme closely associated with the receptor, converts the specifically bound corticosterone to 11-dehydrocorticosterone (Sandor *et al.*, 1977, 1983; Sandor and Mehdi, 1980), which is transported to the nucleus. It is proposed that the receptor binds corticosterone, and its activation requires oxidation of the steroid at C-11.

VI. THE FORMS OF 11-HSD EXPRESSION: UNIQUENESS OR MULTIPLICITY?

A. ON THE QUESTION OF REVERSIBILITY

The evidence presented thus far provides us with a picture of a functionally highly flexible enzyme, capable of adapting to net oxidation or net reduction depending on changing circumstances of age, health, state of gestation, and hormonal status. This remarkable

adaptive process engages the whole animal, and every organ within in unique and distinctive ways. The range of these processes is determined to some degree by genetic endowment (Nguyen-Tong-Thien *et al.*, 1971). However, within these proscribed limits, the ability of the enzyme to respond to changing conditions is so striking that a closer look at it is justified. How is it possible for a single, presumably well-characterized enzyme to express itself as a net dehydrogenase under some conditions, and as a reductase under others?

A number of mechanisms have been proposed to account for the differential behavior of 11- β -HSD in various tissues and the changes in level of activity and in the directional characteristics of oxidorreduction that occur during development. Several rely on the properties of 11-HSD as a reversible pyridine nucleotide dependent oxidorreductase. Nicholae and Lugg (1982) and Torday *et al.* (1976) have postulated that the changing NADP/NADPH ratio is the driving force that determines the relative proportion of 11 β -hydroxy- to 11-oxosteroid in lung in response to stress, castration, or adrenalectomy. Dougherty *et al.* (1960) utilized a similar mechanism to explain the appearance in immature lymphocytes of increased net 11-oxoreduction of cortisol subsequent to the introduction of triiodothyronine.

Other investigators have presented evidence that tissue specific changes in 11-HSD activities are not determined by the oxidation-reduction state of the tissue (Bernal *et al.*, 1980; Bernal and Turnbull, 1986) since they occur when nucleotide cofactors are not rate limiting. The effects of thyroxine in the rat persisted when pyridine nucleotide cofactors were not limiting (Zunoff *et al.*, 1983; Hellman *et al.*, 1961; Koerner and Hellman, 1964).

Product inhibition has been shown not to be responsible for the divergent effects of 11-HSD. Cortisol, even at 100-fold excess did not inhibit 11-oxoreductase activity (Bernal *et al.*, 1980). In the oxidative direction, neither NADPH nor 11-dehydrocorticosterone inhibited rat liver 11-HSD (Monder, 1991a).

An alternative hypothesis based on environmental perturbations is that the equilibrium ratio depends on pH. Changes in pH can theoretically affect the corticosteroid-11-dehydrocorticosteroid ratio, since the equilibrium of the overall redox reaction is dependent on the concentration of protons. The implementation of this hypothesis depends on knowledge of the equilibrium constant of the reaction and the local pH in the environment of the enzyme. Both are unknown. Changes of pH in the physiological range are not large enough to affect the 11-hydroxy/11-oxo ratio to a major degree (Alkalini and Monder, 1985; Monder and Shackleton, 1984). Any large local pH change that would persist for a sufficiently long time to alter the

direction or magnitude of 11-HSD catalyzed reaction would adversely affect other processes in the endoplasmic reticulum.

Of the "environmental" hypotheses, the most likely mechanism accounting for small changes in oxidation-reduction properties is the one that proposes changes in the ratio of pyridine nucleotide cofactors. This may be an occasional mechanism for rapid, local perturbations in 11-oxoreduction. However, it can be readily shown that the ratios of reduced to oxidized pyridine nucleotides [NADPH/NADP] would have to be unrealistically high or low in order to account for the apparent extreme values of 11 β -hydroxy/11-oxo in many tissues during development. Under physiological conditions, where the changes in 11-hydroxy/11-oxo in the whole organism based on urinary excretion patterns are not large, the relative proportion of reduced-to-oxidized cofactors may contribute to the behavior of the steroid at C-11. However, so many reactions occur in which both steroids and other biological substances require pyridine nucleotide cofactors within a cell, that it is not often that a circumstance arises where the machinery of the cell becomes entirely subservient to the metabolic requirements of a single molecule for any finite interval of time. Therefore, it is metabolically more likely and less disruptive for the cell to have developed other more specific mechanisms to control the interconversion of corticosteroids at C-11.

The possibility that the diverse behavior of 11-HSD in tissues is due to distinct, though related, enzymes has been considered by a number of investigators. In general, the view expressed has been that variants of 11-HSD are present in different tissues, representing forms with distinct kinetic properties that express behavior favoring reduction or oxidation. A model for this kind of system is glyceraldehyde phosphate dehydrogenase, in which different isozymes dominate in various tissues, and which have structural characteristics that lead to its preferential reduction to triose phosphate or oxidation to dihydroxyacetate (Kuprian, 1968). Thus the placental and decidual 11-HSD may be isozymes (Bernal *et al.*, 1980) as may also be true of the lung (Nicholae and Lugg, 1982) and liver (Bush and Mahesh, 1959b) enzymes.

B. CHARACTERISTICS OF MITOCHONDRIAL 11-HSD

1. Latency

The catalytic activity of the 11 β -dehydrogenase component of 11-HSD is not fully expressed in liver microsomal preparations. Treatments that disrupt or alter the structure of the microsomal matrix,

such as phospholipase, detergent, and elevated pH, release latent enzyme activity. These processes, by altering membrane integrity, probably make the active site of 11 β -dehydrogenase more accessible to its substrate (Gunderson and Nordlie, 1976). Latency of 11 β -dehydrogenase, first observed in rat liver microsomes (Lakshmi and Monder, 1985b), occurs in the livers of other species as well (Monder and Lakshmi, 1985a). This phenomenon is not unique. Other membrane-based enzymes express latency (Gunderson and Nordlie, 1976; Stetten and Burnett, 1967; Ernster and Jones, 1962; Schulze and Speth, 1980). It is possible that this property is a physiologically significant mechanism for controlling the expression of enzyme activity. In rat liver microsomal preparations, 11-oxoreductase activity is initially fully expressed without the intervention of latency releasing conditions (Lakshmi and Monder, 1985b). The latency behavior of hepatic 11 β -dehydrogenase and 11-oxoreductase are therefore different.

2. Energy of Activation

The temperature dependence of enzyme activity can reveal much about the environment of the enzyme. The relationship of temperature and enzyme activity has been shown to adhere to thermodynamic principles and reflect the environment of the enzyme. The energy of activation is discrete in a homogeneous environment. If, however, the environment shows discontinuities, the energy of activation of an enzymatic process will show corresponding discontinuities should the activity be dependent on the structure of that environment (Raison *et al.*, 1971; Kumamoto *et al.*, 1971).

The energy of activation (E_a) of microsome-bound 11-dehydrogenase is continuous over the entire physiological temperature range and has the same value as the soluble enzyme. In contrast, microsome 11-oxoreductase shows a discontinuity in E_a at 23°C, which is no longer present when the enzyme is solubilized, or when the microsomal lipid matrix is disrupted with phospholipases. The discontinuity coincides with a phase change in the matrix structure.

The differences between reductase and dehydrogenase with respect to latency and activation energy indicate that both activities are in distinct environments within the microsomal membrane. When they are solubilized, these environmental differences are removed, and the behavior of the enzymes reflect this (Lakshmi and Monder, 1985b).

3. Enzyme Stability

The relative stabilities of 11 β -dehydrogenase and 11-oxoreductase provides an additional distinguishing criterion. Oxidation is the more

stable activity. With freshly prepared rat liver microsomal preparations, oxidation proceeds undiminished for at least 2 h at 37°C, whereas reductase is inactivated within 10 min.

We conclude from the literature and our own observations that there is sufficiently wide diversity in the properties and behavior of 11- β SD derived from different sources to be suggestive of multiple enzyme forms. The physicochemical and kinetic characteristics of the enzyme are also consistent with independent 11 β -dehydrogenase and 11-oxoreductase sites. To address this question of enzyme multiplicity, it is necessary to review the molecular properties of 11- β SD. First, however, we will examine how the clinical evidence contributes to our understanding of the properties and physiological functions.

VII. CLINICAL STUDIES

A. 11 β -Dehydrogenase Deficiency

1. Apparent Mineralocorticoid Excess

In humans, an "experiment of nature" has provided insight into the probable function of 11- β SD in at least one organ, the kidney. A disease apparently unique to children was described in the 1970s with a clinical picture consisting of low renin activity, low aldosterone production, hypokalemia, and severe hypertension (Ullick *et al.*, 1977; Werder *et al.*, 1975; Ramirez *et al.*, 1979; Winter and McKenzie, 1977). The first completely described patient with this condition was a Zuni Indian girl in whom the diagnosis was made at the age of 3 years (New and Levine, 1977; New *et al.*, 1977). Urinary cortisol and deoxycorticosterone metabolites were below normal and were not increased after ACTH stimulation. Glucocorticoid administration exacerbated the hypertension, suggesting that it was in some manner linked to endogenous cortisol.

Extreme sodium deprivation lowered blood pressure, possibly by stimulating the rate of conversion of corticosterone to aldosterone (Hanning *et al.*, 1970). High doses of spironolactone, a potassium-sparing diuretic that acts via blockade of the mineralocorticoid receptor, also produced normalization of blood pressure, and on this regimen, plasma renin activity rose. Substitution of triamterene, a potassium-sparing diuretic that does not affect the mineralocorticoid receptor, failed to ameliorate blood pressure.

In vivo measurement of transcutaneous electrical potential difference

In the patient was consistent with mineralocorticoid effect seen in patients with primary hyperaldosteronism; the potential difference was increased with hydrocortisone administration, and diminished with spironolactone administration (New *et al.*, 1982). Sensitivity of glucocorticoid receptors was normal as assayed in lymphocytes (Bigger *et al.*, 1972). Bioassays performed to demonstrate the presence of a steroid hormone in the patient's serum capable of causing sodium retention revealed no mineralocorticoid effect (Marver and Edelman, 1978; Blair-West *et al.*, 1962; Sennett *et al.*, 1976; Adam *et al.*, 1978; Baxter *et al.*, 1976). Because the symptoms and response to treatment were consistent with aldosteronism, despite the low circulating levels of the steroid, the syndrome was referred to as "apparent mineralocorticoid excess" (AME). This designation appears to have gained general acceptance.

Patients with AME have shown (a) low rate of cortisol turnover, with approximately twice the disappearance time of radiolabeled tracer steroid compared with that of a normal subject; (b) low peripheral plasma ACTH levels; (c) normal CBG concentration; (d) greatly diminished level of urinary metabolites of cortisone compared with those of cortisol; (e) no production of tritiated water after infusion of 11-⁴tritiated cortisol, suggesting a defect in the oxidative component of the 11 β -hydroxysteroid dehydrogenase (Ulick *et al.*, 1979); (f) normal metabolism of cortisone to cortisol, suggesting an intact reductive component of 11 β -hydroxysteroid dehydrogenase (Ulick *et al.*, 1979; Mondler *et al.*, 1986); and (g) an abnormal increase in the far relative to 5 β -metabolites of cortisol (Ulick *et al.*, 1977).

Selective glucocorticoid receptor (GR) blockade with RU 38486 (RU 486) did not decrease blood pressure as would be expected if the GR were responsible for the development of hypertension in AME. Instead, a significant increase in mean blood pressure was observed compared with the pretreatment period, indicating that the GR was not contributing to the development of hypertension.

The constellation of clinical, hormonal, and metabolic features that have been described in patients with AME including sib pairs (DiMartino-Nardi *et al.*, 1987; Shackleton *et al.*, 1986) suggests an inborn error of metabolism attributable to a defect in the gene encoding 11 β -hydroxysteroid dehydrogenase (New *et al.*, 1982; Oberfield *et al.*, 1983). Attempts to evince 11-⁴ISD deficiency in parents have yielded positive results in one father whose excretion of tritiated water was slightly low compared with controls (M. I. New, P. Speiser, and H. L. Bradlow, unpublished) and in one mother with mild hypokalemia and hypertension (Stewart *et al.*, 1988). The fact that a subtle enzyme defect could not consistently be demonstrated in parents of these patients

(Shackleton *et al.*, 1985; DiMartino-Nardi *et al.*, 1987) does not negate the genetic theory.

Apparent mineralocorticoid excess occurs in all racial groups and is equally distributed between males and females (Stewart *et al.*, 1987) (Table VIII). Among patients identified to date, ages at diagnosis have ranged from 6 months to 20 years. The fact that no adults with the condition have been described suggests that the disease, if untreated, is invariably fatal. Five patients have died, yielding a mortality rate of 25%. Most patients had some evidence of end organ damage at the time of diagnosis. Two patients had severe complications of aortic insufficiency, one requiring aortic valve replacement. Although the initial therapeutic response to mineralocorticoid blockade with spironolactone is good, patients eventually require two to three antihypertensive medications to maintain their blood pressure within a safe range. It is not well understood why the hypertension in this syndrome follows a more malignant course than in other forms of mineralocorticoid-induced hypertension.

2. Licorice Ingestion

Studies of licorice ingestion provide further insight into the mechanism of glucocorticoid-mediated hypertension. A decade after the first complete description of a patient with AME, Stewart, Edwards, and colleagues were able to show that when healthy adult males were given 200 g/day of licorice (containing 580 mg glycyrrhizic acid, the active component of the confection) their hormonal and metabolic profiles paralleled the profile of AME patients (Stewart *et al.*, 1987). This led to the crystallization of a proposal first promulgated by New in 1982 (New *et al.*, 1982): An increase in cortisol versus metabolically inactive cortisone causes saturation of cortisol binding globulin, allowing cortisol to gain access to the mineralocorticoid receptor (MR), which allows no intrinsic preference for aldosterone as a ligand. Thus, the 11-⁴ISD is the integral link in protecting renal MR from the normally extant 1000-fold excess physiologic concentration of cortisol compared with aldosterone (Edwards *et al.*, 1988; Ponder *et al.*, 1988).

Recent evidence suggests that the metabolic effects of carbenoxolone and glycyrrhethinic acid, in contrast with their clinical effects, may differ from each other. Stewart and Edwards (1991) have shown that carbenoxolone, in contrast with glycyrrhethinic acid, did not change urinary (allo)THF + THF/THF, or alter plasma cortisone in volunteers. The metabolic profile resembles that of a form of AME reported by Ulick.

TABLE VIII
REPORTED CASES OF APPARENT MINERALCORTICOID EXCESS (AME)

| Patient* | Patient age ^b (years) | Sex | R ^c | Blood pressure (mm Hg) | Aldosterone ^d (ng/dl) | Citation* |
|-----------------|-------------------------------------|-----|----------------|------------------------------|-------------------------------------|-------------------------|
| 1 | 3 | F | 10.2 | 175/115 | 1.9 | (1) |
| 2D (14 years) | 3 | F | >7 | 144/104 | ND | (2) |
| 3D (12 years) | 0 1/12 (2 9/12) | F | 9.8 | 180/120 | ND | (3) |
| 4 | 1 7/12 | M | >4 | 140/100 | ND | (3) |
| 5 | 9 | M | >10 | 250/180 | ND | (4) |
| 6 | 3 3/12 | F | 40 | 125/85 | — | G. Phillipou (1978)/ |
| 7 | 2 (4) | M | 15.9 | 140/90 | 1.3 | (5) |
| 8D | 1 7/12 | M | 45 | 150/110 | 3 | (6) |
| 9D (5 1/2 year) | 0 5/12 | M | 70 | 200/100 | 1.1 | (7) |
| 10 | 0 9/12 (19) | F | 15 | 150/100 | 2.4 | (8) |
| 11R | 3 | F | 31.2 | 170/110 | <0.2 | (9) |
| 12R | 3 9/12 | F | 13.4 | 200/129 | — | (9) |
| 13 | 7 | F | 29.8 | 160/120 | ND | (10) |
| 14 | 9 | M | 26.9 | 170/100 | — | J. S. D. Winter (1988)/ |
| 15 | 3 | M | 7.5 | 200/110 | — | Peskovitz (1986)/ |
| 16 | 21 | M | 13.5 | 200/145 | >3.4 | (11) |
| 17R | 2 (9 4/12) | F | 8.9 | 130/90 | ND | (12) |
| 18R | 2 6/12 (4 4/12) | F | 20 | 142/98 | ND | (12) |
| 19 | 14 9/12 | M | 8 | 130/90 | ND | (12) |
| 20 | 2 3/13 | M | 129 | | | Wood (1992)/ |

* D, patient died (age at death); R, the adjacent patients are siblings.

^b Two ages are presented in some cases. The first is the one in which hypertension was reported. The second, in parenthesis, is the age at which AME was diagnosed. A single figure indicates that hypertension was found at the time of AME diagnosis.

^c R = (THF + 5αTHF)/THF.

^d Normal range is 5 to 20 ng/dl. ND, not detected.

(1) Werder *et al.* (1975); (2) New *et al.* (1977); (3) Winter and McKenzie (1977); (4) Ulick *et al.* (1979); (5) Shackleton *et al.* (1980); (6) Fisliler *et al.* (1982); (7) Honour *et al.* (1983); (8) Harinck *et al.* (1984); (9) Shackleton *et al.* (1985); (10) Batista *et al.* (1986); (11) Stewart *et al.* (1988); (12) Monder *et al.* (1986).
/ Unpublished.

3. Alternative Forms of AME

Ulick has described a Type 2 AME in which the cortisol metabolic clearance rate is delayed, but the conversion of cortisol to cortisone is not impaired (Ulick *et al.*, 1989). Unlike patients with the classic form of AME, hypertension in the Type 2 patients is ameliorated by treatment with dexamethasone (Ulick *et al.*, 1990). It has been suggested that these cases might be explained based on a generalized defect in cortisol metabolism. Alternatively, the absence of a discernible alteration in the T11E:THF ratio may reflect equivalent defects in both the oxidative and the reductive components of the 11-HSD system. Support for the latter theory derives from *in vivo* studies with carbonoxalene in which cortisol half-life was prolonged, yet the T11E:THF ratio was not perturbed (Stewart *et al.*, 1989).

Liddle has described a familial hypertensive syndrome with signs of mineralocorticoid excess responsive to fludrocortisone, but not to spironolactone (Liddle *et al.*, 1963). The proposed etiology for this disorder is enhanced renal tubular sensitivity to low levels of mineralocorticoids; several additional cases have since been reported (Mittre *et al.*, 1967; Helbock and Reynolds, 1970; Wachtel *et al.*, 1975; Costin *et al.*, 1979; Wong *et al.*, 1981).

Although the syndrome of AME usually results in severe and often fatal hypertension and has most often been diagnosed in children (New and Levine, 1977; New *et al.*, 1977; Winter and McKenzie, 1977; Werder *et al.*, 1974; Fisliler *et al.*, 1982; Honour *et al.*, 1983; Harinck *et al.*, 1984; Shackleton *et al.*, 1985), one adult-onset case has been recognized (Stewart *et al.*, 1988). This patient was thought to have the classic form of AME, but unlike most of the others reported, he was responsive to dexamethasone treatment in terms of restoring positive potassium balance and elevation of plasma renin activity, although blood pressure was not significantly changed. Other classic cases where dexamethasone was tried with some salutary effects were reported by Werder *et al.* (1974) and others (Fisliler *et al.*, 1982; Honour *et al.*, 1983; Harinck *et al.*, 1984; Shackleton *et al.*, 1980). Secondary effects of apparent mineralocorticoid excess have also been reported, most notably the coexistence of large renal calculi (DiMartino-Nardi *et al.*, 1987), and in one case actual rickets due to secondary hyperparathyroidism (Batista *et al.*, 1986).

Clinical characterization of the syndrome of apparent mineralocorticoid excess has provided unique and powerful insights into the importance of 11β-hydroxysteroid dehydrogenase in blood pressure homeostasis. For the clinician, there are as yet several unresolved questions: (1) Why are these patients not Cushingoid in light of the low

plasma ACTH and accompanying prolonged cortisol half-life? (2) Conversely, if they are not in a state of cortisol excess as reflected by low plasma ACTH levels, how are they able to survive stressful illness without cortisol supplementation? (3) What are the relationships between the variant syndromes that have been described? (4) Why is there so much heterogeneity among patients with respect to the therapeutic efficacy of low-sodium diet, spironolactone, triamterene, and dexamethasone?

4. *The Defect in AME is Mainly in the Kidney*

That the primary defect of AME was in the kidney tubule was deduced from the fact that hypertension and salt imbalance was controlled using the therapeutic regimen utilized for the treatment of aldosteronemia, despite clear evidence for hypoadosteronemia in these patients. How the imbalance in conversion of cortisol to cortisone is related to juvenile hypertension emerges from the observation that, in all species, cortisol (and corticosterone) at sufficiently high concentrations are mineralocorticoids. Under normal physiological conditions, the active mineralocorticoid is aldosterone; glucocorticoids have no important role in salt metabolism. It is now known that MR bind aldosterone and corticosterone (or cortisol) with equally high affinity (Amelung *et al.*, 1953b). Punder (1987) posed the following significant question: how does aldosterone get its message through to the mineralocorticoid target tissues in the face of much higher circulating free levels of the glucocorticoid? In attempting to answer this question, Stewart and Edwards (1990) and Punder (1990a) have presented a refined and expanded version of earlier proposals (Now *et al.*, 1982) that were designed to explain the role of 11β -dehydrogenase in blood pressure control. They proposed that the role of 11β -dehydrogenase in highly vascular tissues, such as the kidney, is to provide an enzymatic barrier to prevent the accumulation of levels of glucocorticoid that would compete with aldosterone for MR. In AME, this barrier is defective, aldosterone secretion is suppressed, and cortisol, because it cannot be inactivated, is utilized by the receptor as if it were a mineralocorticoid.

B. 11-Oxoreductase Deficiency

Independent reports by Taylor *et al.* (1984) in England and Philippou (Philippou and Jiggins, 1985) in Australia described female patients with apparent deficiency of 11-reduction. These women presented with hirsutism and bilaterally enlarged adrenal glands. Plasma androgen

concentrations were about five times above normal; plasma and urinary free cortisol were normal. Examination of the urinary steroid revealed a 7- to 9-fold increase in cortisol metabolites and a 6- to 10-fold increase in androgens. The ratio of $11\text{E}/(11\text{F} + 5\alpha 11\text{F})$ was extremely high (26, normal ca. 1). These are the only recorded examples of selective 11-oxoreductase deficiency. The evidence indicates two conditions, AME and 11-oxoreductase deficiency, in which 11-HSD appears to be expressed in opposite directions with little reversibility.

VIII. ENZYMOLOGY AND MOLECULAR BIOLOGY

A. The Uniqueness of 11-HSD

The metabolism of a steroid in its target cell determines its effective intracellular concentration, its accessibility to its receptor, and its ability to affect cell function. Thus, any catabolizing enzyme could qualify as a candidate for controlling tissue steroid levels. In this sense, the role of 11-HSD is potentially not different from that of any other enzyme. There are, however, characteristic properties of 11-HSD that, taken together, make it unique. First, 11-HSD affects the activity of glucocorticoids and no other steroid class. Second, 11-HSD is the dominant, if not the sole, enzyme responsible for modifying intracellular corticosteroid levels in many tissues. Third, the enzyme is reversible, enabling it to control the direction of corticosteroid metabolism, thus permitting it to catalyze 11-oxidation to diminish intracellular glucocorticoid concentrations, or 11-reduction to increase them. Fourth, in circumstances where selectivity of aldosterone activity is crucial, such as in the kidney or brain, the enzyme specifically depletes glucocorticoid, without affecting mineralocorticoid.

B. PURIFICATION AND PROPERTIES OF HOMOGENEOUS 11-HSD

1. Purification

The selective directionality of 11-HSD catalysis has led to numerous hypotheses, some assuming a unique reversible enzyme, others a complex of separate, intercommunicating proteins expressing either 11β -dehydrogenase or 11-oxoreductase activities. Attempts to separate these activities or purify 11-HSD have, in the past, been unsuccessful (Hurlock and Islay, 1969; Bush *et al.*, 1968). The enzyme of rat liver is embedded in the endoplasmic reticulum, and because of this, its

purification presents particular problems unique to membrane-bound proteins. Release of the protein from the membrane without denaturing it is usually achieved by displacing the detergent-like native environment with a synthetic detergent (Hjelmeland and Chrambach, 1984; Heienius and Simons, 1975; Tanford and Reynolds, 1976; Rujin, 1972; Lakshmi and Monder, 1985a). Detergent extraction releases 11-HSD in a soluble state, but does not separate oxidation and reduction activity (Lakshmi and Monder, 1985a).

To investigate the properties of 11-HSD, it was purified from rat liver using NADP-agarose affinity chromatography. The homogeneous enzyme preferentially used NADP as cosubstrate; NAD was about 30% as effective (A. Marandici and C. Monder, unpublished observations). The enzyme expressed no detectable 11-oxoreductase activity. This observation initially reinforced the conclusion that 11-HSD is a complex of separate 11 β -dehydrogenase and 11-oxoreductase components (Lakshmi and Monder, 1988).

2. Properties of Purified Enzyme

The homogeneous 11 β -dehydrogenase is a glycoprotein with a monomer molecular weight of about 34,000. It readily aggregates into clusters of 6 to 11 units, due to the mutual attraction of its hydrophobic regions. Total liver 11-HSD activity is the sum of high K_m (6 μ M, corticosterone as substrate) and low K_m (50 nM) activities. Purified enzyme expresses the kinetic behavior of the high K_m form (Monder and Lakshmi, 1989b).

Kinetic analysis and ligand binding studies of purified 11-HSD reveals that the behavior of the enzyme conforms to an ordered sequential mechanism (Monder *et al.*, 1991). In the oxidative direction, the obligatory sequence of addition of cosubstrates requires that NADP be bound first, followed by corticosteroid. Because the enzyme does not express 11-oxoreductase activity, no kinetic analysis has been possible in the reductive direction.

3. Antibodies

Monospecific, polyclonal antibodies to homogeneous rat liver 11-HSD generated in rabbits (Monder and Lakshmi, 1990) have been used to investigate the organ-specific distribution and physiological functions of this enzyme in several organs (Monder, 1991a,b). In all tissues of the rat thus far investigated, 11-HSD antibody reveals a 34K protein, indistinguishable from that of the rat liver enzyme (Monder and Lakshmi, 1990). The intensities of the bands on electrophoretograms after Western blot analysis generally corresponded in magnitude with

enzyme activity. A few tissues that expressed 11-HSD activity had no evidence of 11-HSD-like immunoreactivity, suggesting that they contain possible alternative enzyme forms (Monder, 1991a).

C. MOLECULAR ANALYSIS

1. Structure-Function Predictions

As a first step in the molecular genetic analysis of this enzyme, clones encoding 11-HSD were isolated by probing a rat liver cDNA library in the phage λ gt11 with a monospecific antiserum to 11-HSD (Agrawal *et al.*, 1989). Analysis of clones demonstrated that the mRNA encoding this enzyme in the rat has an open reading frame that predicts a polypeptide of 287 residues with a molecular weight of 31,800, in contrast to the purified protein's actual MW of 34,000. The difference may be due to glycosylation; there were two potential sites for N-glycosylation in the predicted sequence. The rat clone was subsequently used to isolate human 11-HSD cDNA clones from a testis library (Thannin *et al.*, 1991). The amino acid sequence of human 11-HSD predicted from the nucleotide sequence is 79% identical to the corresponding rat sequence.

A search of sequence databases revealed that the predicted sequence of 11-HSD was related to several other prokaryotic and eukaryotic enzymes (Baker, 1989, 1990a). These include steroid 3 α ,20 β -hydroxysteroid dehydrogenase from *Streptomyces hydropneumans* (Marakov *et al.*, 1990), a murine 27-kDa adipocyte protein of unknown function, the *nodG* gene product of the nitrogen fixing bacterium *Rhizobium meliloti*, ribitol dehydrogenase from *Klebsiella aerogenes*, the *act III* gene product from *Streptomyces coelicolor*, human estradiol 17 β -hydroxysteroid dehydrogenase, and dihydrodiol dehydrogenase from *Pseudomonas* species. Although it could not be readily aligned directly with 11-HSD, alcohol dehydrogenase of *Drosophila melanogaster* showed significant similarity to several of the other dehydrogenases used in the alignment (Baker, 1990b). Examination of these alignments (excluding *Drosophila* alcohol dehydrogenase) revealed a total of nine residues that were conserved in all proteins. These residues are likely to be structurally or functionally important. Although it, too, could not be aligned with 11-HSD using the computer algorithm, human 3 β -hydroxysteroid dehydrogenase retains six of these nine residues in a similar arrangement (Fig. 4). Three of these residues are in an area near the amino terminus that is similar to known nucleotide cofactor binding sites of other enzymes, including




Fig. 4. (Top) Conserved amino acid sequences in 11 β -hydroxysteroid dehydrogenase (bold letters) and related enzymes, in descending order, the sequences are 11-HSD, 17 β -hydroxysteroid dehydrogenase, 3 β -hydroxysteroid dehydrogenase, ribitol dehydrogenase (Klebsiella aerogenes), and 11 β -protein from Streptomyces coelicolor. Amino acids shown are A, alanine; C, cysteine; D, aspartic acid; E, glutamic acid; F, phenylalanine; G, glycine; H, histidine; I, isoleucine; K, lysine; L, leucine; M, methionine; N, asparagine; P, proline; Q, glutamine; R, arginine; S, serine; T, threonine; V, valine; W, tryptophan; Y, tyrosine. Absolutely conserved residues are boxed. The positions of three sequences are indicated by the dark boxes within the first three sequences. The 11-HSD amino acid sequence, (function) Proposed active site of rat liver 11-HSD showing the spatial relationship of tyrosine-179, the pyridine ring of NADP⁺, lysine-183, and position 11 of the steroid substrate.

yeast alcohol dehydrogenase (Jornvall *et al.*, 1981). If the three absolutely conserved residues distal to the cofactor binding site (Asp-114, Tyr-183 and Lys-187, human 11-HSD; Asp-110, Tyr-179, Lys-183, rat liver 11-HSD) participate in the catalytic function of the enzyme, they should be near the pyridine ring of NADP⁺ and/or the 11a position of the steroid, a hypothesis that could be tested if the three dimensional structure of 11-HSD could be determined by X-ray crystallography. The three-dimensional structure of a related enzyme should also provide useful information concerning the functional significance of the conserved residues. Crystallographic studies of 3 α ,20 β -hydroxysteroid dehydrogenase of *S. hydropneumoniae* were recently reported (Ghosh *et al.*, 1991). In this related enzyme, the conserved region near the amino terminus does form part of the nucleotide cofactor binding site. The conserved tyrosine residue (Tyr-152 in 3 α ,20 β -HSD) is indeed located near the pyridine ring of the cofactor in a cleft that is presumed to be the steroid binding site. The conserved lysine is directly behind the tyrosine (i.e., on the opposite side of the tyrosine ring from the cofactor). There is demonstrable bridging of electrons between the phenolic hydroxyl of tyrosine and the δ -amino group of lysine, suggesting an interaction between these groups. These findings support the idea that the conserved tyrosine and lysine participate in the catalytic function of the enzyme by facilitating the transfer of a hydride radical from the steroid to the cofactor. In contrast, the conserved aspartate (Asp-82 in 3 α ,20 β -HSD) is not located near the cofactor or the steroid, and its functional significance is difficult to assess from these studies. Thus, it will be necessary to test the importance of these residues in 11-HSD by *in vitro* mutagenesis and expression of the cDNA in cultured cells.

2. Functional Characteristics of Recombinant 11-HSD

To determine whether both 11 β -dehydrogenase and 11-oxoreductase activities resided in the same enzyme, a full-length cDNA clone was expressed in Chinese hamster ovary (CHO) cells by transient transfection with a plasmid expression vector. Enzymatic activities were determined by incubating transfected cells with radioactive substrates. Whereas normal CHO cells did not contain significant 11 β -dehydro-

1, leucine; M, methionine; N, asparagine; P, proline; Q, glutamine; R, arginine; S, serine; T, threonine; V, valine; W, tryptophan; Y, tyrosine. Absolutely conserved residues are boxed. The positions of three sequences are indicated by the dark boxes within the first three sequences. The 11-HSD amino acid sequence, (function) Proposed active site of rat liver 11-HSD showing the spatial relationship of tyrosine-179, the pyridine ring of NADP⁺, lysine-183, and position 11 of the steroid substrate.

genase and 11-oxoreductase activities, these cells developed roughly equal levels of both activities (about 40% conversion of substrate to product after 20 h) after transfection with the expression plasmid. Addition of glycylthionine acid, a known inhibitor of 11 β -dehydrogenase, reduced expressed dehydrogenase activity by 50% without affecting reductase activity (Lakshmi and Monder, 1986b).

To obtain kinetic parameters for the two activities, 11-HSD was expressed at higher levels using recombinant vaccinia virus (Agarwal *et al.*, 1990). Dehydrogenase and reductase activities were assessed in cellular lysates in the presence of saturating concentrations of NADP and NADPH, respectively. At pH 7.0, the recombinant enzyme had very similar K_m and first-order rate constants (V_{max}/K_m) for both activities. These results were consistent with the hypothesis that both dehydrogenase and reductase activities reside in a single enzyme. Exposure to NADIP resulted in rapid and irreversible inactivation of the reductase activity of the enzyme, a phenomenon consistent with the instability of the reductase during attempted purification from rat liver.

In contrast, when the recombinant enzyme was prepared from cells grown in the presence of A, tunicamycin (an inhibitor of glycosylation), dehydrogenase activity was reduced by about 50%, whereas reductase activity was unaffected. This was associated with increased amounts of a 31-kDa enzyme species that presumably represented the unglycosylated enzyme. This suggests that the dehydrogenase activity of the enzyme may depend on adequate glycosylation.

3. Tissue Distribution of 11-HSD Expression

In initial studies, the rat cDNA clone hybridized to a single mRNA species of approximately 1600–1700 nucleotides in samples from testis (highest), liver, kidney, and lung but did not hybridize to samples from heart or colon. This distribution roughly paralleled that of 11 β -dehydrogenase activity.

A subsequent study (Krozowski *et al.*, 1990) suggested that the rat kidney actually contains several cross-hybridizing mRNA species of 1900, 1600, and 1500 nucleotides (renal cortex/medulla) and 1700 nucleotides (renal papilla). In this study, the highest level of expression was found in the liver, followed respectively by kidney, lung, testis, hippocampus, heart, and colon.

In further studies of expression in rat brain (Moisin *et al.*, 1990a,b), an apparently identical mRNA species was found in all areas, but at highest levels in the hippocampus and cortex. It is speculated that 11-HSD regulates the access of glucocorticoids to cerebral mineralocor-

ticoid and/or glucocorticoid receptors, thus modulating steroid hormone effects of cerebral function.

The tissue distribution of the human mRNA differs from that in the rat; it is expressed at very high levels in the liver and at much lower levels in the kidney. The significance of these findings, given the importance of this enzyme activity in the kidney, is not yet clear, but it is consistent with the idea that there may be additional proteins with 11-HSD activity in the kidney.

4. Genetic Analysis of Human 11-HSD

To determine the chromosomal location of the human 11 β -hydroxysteroid dehydrogenase (*HSD11*) gene, a cDNA clone was hybridized to DNA samples from a panel of human-rodent somatic cell hybrid lines. Hybridization to human-specific bands was consistent with a location on chromosome 1 (Tannin *et al.*, 1991).

Hybridization of whole of uncloned human genomic DNA that had been digested with restriction endonuclease *Hind*III demonstrated that there was a single *HSD11* gene that was carried on two fragments. Sequence analysis of these fragments showed that they carried a single gene consisting of six exons, the first four of which were contained on the smaller fragment. Comparison of the maps of restriction sites in these fragments with results of hybridization to uncloned DNA revealed that there must be an additional *Hind*III fragment(s) of undetermined size in intron 4 that contains *Eco*RI and *Bam*HI sites.

a. Transcriptional Regulation of the HSD11 Gene. Primer extension analysis indicated that transcription of the human *HSD11* gene starts 93 bp upstream from the start of translation (Tannin *et al.*, 1991). This yields a 5' untranslated region very similar in length to that of rat 11-HSD mRNA. There is no TATA box in the 5' flanking region, but there is a consensus CAAT box (CCAATC) 76 bases upstream from the start of transcription. An 8-bp palindromic sequence (CTGTACAG) was present 188 bp upstream from the start of transcription. It resembles part of a glucocorticoid response element (Evans, 1988), which would be consistent with the known ability of glucocorticoids to increase levels of 11-HSD activity. However, its functional significance requires further study, particularly in light of recent work suggesting that glucocorticoids do not alter the level of *HSD11* gene expression in rat liver, lung, or kidney (Krozowski *et al.*, 1990).

Recent SI nuclease analysis suggests that the different-sized mRNA transcripts observed in rat kidney apparently have different 5' extensions. Cloning studies suggested that some transcripts have a divergent 5' coding sequence that encodes a putative protein with a

truncated amino terminus (Krozowski *et al.*, 1992). Comparison of the sequence of the truncated clones with that of the human gene suggests that these clones originate by transcription within the first intron of the corresponding rat gene. It is not yet known whether the putative protein is functional or even whether it is synthesized *in vivo*.

b. Possibility of Additional 11-HSD Enzymes. In addition to the putative truncated form of the protein, other evidence suggests that there may be an additional enzyme(s) with 11-HSD activity. As mentioned, there appears to be some discrepancy between the levels of 11-HSD activity and *HSD11* mRNA in human kidney. In rat kidney, anti-11-HSD sera react with proximal tubules but not with distal tubules/collecting ducts, although the latter represent the main site of mineralocorticoid action (Rundle *et al.*, 1989a). The 11-HSD activity of isolated rabbit distal tubules and collecting ducts differs markedly in kinetic properties from the enzyme in the proximal tubule (which is similar or identical to the enzyme in the liver) in having a K_m about 100-fold lower (Naray-Fejes-Toth *et al.*, 1990). A similar low K_m form has been detected as a minor species in the liver (Monder and Lakshmi, 1989b). Furthermore, histochemical studies have suggested that the distal tubule contains an 11-HSD activity that requires NAD rather than NADP (Mercer and Krozowski, 1992). No mRNA was detected in heart, yet heart readily converts cortisol to cortisone (C. Monder and A. Marandici, unpublished observations) (Kolanowski *et al.*, 1991). Because Southern blotting studies indicate that rats and humans carry only one *HSD11* gene, the gene(s) encoding any additional 11-HSD activities must be sufficiently different from *HSD11* in their nucleotide sequences that they do not cross-hybridize.

A number of questions regarding the functions of 11-HSD may be answered by molecular genetic analysis of patients with inherited enzymatic deficiencies. Because both dehydrogenase and reductase activities apparently reside in the same enzyme, it will be of obvious interest to search for mutations in the *HSD11* gene(s) associated with AME and 11-oxoreductase deficiencies and correlate their effects on enzymatic function with clinical phenotype.

IX. 11-HSD FUNCTION IN SPECIFIC ORGANS

A. Kidney

1. Mineralocorticoid Receptors and 11-HSD

We have discussed the fact that the characteristic biochemical abnormality of AME is a severe loss in the ability of patients with this

disability to oxidize cortisol. A working model connecting 11-HSD activity and blood pressure control evolved from the convergent findings of many laboratories. It was discovered that, *in vitro*, the renal MR bound aldosterone and the glucocorticoids, corticosterone and cortisol, with comparable affinity (Krozowski and Funder, 1983; Arriza *et al.*, 1987; Armanini *et al.*, 1985). That this behavior is an intrinsic property of MR was shown by Arriza *et al.* (1987) using cloned recombinant MR derived from placental cDNA expressed in COS cells. However, *in vivo*, aldosterone was selectively taken up by the MR of some tissues, such as kidney, parotid, and colon (Sheppard and Funder, 1987a, b), to the exclusion of glucocorticoids. That these tissues are aldosterone selective appeared to be in conflict with the *in vitro* data. To reconcile the *in vitro* and *in vivo* evidence, a hypothesis to explain steroid selectivity was developed based on the observation that glucocorticoids are uniquely sequestered to corticosteroid binding globulin (CBG) and are thus made unavailable to MR (Krozowski and Funder, 1983; Sheppard and Funder, 1987a). It was, however, found that aldosterone selectively persisted *in vivo* in young rats with little or no circulating CBG (Sheppard and Funder, 1987a, b). Since MR is coded for by a single gene (Arriza *et al.*, 1987), it was considered unlikely that selectivity depends on tissue- or age-specific variations in its intrinsic properties. However, hormone-dependent gene regulation by MR showed a preference of aldosterone over cortisol (Arriza, 1991). It was recently suggested that post-translational modifications of MR may play a role in mineralocorticoid binding and specificity, but this possibility remains to be evaluated (Doyle *et al.*, 1988). A search for specific regions of MR localization in the rat nephron revealed that MR are not distributed uniformly throughout the renal tubule and are localized to the distal tubule. Krozowski *et al.* (1989), using an antiserum corresponding to the hinge region of human MR (Arriza *et al.*, 1987), showed that MR are localized in the principal cells of the cortical collecting duct. MR were not detected in the proximal tubule and glomerulus (Wrange and Yu, 1983). They have been reported to be present in the distal convoluted tubule and the thick ascending limb of the loop of Henle (Dovic and Kutz, 1981; Kutz, 1980; Patman *et al.*, 1983). Thus, although there is some uncertainty about the range of MR distribution in the distal tubule, its localization to this region is not questioned (Krozowski *et al.*, 1989; Rundle *et al.*, 1989b; Bonvalet, 1991).

The renal distribution of immunoreactive 11-HSD and MR was very different. Polyclonal antibodies (Edwards *et al.*, 1988; Rundle *et al.*, 1989b) and monoclonal antibodies (Casalillo *et al.*, 1988) to 11-HSD revealed specific immunoreactive staining in the proximal tubules of the inner cortex. The focal distribution of 11-HSD and MR within the

tubule suggested a model in which blood filtrate containing aldosterone and a thousandfold greater concentration of corticosterone must pass through a region of high 11-DII activity, located in the proximal tubule and vasa recta. During this passage corticosterone is completely inactivated, leaving the aldosterone, which is not a substrate of the enzyme, unaltered, permitting its binding to MR to occur unopposed. Since 11-DII and MR do not colocalize, a paracrine relationship between them is inferred. An explanation of the consequences of the deficiency of 11-DII in humans is predicated on the assumption that these structural and functional relationships apply to human as well as to rat kidney.

X2. Role of Glucocorticoid Receptors

The hypothesis based on this model assumes that all available glucocorticoid is oxidatively inactivated as it passes through the region of high 11-HSD activity. The model in this form proved to be extremely useful and enabled puzzling aspects of AME to be explained. It is, however, known that the renal tubule contains glucocorticoid receptors. These mediate glucocorticoid-specific effects on the kidney including effects on renal hemodynamics, acid and water excretion, gluconeogenesis, and sodium-potassium ATPase (Katz, 1990; Kinsella, 1990). Consequently, complete inactivation of corticosterone or cortisol would not be a desirable option for the kidney. Therefore, the hypothesis originally proposed that envisioned a relationship between the MR and 11-HSD that excludes GR-mediated effects must be revised. Glucocorticoid receptors are known to be distributed along the nephron (Parman *et al.*, 1991; Katz, 1990). The number of glucocorticoid binding sites in the thick ascending limb of the loop of Henle, for example, is 100-fold higher than the aldosterone binding sites (Lee *et al.*, 1983). Maximal binding capacity of the cortical collecting tubule for corticosterone is greater than for aldosterone (Katz, 1990). Thus, though glucocorticoids at moderate concentrations may mediate renal function by way of MR, GR are important, as well (Naray-Fejes-Toth and Fejes-Toth, 1990; Clore *et al.*, 1988). The emerging concept that 11-HSD plays an important role in mediating GR-dependent processes is supported by the observation that there is a strong correlation between GR and 11-HSD distribution in tissues (Whorwood *et al.*, 1992).

3. The Protector Role of 11-HSD: Modifying the Hypothesis

The hypothesis that renal 11-HSD enables MR to interact selectively with aldosterone as both ligand and effector by inactivating potentially competing glucocorticoids was supported by extensive laboratory

and clinical data (Funder, 1990a,b; Stewart and Edwards, 1990; Edwards *et al.*, 1989; Monder and Shackleton, 1984; Monder, 1991b). However, the proximal 11-HSD-distal MR model initially proposed to explain the protector function of 11-HSD proved to be less and less adequate as it was reexamined. The great physical distance between 11-HSD and MR appeared to result in an inefficient functional unit. Furthermore, that all available corticosteroid must be inactivated by 11-HSD, implicit in the model, was an untested assumption. The functions of renal GR and the possibility of a functional link between active (steroid bound) GR and the expression of MR remained to be investigated.

The requirement of the original model that access of aldosterone to MR could be achieved only if the tubular filtrate were completely cleansed of glucocorticoid was an extremely stringent one. In order to accommodate this requirement, it was proposed that 11-HSD is strategically distributed along the nephron in order to oxidize residual glucocorticoid. Bonvalet *et al.* (1990) found 11-HSD activity in the distal as well as the proximal portions of the rabbit kidney tubule. Possibly, it was suggested, 11-HSD and MR may coexist in distal cells (Naray-Fejes-Toth *et al.*, 1991; Stewart *et al.*, 1991; Bonvalet, 1991). Gradient fractionation of rat kidney tubules indicated that 11-HSD was indeed in the distal as well as proximal regions (Edwards *et al.*, 1988). The immunohistochemical studies that led to the conclusion that 11-HSD was localized solely in the proximal region of the nephron were in obvious conflict with the enzyme activity data (Rundle *et al.*, 1989a; Castello *et al.*, 1989). The lack of immunoreaction of the distal tubular enzyme with rat liver 11-HSD antibody could be explained by assuming that the enzyme was not easily accessible to antibody because of a transcellular barrier, or was a distantly related antigen. Naray-Fejes-Toth *et al.* (Naray-Fejes-Toth and Fejes-Toth, 1990; Bonvalet *et al.*, 1990), using rabbit kidney cortical collecting tubules isolated by solid-phase immunoadsorption, conclusively showed that there was 11-HSD activity in this region that did not react with antibody on Western blot analysis.

Naray-Fejes-Toth *et al.* (1991) found that the level of 11-HSD activity in monolayer preparations of CCD cells was sufficient to completely convert corticosterone to 11-dehydrocorticosterone in a single pass through the membrane. Therefore, in an individual MR-binding cell, there may be enough 11-HSD to inactivate glucocorticoid completely, thus satisfying the requirement that aldosterone bind MR unencumbered by competing steroids. Differences were also noted in the distribution of 11-HSD in the renal cortex and medulla

(Castello *et al.*, 1989). Consistent with the above postulated autocrine role, the 11-HSD of the distal tubule was more active than the proximal tubule (Edwards *et al.*, 1988; Castello *et al.*, 1989; Bonvalet *et al.*, 1990). A schematic view of the current understanding of corticosteroid associated interactions in normal kidney is shown in Fig. 5.

Evidence that salt metabolism may be mediated through GR as well as MR have been presented by Naray-Fejes-Toth and Fejes-Toth (1990) and Funder *et al.* (1990). The following observations support this conclusion: (a) AME patients are more sensitive to cortisol than aldosterone in terms of increased blood pressure and sodium retention; (b) in pseudohypoaldosteronism, a condition characterized by low or no MR, the electrolyte effect of cortisol results in part from occupancy of GR; (c) RU 28362, a GR-specific glucocorticoid that does not bind MR, affects electrolyte excretion via GR; (d) RU 28318, a specific MR antagonist, does not diminish the electrolyte effect of RU 28362; (e)




FIG. 5. A current view of corticosteroid associated interactions in normal and 11-HSD-deficient kidney. Heavy arrows indicate dominant pathway. (1a) Normal kidney. Cortisol (F) in the proximal (and possibly distal) tubule mediates glucocorticoid dependent events via glucocorticoid receptor (GR). (●) The level of F available to GR is mediated by 11-HSD (1). Steroid dissociated from GR is oxidized to cortisone (E) (2) to prevent its reentry into the system. F entering from the proximal tubule and other sources may compete with aldosterone (M), for mineralocorticoid receptor (MR). (■) This competition is prevented by oxidation of F to E (2) in the distal tubule and collecting duct (RTN) (11-HSD) deficient kidney. In the absence of functioning 11-HSD, cortisol cannot be oxidized and accumulates, preferentially binding to MR, displacing M, and initiating a sequence of aldosterone-minimetic events.

immunodissociated rabbit cortical collecting tubule cells responded similarly to aldosterone, dexamethasone and RU 28362; (f) The glucocorticoid receptor antagonist RU 486 blocked the effect of RU 28362, but the MR antagonist 2K 91567 did not; (g) kaliuresis caused by cortisol is blocked by RU 486 (Cloré *et al.*, 1988). Localization of 11-HSD mRNA by *in situ* hybridization using a cRNA probe (Agarwal *et al.*, 1989) indicated its location in the proximal tubules and in the cortical and medullary collecting tubules, a finding that accords with the enzyme distribution studies. The presence of multiple 11-HSD mRNA species in kidney is consistent with the possibility of a heterogeneous population of 11-HSD proteins that may be generated from them, some of which may be recognized by 11-HSD antibody (Krozowski *et al.*, 1990). These results also indicate that the variant forms of 11-HSD that have been proposed may be generally similar in structure.

4. Licorice, Hypertension, and Kidney Function

a. *The Active Agent of Licorice.* Valuable evidence supporting the role of 11-HSD in kidney function emerged from studies on the pharmacological behavior of licorice, a flavoring agent extracted from the roots of *Glycyrrhiza glabra*. Licorice has been used as a medicine and condiment for at least 5000 years (Davis and Morris, 1991). Glycyrrhetic acid (GA), its active ingredient, is a cyclic triterpene whose fused ring structure, illustrated in Fig. 6, closely resembles that of the glucocorticoids. A synthetic agent developed for the treatment for gastric and duodenal ulcers, carbenoxolone (CA), is the 3-O- β -carboxypionyl ester of glycyrrhetic acid. Ingestion of either GA or CA causes




FIG. 6. Glycyrrhetic acid (GA).

clinical effects that resemble those of aldosterone excess, including hypertension, hypokalemia, edema, polyuria, polydipsia, heart failure, and muscle weakness (Pinder *et al.*, 1976; Barron, 1983; Werning *et al.*, 1971).

b. *Possible Explanation of Licorice Actions.* Reevers (1948) first documented the aldosterone mimetic behavior of GA. Explanations for its properties have included suggestions that it (a) stimulates aldosterone production; (b) displaces aldosterone from nonspecific binding sites, thus increasing its effective concentration (Humphrey *et al.*, 1979; Porter, 1970); (c) demonstrates intrinsic mineralocorticoid activity; or (d) potentiates the effects of aldosterone (Humphrey *et al.*, 1979; Armanini *et al.*, 1989b). All except the last two are unlikely mechanisms (Hausemann and Tarnoky, 1968; Porter, 1970). Evidence for binding of glycyrrhetic acid to kidney mineralocorticoid receptors was presented by Ujmann *et al.* (1976), Armanini *et al.* (1983), Takeda *et al.* (1987), and Hayashi *et al.* (1984). There is general agreement that binding of GA to MR is about 1/10,000 that of aldosterone based on competitive binding studies. It is unclear whether significant binding to human or rat kidney MR occurs even under the conditions of massive GA or CA intake. In a load bladder model, 2.5×10^{-5} M CA had no effect on short-circuit current over 360 min of exposure (Gaeggele *et al.*, 1989). Humphrey *et al.* (1979) found that CA did not displace 3 H aldosterone from rat kidney nuclei. The reason may be, as Armanini *et al.* (1989a) suggest, that binding of CA to MR requires its prior hydrolysis to (1A).

It has been estimated that subjects consuming 100 to 200 g of licorice per day have total circulating plasma GA levels of 80 to 480 ng/ml (Hughes and Cowles, 1977; Stewart *et al.*, 1987). The concentration of free circulating GA is lower, since 95% of GA is bound to plasma proteins (Ishida *et al.*, 1988). Thus, the concentration of GA potentially accessible to MR is too low to measurably bind to the receptor under physiological conditions. It is, however, theoretically possible that specific ligand-receptor interaction may lead to some responses resembling that of the binding of mineralocorticoid. The availability of radioactive glycyrrhetic acid (Kanaoka *et al.*, 1988) should make it possible to determine whether its interaction with MR leads to nuclear translocation.

Additional evidence that cannot currently be reconciled with the postulated mineralocorticoid-mimetic behavior of GA is the observation that the effectiveness of GA is abolished in adrenalectomized rodents (Gard *et al.*, 1953; Circed *et al.*, 1960) and humans (Hort *et al.*, 1983; Elmudjani *et al.*, 1986) and is restored when glucocorticoids are administered (Hort *et al.*, 1983). The results indicate that a secretory

product of the adrenal cortex is an essential participant of GA action. Normal individuals ingesting glycyrrhetic acid under controlled conditions for brief periods of time (3–10 days) showed significant decrease in cortisol oxidation to 11-oxo metabolites (MacKenzie *et al.*, 1990), a finding consistent with an inhibitory effect on 11-HSD (Matingly *et al.*, 1970; Chen *et al.*, 1990; Ojima *et al.*, 1990).

c. *Glycyrrhetic Acid and Other Inhibitors of Renal 11-HSD.* Other agents affect the activity of renal 11-HSD. The inhibition of 11-HSD by gossypol, a potential male contraceptive agent extracted from cottonseed oil, resembles that of GA and CA. This observation has led to the suggestion that the hypokalemia observed in men taking this agent has the same cause as that of men ingesting licorice (Sang *et al.*, 1991). Hersholtzer and co-workers (1990b) have found that bile acids, though of low inhibitory potency, are present in the human circulation at concentrations that indicate that they have the potential to modulate 11-HSD activity.

Toulou *et al.* (1984) have made the surprising observation that triolene, a cytochrome derivative known to inhibit 3 β -hydroxysteroid dehydrogenase, increased 1 β -hydroxy oxidation in sheep liver homogenates, a phenomenon that may be species specific. Perschel *et al.* (1991) found that pooled rabbit bile at low concentrations increased rat renal 11-HSD. Whether these examples represent stimulation of 11-HSD, as the authors suggest, or an expression of the ability of P450₁₁ to catalyze the oxidation of cortisol to cortisone (Suhara *et al.*, 1986) remains undecided.

In a recent study, GA and CA were found to be extremely potent inhibitors of 11-HSD in isolated rat kidney microsomes, with K_i values of 3 nM (Monder *et al.*, 1989). In the range 1 to 20 nM, reductase was inhibited poorly (Monder *et al.*, 1989; Hersholtzer *et al.*, 1991). Glycyrrhetic acid is the most powerful known inhibitor of 11-HSD (Monder *et al.*, 1989), but it is 10-fold less potent in intact cells. The basis for this difference is unknown. A transmembrane barrier to GA or sequestration to proteins and other macromolecules has been suggested (Monder, 1991c).

The load bladder, the amphibian counterpart of the nephron, has proven to be a useful model for studying the pharmacological action of CA and GA on the kidney tubule. Using this system, Gaeggele *et al.* (1989) and Brem *et al.* (1989, 1991) have shown that CA allows corticosterone to be as potent as aldosterone in eliciting the mineralocorticoid response, in accord with the proposed role of CA as an inhibitor of 11-HSD.

d. *Glycyrrhetic Acid: An Inhibitor of Broad Specificity.* It is not

worthy that Morris *et al.* have shown that the metabolism of aldosterone is slowed by glycyrrhetic acid, a potent inhibitor of cytosolic 5 β -reductase and microsomal 3 β -hydroxysteroid dehydrogenase (Latif *et al.*, 1990; Tamura *et al.*, 1979; Yoshida *et al.*, 1992). By slowing the rate of inactivation of aldosterone and 11-deoxycorticosterone, these agents potentiate the activity of mineralocorticoids. The two-pronged effect of GA and CA on mineralocorticoid and 11 β -hydroxysteroid metabolism would therefore provide a mechanism for inactivating glucocorticoids and simultaneously enhancing the activity of mineralocorticoids.

There are other ways in which GA or CA can affect renal function (Monder, 1991c). Indirect evidence suggests that GA may inhibit glucuronide formation, since it increases the proportion of unconjugated cortisol in urine of people given massive doses of licorice (equivalent to 0.7 to 1.4 g of GA per day for 1-4 weeks) (Epstein *et al.*, 1978). A possible direct effect of glycyrrhetic acid on (Na⁺ - K⁺) ATPase (Itoh *et al.*, 1989; Baron and Greene, 1986) may account for some of the effects of GA on the kidneys of adrenalectomized animals. The combined effects of GA on glucocorticoid oxidation at C-11, A-ring reduction, and excretion of unconjugated steroids bear a striking resemblance to the metabolic changes characteristic of AME (Monder *et al.*, 1986). The possibility that an endogenous glycyrrhetic acid-like compound contributes to the pathology of AME cannot be excluded.

B. The Vascular Bed

It has been known for about 50 years that adrenocortical hormones influence the behavior of the peripheral blood vessels (Swingle and Remington, 1944). These influences include alterations in intra- and extracellular levels of Na⁺ and K⁺ critical for maintaining vascular tone (Zweifach *et al.*, 1953), and maintenance of the sensitivity of the peripheral vasculature to pressor agents (Darlington *et al.*, 1959; Grunfeld and Eloy, 1987; Ashton and Cook, 1992). These and other effects (Mouré and Worcel, 1984; Nichols *et al.*, 1983, 1984; Jazayeri and Meyer, 1988; Haigh and Jones, 1990; Yamanari *et al.*, 1989) are mediated by MR and GR in vascular smooth muscle (VSM) cells. The presence of MR and GR in vascular smooth muscle provides evidence of direct action of corticosteroids on the arterial wall affecting muscle tone and responsiveness to humoral and neurogenic vasoconstrictive stimuli (Kornel *et al.*, 1975; Onoyama *et al.*, 1979).

The whole arterial tree appears to be a target organ for both mineralocorticoids and glucocorticoids (Kornel *et al.*, 1982). There is evidence that the effects of both steroid classes on vascular tissue proceed by

independent processes (Jazayeri and Meyer, 1988, 1989). Vascular smooth muscle cells in culture are affected differently by mineralocorticoid and glucocorticoids. The glucocorticoid effects are blocked by RU 486, indicating GR dependence (Kornel, 1988; Nichols *et al.*, 1985; Meyer and Nichols, 1981). High levels of glucocorticoids could, by binding both the MR and GR, contribute to the pathogenesis of essential hypertension by stimulating vasoconstriction. Funder *et al.* (1989) found that MR of the mesenteric vascular arcade is aldosterone specific *in vivo*. They suggested that, as in the kidney, 11-HSD may mediate the selective mineralocorticoid response. Funder *et al.* (1989) and Walker *et al.* (1991) have confirmed the original report of Kornel *et al.* (1982) that the vessels of the circulatory system express 11-HSD activity.

The enzyme appears to be predominantly in the smaller vessels, a finding that has been interpreted to indicate that by catalyzing the reversible inactivation of glucocorticoids, it modulates tone in the peripheral resistance beds and thereby influences blood pressure. Alternatively, as occurs in the brain (see later), the availability of NADP may affect 11-HSD activity. There appears to be insufficient NADP in VSM cells to fully activate the available 11-HSD, thus making the nucleotide a limiting factor in the expression of enzyme activity. Consistent with the proposed role of 11-HSD, the enzyme and VSM glucocorticoid receptors are colocalized, indicating that the regulation of GR-steroid interaction occurs by an autocrine mechanism.

C. The Skin

The modulation of corticosteroid effects by 11-HSD appears to extend to the superficial peripheral vessels. The potency of topical corticosteroid on suppression of the inflammatory response is determined in part by its local persistence; oxidative inactivation by dermal 11-HSD diminishes its effectiveness. It has been proposed that the vasoconstrictor action of corticosteroid contributes to the potentiation of its action, by preventing its loss. Thus, dermal 11-HSD, which accelerates the destruction of inactivation of the steroid in skin, would diminish its topical effectiveness. Consequently, inhibition of 11-HSD activity in target tissues should potentiate the local action of glucocorticoids.

This concept has been put to the test by Tealucksingh *et al.* (1990) who investigated the activity of hydrocortisone on skin. Topical application of glycyrrhetic acid inhibited dermal 11-HSD, reducing inactivation of cortisol by skin, prolonging and enhancing its topical anti-inflammatory activity. It has been proposed that this property of GA and CA explains their beneficial effects in inflammatory cut-

aneous disorders (Colin Jones, 1957). However, recent studies have shown that 11-oxoreductase exceeds 11 β -dehydrogenase activity in human skin fibroblasts. Whether this is due to the intrinsic character of the skin enzyme or to another rate-limiting step, such as lack of pyridine nucleotide (as is found in brain), is not known. These observations suggest that in human skin the preferred direction of corticosteroid metabolism is reductive (Hammani and Sileri, 1991; Monder *et al.*, 1986) and, therefore, that the anti-inflammatory effects of topical application of GA on human skin cannot be fully explained by the inhibition of 11- β SD.

D. The Nervous System

1. Neural 11- β SD

Interest in the metabolism of corticosteroids in brain and pituitary evolved simultaneously with the recognition of the importance of steroids on brain function (Woodbury, 1958), on the one hand, and the importance of neuroendocrine influences on steroid secretion, on the other. Soon after cortisol had been isolated from human nerve tissue (Touchstone *et al.*, 1963), evidence for the oxidation of corticosteroids to 11-dehydrocorticosteroids by brain tissue was obtained for rat (Petersen *et al.*, 1965; Sholtz *et al.*, 1965), mouse (Grosser, 1966; Ye and Burton, 1980), dog (Miyabu *et al.*, 1973; Elk-Nas and Brizoe, 1965), and primate (Grosser and Axelrod, 1968). Despite the fact that the presence of 11- β SD in nervous tissue had been known for many years, its possible function in the central nervous system has only recently come under investigation. The working assumption is that brain 11- β SD plays an important role in the expression of glucocorticoid-dependent processes.

2. Receptor-Mediated Selectivity of Corticosteroid Effects

As with kidney, central MR and GR mediate corticosteroid-specific effects. Neutral MR, with properties identical to those of the renal mineralocorticoid binder (Tashima *et al.*, 1989), interacts with corticosterone (or cortisol) and aldosterone with comparable affinity, and binds dexamethasone, a synthetic glucocorticoid, much less efficiently (Beaumont and Fancelli, 1983; Krasowski and Funder, 1983; Wrangé and Yi, 1983). The classic glucocorticoid receptor also uses corticosterone as ligand, but prefers dexamethasone. The equivalent affinity of MR for corticosterone and aldosterone in the rat brain contrasts sharply with the clear preference of the receptor for aldosterone

in the kidney. It therefore follows that the overwhelmingly greater concentration of corticosterone (10^2 - to 10^3 -fold that of aldosterone) in the circulation of the rat would result in MR saturated with and largely dependent for its activity on the circulating corticosterone. The system would thus be insensitive to aldosterone, leading to the conclusion that aldosterone can have no effect on brain function.

There is strong evidence for selective aldosterone effects in the central nervous system. It is known that rat brain takes up both aldosterone and corticosterone, with a similar regional distribution. Highest uptake occurs in the hippocampus, septum, and amygdala (Birmingham *et al.*, 1979; Gertsch and McEwen, 1972; Mogulewsky and Raynaud, 1980). The analysis of brain receptor distribution using RU 26988 (a pure glucocorticoid) and RU 28318 and RU 26752 (pure mineralocorticoids) (Coirini *et al.*, 1985) has led to the conclusion that there are only the two receptor subtypes, MR and GR. These are distributed in neurons and glial cells (Bohn *et al.*, 1991). Despite the 100- to 1000-fold excess of corticosterone over aldosterone in the circulation (Eliere and Peterson, 1964), MR receptors recognize aldosterone in the presence of corticosterone in signaling changes in salt balance, and this effect persists at physiological levels of both mineralocorticoid and glucocorticoid (McEwen *et al.*, 1986; Forman and Mulrow, 1973; Fregly and Rowland, 1985).

Intracerebroventricular (ICV) administration of aldosterone to hominephrectomized rats caused elevated blood pressure. The effect was blocked by the mineralocorticoid receptor antagonist RU 28318. Corticosterone could not replace aldosterone, nor could systemic administration of steroids reproduce these effects (Gomez-Sanchez, 1991). The hypertension induced by aldosterone administered ICV requires corticosterone, for the effect is prevented in bilaterally adrenalectomized rats, and restored by exogenous corticosterone (Gomez-Sanchez, 1991). Arriza and Evans found in a cotransfection assay that MR was more sensitive to mineralocorticoid than to glucocorticoids, despite equivalent binding affinities (Arriza *et al.*, 1988). It has been reported that corticosteroids differentially modulate nutrient intake in rats through central receptor-mediated processes. Implant of aldosterone in the paraventricular nucleus of adrenalectomized rats stimulated ingestion of fat; corticosterone stimulated carbohydrate intake (Temple and Liebowitz, 1989).

The interrelationships between GR, MR, and corticosteroids in the central nervous system are complex. Receptor specificity varies in ways that are not immediately obvious. In early studies, the differential binding of corticosterone to receptors suggested that there may be

three receptor types in the nervous system: the classical GR and MR, and a corticosterone binding subset of MR termed CR. Binding studies with corticosterone and aldosterone *in vitro* showed no distinction between CR and MR, and the former term was abandoned, since its retention obscured the question of the specifically-conferring mechanism. To illustrate this point, MR in the circumventricular region is aldosterone selective; MR in the neurons of the limbic region is corticosterone selective. This selectivity is reflected throughout the nervous system and shows up as differential retention of corticosterone and aldosterone in different subregions.

a. Antagonistic and Synergistic Mechanisms. DeKloet and co-workers have developed a functional rationale for the preferential binding of glucocorticoids to both GR and MR under normal physiological conditions (de Kloet and Reul, 1987). They have shown that the circulating concentration of corticosterone results in 80 to 90% occupancy of cerebral MR. This generates a baseline level of continuously activated receptor that serves to monitor and interpret the animal's external environment. At basal levels of circulating corticosterone, specifically at the diurnal trough, the levels of occupancy of GR is low, under stress, or to a lesser extent at the diurnal peak, the level of circulating corticosterone increases, leading to GR occupancy, generating a negative feedback on stress-activated brain mechanisms. There are, thus, reciprocal balancing, tonic-activating actions and feedback-damping mechanisms. This continuum reflects corticosteroid concentration, and the sequential, selective occupancy of corticosterone to MR and GR. Evans and Arriza *et al.*, 1988) have suggested that MR and GR act as a binary response system for corticosterone. Their model depends on the coordinated synergistic interaction of MR and GR with overlapping sets of genes, the magnitude of the response being dependent on the circulating glucocorticoid levels. This model may be compared with the coordinated antagonistic MR- and GR-mediated effects proposed by DeKloet. Van den Berg *et al.* (1990) suggest that central MR and GR mediate opposing effects of glucocorticoids and mineralocorticoids on blood pressure, consistent with other evidence that MR and GR in the brain mediate reciprocal neurochemical, neuronal, and behavioral responses.

b. Hippocampal Degeneration. The hippocampus illustrates the importance of regional optimization of corticosteroid concentration. This organ contains the highest concentration of MR in the central nervous system (Reul and de Kloet, 1986; Krozowski and Pinder, 1989; de Kloet *et al.*, 1984). It is extremely vulnerable to corticosteroid hormones. Chronic glucocorticoid exposure coupled with other chronic

insults (e.g., aging, ischemia) results in the degeneration of specific hippocampal (CA1, CA3) fields (Supolsky and Pulsinelli, 1985; Dokae, 1990). Glucocorticoid absence also results in specific hippocampal degeneration (Sloviter *et al.*, 1989). There must therefore be a mechanism to maintain glucocorticoids (e.g., corticosterone) in specific hippocampal fields within a defined range of intracellular concentration. Possibly, the survival of these cells requires a persistent occupation of the MR or GR by steroid ligand at some tonic level (Sloviter *et al.*, 1989). That the effects are direct is supported by the observation that cells in culture as well as in intact brain are sensitized by excess glucocorticoid (Masters *et al.*, 1989; Fehsenh *et al.*, 1980).

c. Selectivity of Brain Corticosteroid Receptors: Proposed Role of 11-HSD. The source of the regional selectivity of brain receptors for mineral- or glucocorticoid has not yet been determined. The evolution of hypotheses designed to investigate this question has paralleled those proposed for the kidney. For several years, the only serious contender for a selection mechanism was CBG, but this was withdrawn for the same reason as in the kidney (de Kloet and Reul, 1987; Pinder, 1986); i.e., selectivity was not altered in animals with little or no circulating CBG. The suggestion that receptor selectivity was mediated by the local action of 11-HSD on glucocorticoids, similar to a parallel process in kidney initiated a series of promising investigations. The starting point was the hypothesis that the excess corticosterone in blood entering cells represents an impediment to the ability of aldosterone to gain access to the MR in the absence of 11-HSD.

Oxidation of corticosterone to 11-dehydrocorticosterone by 11 β -dehydrogenase occurred in widely distributed regions of the brain. Activities were highest in the hippocampus and cortex (Lakshmi *et al.*, 1991; Moisin *et al.*, 1990a), an observation that was confirmed by immunohistochemical staining of brain regions with 11 β -dehydrogenase antibody (Lakshmi *et al.*, 1991; Roussseau *et al.*, 1972) and by *in situ* hybridization (Moisin *et al.*, 1990a) using cDNA corresponding to rat liver 11 β -dehydrogenase (Agarwal *et al.*, 1989).

There is as yet no direct experimental evidence to show that 11-HSD is the selection mechanism for brain receptor. Correlation of 11-HSD activity and intensity of immunoreactive labeling is consistent with a protective mechanism. Using neuronal and glial markers to measure the distribution of 11-HSD-like antigen, it was found that 11-HSD was distributed in the hippocampus in the CA1-4 regions and the dentate gyrus. The distribution in the hippocampus and cortex coincided with the distribution of MR. Neuronal 11-HSD was found throughout the cell body and its projections. Consistent with the hypothesis that 11-

11SD is the selection mechanism for MR, receptor and enzyme were located within the same cell (Sakai *et al.*, 1990). The distribution was heterogeneous, with some neurons that contained MR showing no detectable 11-11SD immunoreactivity. The distribution studies suggest that the selection mechanism of 11-11SD protection of MR is related within the individual neuron. The observation that the enzyme is localized in the neuronal nuclei and the discovery that in all brain regions investigated 11-11SD is found in glial cells suggest that its functions are complex.

In the hippocampus, as in other brain regions, glucose- and mineralocorticoids must both be present in some crucial, though as yet unknown, relationship for optimal function to occur. A neuron containing both GR and MR must be able to manipulate both glucocorticoid and mineralocorticoid levels to permit functionally adequate binding to the available receptors. This may require that corticosterone concentrations be adjusted to permit its occupancy of MR and/or GR in a way that is in accord with the needs of the cell, or alternatively, to permit glucocorticoid metabolism to proceed extensively in order for the MR to bind aldosterone. How 11-11SD activity is controlled to permit optimal neuronal function is not known. Several mechanisms are possible: (a) controlled synthesis and inactivation of enzyme; (b) control of activity based on availability of cofactor; (c) reversibility of enzyme, permitting net oxidation or reduction of 11-oxygenated steroid to occur.

In some regions of the brain, 11-11SD may mediate GR-dependent events. Cerebellum contains no measurable MR, but does have well-defined GR; 11-11SD is expressed as high activity accompanied by high levels of 11-11SD mRNA (Moish *et al.*, 1990a). If 11-11SD serves any receptor-related function in cerebellum, it must only influence GR-dependent events. It has been suggested that 11-11SD may control glucose metabolism in the brain via GR. Inhibition of 11-11SD by glycyrrhetic acid increased steroid-dependent uptake of 2-¹⁴C-deoxyglucose in the arcuate nucleus, prople area, cortex, hippocampus, and paraventricular nucleus (Seckl *et al.*, 1991). Glial cells contain GR (McGinnis and de Vellis, 1981), but no MR, respond to glucocorticoids, and contain 11-11SD. There are thus several examples of cell types in which the resident 11-11SD may serve cell-specific functions depending on their receptor content.

E. Leydig Cells, Strauss, and 11-11SD

An extensive literature has accumulated that shows that the *luteal* synthesizes less testosterone when exposed to pharmacological levels

of circulating corticosteroid, and that the diminished responses are receptor mediated (Phillips *et al.*, 1989). Extending the idea first proposed for kidney function, it was suggested that 11-11SD protects the production of testosterone by Leydig cells against the inhibitory effects of glucocorticoids. That testicular 11-11SD is restricted to the Leydig cells is consistent with this hypothesis.

By inactivating cortisol (corticosterone in the rat) 11-11SD acts as an enzymatic barrier. This testicular barrier is overwhelmed when the level of circulating glucocorticoid exceeds a threshold defined by the enzyme's ability to oxidize the steroid. It was recently found that 11-11SD is absent from rat Leydig cells prior to the twenty-fifth postnatal day (Phillips *et al.*, 1989; Haider *et al.*, 1990). This observation suggested that prior to 25 days of age, corticosterone cannot be inactivated and thus contributes to the prepubertal suppression of testosterone production. Subsequently, as enzyme is expressed and corticosteroid is oxidized, inhibition of testosterone is overcome. As the animal ages, it is only possible to inhibit testosterone production with amounts of cortisol and corticosterone that exceed the oxidative capacity of 11-11SD, or by glucocorticoid analogs, such as dexamethasone, that are poor substrates for the enzyme. The 11-11SD inhibitor carbenoxolone increases the testosterone suppressive effects of corticosterone, an effect in accord with predictions (Abayasekara *et al.*, 1990). The synthetic glucocorticoid dexamethasone inhibits testosterone secretion by Leydig cells, but since, unlike corticosterone, it is not a substrate of 11-11SD, its effect is not increased by 11-11SD inhibitors (Monder *et al.*, 1992). The mineralocorticoid aldosterone has no effect on testosterone production (Abayasekara *et al.*, 1990; Monder *et al.*, 1992), since Leydig cells have no MR (R. R. Sakai, M. Hardy, and C. Monder, unpublished observations).

F. Mammary Gland

In the mammary gland, glucocorticoids are required for the synthesis of casein, lactalbumin, and other proteins, through a GR-dependent process (Jahn *et al.*, 1987). Quirk *et al.* (1990a) have found 11-11SD in the epithelial and adipose tissue of pregnant and lactating mammary gland of rats. The enzyme is 20-fold higher in adipocytes than epithelial cells and diminishes in both cell types as pregnancy progresses to reach low levels in lactating glands. The authors propose that 11-11SD decreases local concentration of corticosterone by the formation of the inactive 11-dehydrosteroid metabolite, and thus prevents premature milk production (Quirk *et al.*, 1990a,b). The presence of MR in breast tissue (Quirk *et al.*, 1993) suggests that the role of 11-11SD in

the mammary gland may involve the participation of corticosteroids on salt and water metabolism (Mollna *et al.*, 1990) as well as on milk protein production.

X. Epilogue

In this article, we have attempted to provide a historical perspective into the conceptual evolution of 11 β -hydroxysteroid dehydrogenase from its pedestrian origin as an enzyme that catalyzes reversible inactivation of corticosteroids to its currently more prestigious role as mediator of steroid-receptor interactions. The recent surge of interest in 11- β SD was powered by two factors. The first was the recognition of clinical disorders whose symptomatology could be rationalized as being due to defects in 11- β SD expression. The second was the development of the tools—antibodies, cDNA—that facilitated exploration of the enzyme at the molecular level. With the use of these probes, investigators have developed hypotheses implicating the interplay of corticosteroids, 11- β SD, and steroid receptors in the etiology of juvenile hypertension, and in normal renal function. The success of this endeavor has inspired further exploration of the possibility that the principle that emerged from the study of the kidney also apply to other organs. The current state of these investigations has been summarized in this article. The significance of 11- β SD as a mediator of steroid-receptor interaction cannot, however, be unique. The concepts emerging from studies with this corticosteroid-metabolizing enzyme ought to apply as well to enzymes participating in the metabolism of other classes of steroid. Thus, there is reason to believe that the answers emerging from the questions posed in this article may find application elsewhere in steroid biology.

ACKNOWLEDGMENTS

We thank Maria I. New and Phyllis Speiser for their contributions to this work, and Dr. Cedric Shekleton for his help in compiling Table VIII. Original research described was supported by U.S. Public Health Services Grants DK 37094 and DK 42169 and March of Dimes Grant 6-609.

REFERENCES

- Abeysekera, D. R. E., Iland, A. M., and Cooke, D. A. (1990). Inhibition of Leydig cell steroidogenesis by adrenal steroids. Specifically, time dependency and effects of an 11 β -hydroxysteroid dehydrogenase inhibitor. *J. Endocrinol.* 124, Suppl. Abstr 73.
- Abrahamovitz, M., Dinnendahl, C. L., and Murphy, B. E. P. (1992). Cortisol-cortisone interconversions in human fetal lung: Contrasting results using explant and monolayer cultures suggest that 11 β -hydroxysteroid dehydrogenase (EC 1.1.1.46) comprises two enzymes. *J. Clin. Endocrinol. Metab.* 64, 653-659.
- Abrahamovitz, M., Gertler, R., and Murphy, B. E. P. (1984). Investigation of factors influencing 11 β -hydroxysteroid dehydrogenase (EC 1.1.1.46) activity in midgestation human fetal lung monolayer and explant cultures. *J. Steroid Biochem.* 21, 677-683.
- Adam, W. R., Punder, J. W., Mercer, J., and Ullick, S. (1978). Amplification of the action of aldosterone by 5 α -dihydrocortisol. *Endocrinology (Baltimore)* 103, 465-471.
- Agarwal, A. L., Mondel, C., Eckstein, B., and White, P. G. (1989). Cloning and expression of rat cDNA encoding corticosteroid 11 β -dehydrogenase. *J. Biol. Chem.* 264, 18939-18943.
- Agarwal, A. K., Tsai-Luna, M. T., Mondel, C., and White, P. G. (1990). Expression of 11 β -hydroxysteroid dehydrogenase using recombinant vaccinia virus. *Mol. Endocrinol.* 4, 1827-1832.
- Althaus, Z. R., Bailey, J. R., Leskey, J. E. A., and Slikker, W. Jr. (1982). Transplacental metabolism of dexamethasone and cortisol in late gestational age of rhesus monkey (Macaca mulatta). *Dev. Pharmacol. Ther.* 8, 332-349.
- Ameling, D., Hubner, H. J., and Roke, L. (1953). Enzymatische Umwandlungen von Steroiden III Über die Einführung der 11-Oxygruppe und die Reduktion der 11-Keto-Gruppe. *Hoppe-Seyler's Z. Physiol. Chem.* 304, 36-48.
- Ameling, D., Hubner, H. J., Roke, L., and Meyerheim, G. (1953b). Conversion of corticosterone to compound F. *J. Clin. Endocrinol. Metab.* 13, 1125.
- Anderson, D. F., Buck, M. K., and Rankin, J. H. G. (1979). Placental transfer of dexamethasone in near term sheep. *J. Dev. Physiol.* 1, 431-436.
- Armenini, D., Karbowiak, L., and Punder, J. W. (1983). Affinity of liquorite derivatives for mineralocorticoid and glucocorticoid receptors. *Clin. Endocrinol. (Oxford)* 19, 609-612.
- Armenini, D., Straaser, T., and Weber, P. G. (1983). Characterization of aldosterone binding sites in circulating human mononuclear leukocytes. *Am. J. Physiol.* 246, E338-E339.
- Armenini, D., Wehling, M., and Weber, P. G. (1989a). Mineralocorticoid effector metabolism of liquorite derivatives in human mononuclear leukocytes. *J. Endocrinol. Invest.* 12, 303-306.
- Armenini, D., Scall, M., Zennaro, M. G., Karbowiak, L., Wallace, C., Lewitch, B., Vesel, P., and Mianero, F. (1989b). The pathogenesis of pseudohyperaldosteronism from carbonoxolone. *J. Endocrinol. Invest.* 12, 337-341.
- Arriza, J. L. (1991). Aldosterone action: Perspectives from the cloning of the mineralocorticoid receptor. *Colloq.-Int. Natl. Suisse Rech. Med.* 316, 13-21.
- Arriza, J. L., Weinberger, C., Cerrelli, G., Ghier, T. M., Handelin, B. L., Jonsson, D. E., and Evans, R. M. (1987). Cloning of human mineralocorticoid receptor complementary DNA: Structural and functional kinship with the glucocorticoid receptor. *Science* 237, 268-276.
- Arriza, J. L., Simerly, R. B., Swanson, L. W., and Evans, R. M. (1988). The neuronal mineralocorticoid receptor as a mediator of glucocorticoid responses. *Neuron* 1, 887-900.
- Ashkin, N., and Cook, C. (1953). *In vivo* observations of the effects of cortisone upon blood vessels in rabbit ear chambers. *Br. J. Exp. Pathol.* 33, 445-450.
- Avery, M. E. (1976). Pharmacological approaches to the acceleration of fetal lung maturation. *Br. Med. J.* 31, 13-17.
- Bingli, S., Albrecht, E. D., and Type, G. J. (1990). Regulation of 11 β -hydroxysteroid

- dehydrogenase activity in baboon placenta by estrogen. *Endocrinology (Baltimore)* 120, 2742-2748.
- Bailey, E., and West, H. F. (1969). The secretion, interconversion and catabolism of cortisol, cortisone and some of their metabolites in man. *Acta Endocrinol. (Copenhagen)* 42, 339-359.
- Baird, C. W., and Bush, I. E. (1960). Cortisol and cortisone content of amniotic fluid from diabetic and non diabetic women. *Acta Endocrinol. (Copenhagen)* 34, 97-104.
- Baker, M. E. (1969). Human placental 17 β -hydroxysteroid dehydrogenase (homologous to NADH protein of *Rhizobium meliloti*). *Mol. Endocrinol.* 3, 681-684.
- Baker, M. E. (1970a). Sequence similarity between *Pseudomonas* dihydroliol dehydrogenase, part of the gene cluster that metabolizes polychlorinated biphenyls and dehydrogenase involved in metabolism of ribitol and glycerol and synthesis of antibiotics and 17 β -oestradiol, testosterone and corticosterone. *Biochem. J.* 207, 639-641.
- Baker, M. E. (1970b). A common ancestor for human placental 17 β -hydroxysteroid dehydrogenase, *Streptomyces coelicolor* act III protein, and *Phenylalanine methyltransferase* hol dehydrogenase. *FASEB J.* 4, 222-226.
- Barton, D. N., and Greene, R. J. (1965). Action of compounds with effective in vivo mineralocorticoid activity on ion transport in leucocytes. *Br. J. Clin. Pharmacol.* 21, 27-34.
- Barron, J. H. (1963). Side effects of carbonazone. *Acta Gastroenterol. Belg.* 46, 469-484.
- Barzeghian, O., Levine, R., and Eppa, P. (1962). Direct effect of cortisol and cortisone on insulin and glucagon secretion. *Endocrinology (Baltimore)* 111, 1648-1651.
- Bellata, M. C., Mendonca, D. B., Kater, C. E., Arnold, I. J. P., Rocha, A., Nicolau, W., and Molde, W. (1966). Spironolactone-reversible nickels associated with 11 β -hydroxysteroid dehydrogenase deficiency syndrome. *J. Pediatr.* 109, 889-893.
- Bouillon, E. F., and Jayle, M. F. (1957). Etude de l'équilibre entre 11 β -hydroxy et 11-tétrahydroxy urinaires après administration de cortisone, cortisol et ACTH A. *Thieme. Bull. Soc. Chim. Biol.* 39, 37.
- Baxter, J. D., Schambelan, M., Matulich, D. T., Spindler, B. J., Taylor, A. A., and Bartter, F. C. (1976). Aldosterone receptors and the evaluation of plasma mineralocorticoid activity in normal and hypertensive states. *J. Clin. Invest.* 66, 579-589.
- Beaumont, K., and Faneuil, D. D. (1963). Characterization of rat brain aldosterone receptors reveals high affinity for corticosterone. *Endocrinology (Baltimore)* 113, 2043-2051.
- Beilman, I. Z., Bayard, F., Ance, I. O., Kowarski, A., and Migeon, C. J. (1972). The transplacental passage of prednisone and prednisolone in pregnancy near term. *J. Pediatr.* 81, 836-845.
- Berliner, D. L. (1965). Studies of the mechanisms by which cells become resistant to steroids. *Cancer Res.* 25, 1085-1090.
- Bierlmeier, D. L., and Ruhmann, A. G. (1966). Comparison of the growth of *Rhizobium* under the influence of 11 β -hydroxy and 11-keto compounds. *Endocrinology (Baltimore)* 79, 373-382.
- Bernal, A., Lopez, and Graf, I. L. (1961). Corticosteroid metabolism in vitro by human placenta, fetal membranes and decidua in early and late gestation. *Placenta* 2, 279-286.
- Bernal, A., Lopez, and Turnbull, A. G. (1965). Cortisol metabolism in human placenta and decidua in relation to enzyme. *Ann. N.Y. Acad. Sci.* 127, 107.
- Bernal, A., Lopez, Piant, A. P. F., Anderson, A. H. M., and Turnbull, A. G. (1968). 11 β -hydroxysteroid dehydrogenase activity (E.C. 1.1.1.146) in human placenta and decidua. *J. Steroid Biochem.* 13, 1081-1087.
- Bernal, A., Lopez, Anderson, A. B. M., and Turnbull, A. G. (1962). The lack of influence of labor on human placental 11 β -hydroxysteroid dehydrogenase activity. *J. Clin. Endocrinol. Metab.* 64, 1261-1264.
- Bigger, J. F., Fahnberg, P. F., and Becker, B. (1972). Increased cellular sensitivity to glucocorticoids in primary open angle glaucoma. *Invest. Ophthalmol.* 11, 832-837.
- Birmingham, M. K., Slumpf, W. S., and Sar, M. (1970). Nuclear localization of aldosterone in rat brain cells assessed by autoradiography. *Experientia* 26, 1240-1241.
- Blair-West, J. R., Coghan, D. A., Denton, D. A., Goding, J. R., Munro, J. A., Peterson, R. E., and Winlow, M. (1962). Humoral stimulation of adrenal cortical secretion. *J. Clin. Invest.* 41, 1600-1627.
- Blumck, W. (1968). Die α -ketoacidien cortisol-und corticosteronmetaboliten sowie die 11-oxo und 11-oxo-17-ketosterole im urin von Kindern. *Acta Endocrinol. (Copenhagen)*, Suppl. 134, 9-112.
- Bohn, M. C., Howard, E., Vitek, U., and Kozswaki, Z. (1991). Glial cells express both mineralocorticoid and glucocorticoid receptors. *J. Steroid Biochem. Mol. Biol.* 40, 105-111.
- Boland, E. W. (1962). Antihypertensive effects of hydrocortisone (free alcohol), hydrocortisone acetate and cortisone (free alcohol) as compared with cortisone acetate. *Br. Med. J.* 1, 658-664.
- Bonvalet, J. P. (1991). Aldosterone-sensitive cells in the kidney: New insights. *Neuro Physiol. Sci.* 6, 201-205.
- Bonvalet, J. P., Dujardin, J., Bial-Chabaud, M., Pradelle, P., and Fermand, N. (1990). Distribution of 11 β -hydroxysteroid dehydrogenase along the rabbit nephron. *J. Clin. Invest.* 86, 832-837.
- Borrel, J. G., Van Loth, S. P., De Vries, L. A., and Mulhysen, J. A. (1993). Syncretic action of liquorice and cortisone in Addison's and Simmonds' disease. *Lancet* 1, 657-663.
- Bradlow, H. L., Zimmet, B., Gallagher, T. F., and Hellman, L. (1968). Tetrahydrocortisol metabolism in man. *Steroids* 12, 303-308.
- Brem, A. S., Matheson, K. L., Conca, T., and Horita, D. J. (1969). Effect of carbonazone on glucocorticoid metabolism and Na transport in toad bladder. *Am. J. Physiol.* 267, F700-F704.
- Brem, A. S., Matheson, K. L., Barnes, J. L., and Horita, D. J. (1991). 11-dehydrocortisone, a glucocorticoid metabolite, inhibits aldosterone action in toad bladder. *Am. J. Physiol.* 261, F873-F879.
- Bro-Rasmussen, F., Buus, O., and Trolle, D. (1962). Ratio of cortisone/cortisol in mother and infant at birth. *Acta Endocrinol. (Copenhagen)* 40, 679-683.
- Brown-Sequard, C. E. (1956). Recherche expérimentale sur la physiologie des capsules surrénales. *C. R. Hebd. Seances Acad. Sci.* 43, 422.
- Burslein, S., Savard, K., and Dorfman, R. I. (1963). The in vivo metabolism of hydrocortisone. *Endocrinology (Baltimore)* 63, 88-97.
- Burton, A. F. (1965). Inhibition of 11 β -hydroxysteroid dehydrogenase activity in rat and mouse in vitro and in vivo. *Endocrinology (Baltimore)* 77, 325-331.
- Burton, A. F., and Anderson, F. H. (1968). Inactivation of corticosteroids in intestinal mucosa by 11 β -hydroxysteroid NADP oxidoreductase (EC 1.1.1.146). *Am. J. Gastroenterol.* 76, 627-631.

- Edwards, C. R. W., Stewart, P. M., Burt, D., Brett, L., McIntyre, M. A., Salama, W. S., de Kloet, E. R., and Mondet, C. (1981). Localization of 11β -hydroxysteroid dehydrogenase-11 α -specific protector of the mineralocorticoid receptor. *Lancet* 2, 986-989.
- Edwards, C. R. W., Burt, D., and Stewart, P. M. (1983). The specificity of the human mineralocorticoid receptor: Clinical clues to a biological conundrum. *J. Steroid Biochem.* 32, 213-216.
- Elk-Nes, K. B., and Brizze, K. R. (1963). Concentration of tritium in brain tissue of dogs given $11,2311$ corticoid intravenously. *Biochim. Biophys. Acta* 97, 320-333.
- Ellers, E. A., and Peterson, R. E. (1964). Aldosterone secretion in the rat. In "Aldosterone" (E. F. Hollen and J. Reibel, eds.), pp. 251-264. Blackwell, Oxford.
- Ellenberg, A. H. (1962). Steroid compounds: Isolating from incubation of cortisone with surviving liver slices. *Science* 116, 520-521.
- Elmadjian, F., Ilope, J. M., and Pincus, G. (1960). The action of mono-ammonium glycerinate on adrenocortical nuclei and the synergism with hydrocortisone. *J. Clin. Endocrinol. Metab.* 16, 338-349.
- Endoh, G. I., and Kochakian, C. D. (1962). Partial purification and further characterization of the triphosphopyridine nucleotide specific G_{12} - 17β -hydroxysteroid dehydrogenase of guinea pig liver. *Biochim. Biophys. Acta* 62, 245-250.
- Endoh, G. I., Kochakian, C. D., and Hamm, D. (1960). Separation of a triphosphopyridine nucleotide-specific from a diphosphopyridine nucleotide-specific 17β -hydroxysteroid dehydrogenase of guinea pig liver. *J. Biol. Chem.* 235, 2792-2796.
- Engel, L. L., Carter, P., and Fielding, L. L. (1955). Urinary metabolites of administered corticosterone. I. Steroid liberated by glucuronidase hydrolysis. *J. Biol. Chem.* 213, 99-106.
- Eppstein, M. T., Enginer, E. A., Donah, R. A., Hughes, H., Cowles, R. J., and Lam, S. (1978). Licorice raises urinary corticoid in man. *J. Clin. Endocrinol. Metab.* 47, 397-400.
- Eriksson, H., and Gustafsson, J. A. (1971). Metabolism of corticosterone in the isolated perfused rat liver. *Eur. J. Biochem.* 20, 231-236.
- Ernst, L., and Jones, L. C. (1962). A study of nucleotide tri- and diphosphate activities of rat liver microsomes. *J. Cell Biol.* 15, 563-578.
- Evans, R. M. (1988). The steroid and thyroid hormone receptor superfamily. *Science* 240, 889-895.
- Farmah, N., Vandewalle, A., and Bonvolet, J. P. (1983). Autoradiographic determination of desamethasone binding sites along the rabbit nephron. *Ann. J. Physiol.* 244, F325-F331.
- Farmah, N., Ohlin, M. E., Lamhon, M., Ikehaya, F., Wenzel, H. M., Bonvolet, J. P., and (1982). Immunolocalization of glucocorticoid and mineralocorticoid receptors in rabbit kidney. *Am. J. Physiol.* 240, C226-C233.
- Fazekas, A. G., Sander, T., and Janhies, A. (1970). Conversion of corticosterone to 11β -dehydrocorticosterone by adrenal gland preparations of different animal species. *Endocrinology (Baltimore)* 89, 438-440.
- Flemer, L. F., and Flemer, M. (1953). "Steroids." Reinhold, New York.
- Fletcher, W., Olson, B. J., Monahan, L. A. H., Hanover, J. W., and van Muijster, P. J. J. (1982). Low-renin, low-aldosterone hypertension and abnormal cortisol metabolism in a 18-month-old child. *J. Form. Res.* 19, 107-114.
- Fish, C. A., Hysano, M., and Pincus, G. (1953). Conversion of cortisone to 17β -hydroxycorticosterone by liver homogenates. *Arch. Biochem. Biophys.* 42, 480-481.

- Forman, B. H., and Mulrow, P. J. (1973). Effect of corticosteroids on water and electrolyte metabolism. In "Handbook of Physiology" (R. O. Greep and E. B. Astwood, eds.), Sect. 7, Vol. 6, pp. 179-180. Am. Physiol. Soc., Washington, DC.
- Fregly, M. J., and Rowland, N. E. (1985). Role of renin-angiotensin-aldosterone system in NaCl appetite of rats. *Am. J. Physiol.* 248, R1-R11.
- Freeman, H. L., Sherry, A., Haseenzadeh, M., Freeman, M., Cilly, P., and Morgan, D. (1980). Control of cell proliferation in human gliomas by glucocorticoids. *Br. J. Cancer* 41, 887-889.
- Pukuhila, D. K., Gallagher, T. F., Greenberg, W., and Pearson, O. H. (1960). Studies with an adrenal inhibitor in adrenal carcinoma. *J. Clin. Endocrinol. Metab.* 20, 1234-1240.
- Punder, J. W. (1980). Adrenocortical receptors in the brain. In "Frontiers in Neuroendocrinology" (W. F. Gunning and L. Martin, eds.), Vol. 9, pp. 169-189. Raven Press, New York.
- Punder, J. W. (1982). Adrenal steroids: New answers, new questions. *Science* 237, 235.
- Punder, J. W. (1980a). Target tissue specificity of mineralocorticoids. *TEM* 1, 145-148.
- Punder, J. W. (1980b). Corticosteroid receptors and renal 11β -hydroxysteroid dehydrogenase activity. *Semin. Nephrol.* 10, 311-319.
- Punder, J. W., Lawrence, J. T., Smith, H., and Smith, A. I. (1988). Mineralocorticoid action: Target tissue specificity is enzyme, not receptor, mediated. *Science* 242, 583-585.
- Punder, J. W., Pearce, P. T., Smith, R., and Campbell, J. (1989). Vascular Type I aldosterone binding sites are physiological mineralocorticoid receptors. *Endocrinology (Baltimore)* 125, 2224-2226.
- Punder, J. W., Pearce, P. T., Myler, K., and Roy, L. P. (1990). Apparent mineralocorticoid excess, pseudohypoaldosteronism and urinary electrolyte excretion: Toward a redefinition of mineralocorticoid action. *FASEB J.* 4, 3234-3238.
- Punkshewer, J. D., Veary, K. J., Mockridge, P. R., and Hughes, E. R. (1978). Distribution of dexamethasone between mother and fetus after maternal administration. *Pediatr. Res.* 12, 1063-1066.
- Purvison, M. M., and MacPhee, G. B. (1976). Kinetic study of 11β -hydroxysteroid dehydrogenase in rat submandibular salivary gland. *Arch. Oral Biol.* 20, 241-245.
- Regehr, H. J., Edwards, C. R. W., and Rossetti, B. C. (1989). Steroid Metabolism determines mineralocorticoid specificity in the testis. *Am. J. Physiol.* 257, F690-F695.
- Gallagher, T. F., Jellman, L., Zano, R., and Miller, D. O. (1983). Steroid hormone metabolism in chronic myelogenous leukemia. *Blood* 28, 743-748.
- Gerlach, J., and McEwen, B. S. (1972). Rat brain binds adrenal steroid hormone: Autoradiography of hippocampus with corticosterone. *Science* 175, 1133-1136.
- Thom, D., Wicks, C. M., (1981a). Dux, W. L., Erman, M., Himmey, R. L., and Orr, J. C. (1981b). Three-dimensional structure of holo 3α , 20β -hydroxysteroid dehydrogenase: A member of a short-chain dehydrogenase family. *Proc. Natl. Acad. Sci. U.S.A.* 78, 10064-10068.
- Graf, R., Vetter, U., Zandvold, J. M., and Schriber, H. (1976a). Organ specific ontogenesis of steroid metabolizing activities in the rat. 11β and 17β hydroxysteroid dehydrogenase. *Acta Endocrinol. (Copenhagen)* 79, 192-201.
- Thiel, R., Hoef, J. G., Iaz, E. R., and Schriber, H. (1976b). Enzyme activity in kidney, adrenal and gonadal tissue of rats treated neonatally with androgen or estrogen. *Endocrinology (Baltimore)* 87, 317-326.
- Glanopoulos, G. (1974). Uptake and metabolism of cortisone and cortisol by fetal rabbit lung. *Steroids* 29, 845-853.

- Chenopoulou, G., Jackson, K., and Tulechinsky, D. (1982). Glucocorticoid metabolism in human placenta, decidua, myometrium and fetal membranes. *J. Steroid Biochem.* 17, 371-374.
- Clayton, H. J., Linnemann, G. L., Hwang, C. L., and Kree, H. L. (1980). Endocrine involvement in luteal hyperandrogenism. *Am. J. Physiol.* 199, 718-720.
- Gomez-Sanchez, E. P. (1991). What is the role of the central nervous system in mineralocorticoid hypertension? *Am. J. Hypertens.* 4, 374-381.
- Gordon, Q. O., and Souliere, A. L. (1971). Thyroid hormone effects on steroid hormone metabolism. *Bull. N.Y. Acad. Med.* 45, 241-259.
- Gottfried, H. (1964). The occurrence and biological significance of steroids in lower vertebrates. *Steroids* 3, 219-242.
- Grey, C. H., Greenaway, J. M., Holmes, N. J., and Shaw, D. A. (1962). Metabolism of Cortisol-4-C¹⁴ in a patient with Cushing's Syndrome. *J. Endocrinol.* 24, 199-214.
- Graesser, B. L. (1960). 11 β -hydroxysteroid metabolism by mouse brain and plasma. *J. Neurochem.* 13, 475-478.
- Grimmer, H. L., and Axelrod, L. H. (1968). Conversion of cortisol to cortinol acetate, cortisone acetate and cortisone by the developing primate brain. *Steroids* 11, 827-836.
- Grimmer, H. L., and Unger, F. (1964). Conversion of pregnenolone and 4-androstene-3 β , 17 β -diol to testosterone by mouse testes *in vitro*. *Steroids* 3, 67-76.
- Grunfeld, J.-P., and Eloy, L. (1987). Glucocorticoids modulate vascular reactivity in the rat. *Hypertension* 10, 608-618.
- Gugnier-de Maeyer, J. A., Crigler, J. F., and Told, N. L. (1963). An alteration in cortisol metabolism in patients with Cushing's syndrome and bilateral adrenal hyperplasia. *J. Clin. Endocrinol. Metab.* 23, 1271-1284.
- Gunderson, H. M., and Nordlie, R. C. (1973). Carbanion phosphate: Glucose phosphate transferase and glucose-6-phosphate phosphohydrolyase of nuclear membrane. Interrelationships between membrane integrity, enzyme latency and catalytic behavior. *J. Biol. Chem.* 248, 3562-3569.
- Gustafsson, J. A., and Gustafsson, S. A. (1974). Delayed expression of neonatal and differentiation of corticosteroid patterns in rat bile. *Eur. J. Biochem.* 44, 225-233.
- Gustafsson, J. A., and Stenberger, A. (1976). Obligatory role of hypophysis in sexual differentiation of hepatic metabolism in rats. *Proc. Natl. Acad. Sci. U.S.A.* 73, 1462-1465.
- Haidler, S. G., Pissia, D., and Linnemann, F. G. (1980). Histoclinical demonstration of 11 β -hydroxysteroid dehydrogenase as a marker for Leydig cell maturation in rat. *Acta Histochem. Suppl.* 28, 203-207.
- Haleigh, R. M., and Jones, C. T. (1980). Effect of glucocorticoids on alpha 1-adrenergic receptor binding in rat vascular smooth muscle. *J. Mol. Endocrinol.* 5, 41-48.
- Hammami, M. M., and Sitter, P. K. (1980). Cortisol resistance and altered 11 β -hydroxysteroid dehydrogenase activity. *Abstr. 72nd Annu. Meet. Endocr. Soc. Abstr.* 1026, p. 281.
- Hammami, M. M., and Sitter, P. K. (1991). Regulation of 11 β -hydroxysteroid dehydrogenase activity in human skin fibroblasts. Enzymatic modulation of glucocorticoid action. *J. Clin. Endocrinol. Metab.* 73, 326-334.
- Hanington, R., Tait, S. A. S., and Tait, J. F. (1970). *In vitro* effects of ACTH, angiotensin, serotonin and potassium on steroid output and conversion of cortisone to aldosterone by isolated adrenal cells. *Endocrinology (Baltimore)* 87, 1141-1167.

- Hartnack, H. L. J., Van Dammelen, P., Van Seiter, A. F., and Noolenar, A. J. (1981). Apparent mineralocorticoid excess and deficient 11 β -oxidation of cortisol in a young female. *Clin. Endocrinol. (Oxford)* 21, 605-614.
- Huismann, W., and Turney, A. L. (1989). Clinical and biochemical effects of carbamazepine. In "A Symposium on Carbamazepine Sodium" (J. M. Johnson and F. M. Sullivan, eds.), pp. 169-172. Butterworth, London.
- Ilayoshi, T., Nakai, T., Uchida, K., Morimoto, S., and Takeda, R. (1984). The characteristics of renal mineralocorticoid receptors in glycyrrholic acid or dexamethasone-induced hypertensive rats. *Clin. Exp. Hypertens—Theory Pract.* A6, 1625-1640.
- Ilechner, O., Zedler, A., Jacobsen, R. F., Levy, H., Jeanloz, R. W., Schenker, V., and Piccus, G. (1981). The nature and biosynthesis of the adrenal secretory product. *Recent Prog. Horm. Res.* 6, 215-241.
- Ilechner, O., Solomon, M. M., and Camp, E. (1983). Corticosteroid metabolism in liver. Studies on perfused rat liver. *Endocrinology (Baltimore)* 63, 202-215.
- Ilklick, H. J., and Hargreaves, J. W. (1970). Pseudoaldosteronism (Liddle's syndrome): Evidence for increased cell membrane permeability to Na⁺. *Endocrinol. Res.* 456.
- Ilklick, H. J., and Simone, K. (1976). Solubilization of membranes by detergent. *Biochim. Biophys. Acta* 416, 29-79.
- Ilklick, H. J., Hargreaves, J. W., Zurnoff, B., and Gallagher, T. F. (1981). The influence of thyroid hormone on hydrocortisone production and metabolism. *J. Clin. Endocrinol. Metab.* 21, 1231-1247.
- Ilklick, H. J., Castello, R., Kobayashi, N., and Fromm, M. (1980a). Sites and significance of renal corticosteroid metabolism. *Int. Congr. Ser.—Endocrinol.* 877, 67-76.
- Ilklick, H. J., Siebe, H., and Fromm, M. (1980b). Inhibition of 11 β -hydroxysteroid dehydrogenase and its effect on epithelial sodium transport. *Kidney Int.* 38, 673-678.
- Ilklick, H. J., Ilklick, H., and Yershel, F. H. (1981). Target cell metabolism of corticosteroids modulating antidiuretic effect. In "Nephrology" (M. Italiano, ed.), pp. 1122-1126. Springer-Verlag, Tokyo.
- Ilklick, H. J., M., and Chrambach, A. (1981). Solubilization of functional membrane-bound receptors. In "Receptor Biochemistry and Molecular Biology" (J. Craig Venter and L. C. Harrison, eds.), Vol. 1, pp. 35-46. Liss, New York.
- Ilklick, H. J., Chrambach, R., Ralbe, M., and Schreier, H. (1973). Ontogenesis von 11 β -hydroxysteroid-Dehydrogenase-Aktivitäten in der Ratte. *Leber. Hoppe-Seyler's Z. Physiol. Chem.* 354, 306-311.
- Ilklick, H. J., Brown, E. M., Jansen, R. A., and Brown, C. Y. (1981). Hydrocortisone and cortisone injected into arthritic joints. Comparative effects of and use of hydrocortisone as a local anti-arthritis agent. *JAMA. J. Am. Med. Assoc.* 147, 1629-1636.
- Ilklick, H. N., Brook, R. L., and Devlin, J. M. (1974). Thylakoid corticosteroid metabolism in intact and adenohypophysectomized ducts. *Gen. Comp. Endocrinol.* 22, 417-427.
- Ilklick, H. J., Dillon, M. J., Levin, M., and Shub, V. (1983). Fetal, low renin hypertension associated with a disturbance of cortisol metabolism. *Arch. Dis. Child.* 68, 1018-1020.
- Ilklick, H. J., and Moller, M. (1977). Histochemistry of 11 β -hydroxysteroid dehydrogenase in rat submandibular gland. Effect of cortisol stimulation. *Histochem. J.* 9, 699-708.

- terization of 11 β -hydroxysteroid dehydrogenase gene expression: Identification of multiple unique forms of messenger ribonucleic acid in the rat kidney. *Endocrinology (Baltimore)* 127, 3008-3013.
- Krozowski, Z. S., Obeyesekere, V., Smith, R., and Mercer, W. (1992). Tissue-specific expression of an 11 β -hydroxysteroid dehydrogenase with a truncated N-terminal domain—a potential mechanism for differential intracellular localization within mineralocorticoid target cells. *J. Biol. Chem.* 267, 2659-2674.
- Kumamoto, J., Ralston, J. K., and Lyons, J. M. (1971). Temperature trends in Arrhenius plots. A thermodynamic consequence of a phase change. *J. Theor. Biol.* 31, 47-61.
- Lakehmi, V., and Mondor, C. (1985a). Extraction of 11 β -hydroxysteroid dehydrogenase from rat liver by detergent. *J. Steroid Biochem.* 22, 331-340.
- Lakehmi, V., and Mondor, C. (1985b). Evidence for independent 11-oxidase and 11-reductase activities of 11 β -hydroxysteroid dehydrogenase: Enzyme latency, phase transitions and lipid requirements. *Endocrinology (Baltimore)* 116, 662-666.
- Lakehmi, V., and Mondor, C. (1988). Purification and characterization of the corticosteroid 11 β -dehydrogenase component of the rat liver 11 β -hydroxysteroid dehydrogenase complex. *Endocrinology (Baltimore)* 123, 2390-2398.
- Lakehmi, V., Sakel, R., McEwen, B. S., and Mondor, C. (1991). Regional distribution of 11 β -hydroxysteroid dehydrogenase in rat brain. *Endocrinology (Baltimore)* 128, 1741-1748.
- Latif, S. A., Conca, T. J., and Norris, D. J. (1990). The effects of the licorice derivative, glycyrrhetic acid, on hepatic 3 α and 3 β -hydroxysteroid dehydrogenase and 5 α and 5 β -reductase pathway of metabolism of aldosterone in male rats. *Steroids* 55, 52-58.
- Lax, E. R., Ghraf, R., and Schrieffers, H. (1978). The hormonal regulation of hepatic microsomal 11 β -hydroxysteroid dehydrogenase activity in the rat. *Acta Endocrinol. (Copenhagen)* 80, 352-358.
- Lax, E. R., Ghraf, R., Schrieffers, H., and Volgt, K. H. (1979). The involvement of the thyroid and adrenal in the regulation of enzyme activities of hepatic and renal steroid metabolism in the rat. *Hoppe-Seyler's Z. Physiol. Chem.* 300, 137-143.
- Lee, S.-M. K., Chiehol, M. A., and Katz, A. L. (1983). Corticosteroid binding sites along the rat nephron. *Am. J. Physiol.* 245, F604-F609.
- Leitz, T., and Reinbohl, R. (1987). The biosynthesis of 11-ketotestosterone by the testis of the Siamese fighting fish, *Betta splendens* (Anabantidae, Poeciliidae). *Gen. Comp. Endocrinol.* 68, 145-157.
- Lavitz, M., Jansen, V., and Jancia, J. (1978). The transfer and metabolism of corticosteroids in the perfused human placenta. *Am. J. Obstet. Gynecol.* 132, 363-366.
- Liddle, G. W., Blicke, T., and Coppage, W. S. (1963). A familial renal disorder simulating primary aldosteronism but with negligible aldosterone secretion. *Trans. Assoc. Am. Physicians* 76, 199-211.
- Minkin, O. C. (1976). Adrenocortical fetal malnutrition syndrome in the fetus. *Am. J. Obstet. Gynecol.* 128, 681-689.
- Migawa, G. C., and Howie, R. N. (1972). A controlled trial of antepartum glucocorticoid treatment for prevention of the respiratory distress syndrome in premature infants. *Pediatrics* 50, 515-526.
- Logg, M. A., and Nicholson, T. E. (1978). The effect of dexamethasone on the activity of 11 β -hydroxysteroid dehydrogenase in the foetal rabbit lung during the final stages of gestation. *J. Pharm. Pharmacol.* 30, 687-688.
- Muecklenstein, M. A., Ilofengier, W. H. L., Jansen, R. W. M., Hamerling, T. J., and Kluybenburg, P. W. G. (1980). The influence of glycyrrhetic acid on plasma cortisol

- and cortisone in healthy young volunteers. *J. Clin. Endocrinol. Metab.* 70, 1637-1643.
- Nashab, V. B., and Ulrich, F. (1960). Metabolism of cortisol and cortisone by various tissues and subcellular particles. *J. Biol. Chem.* 235, 356-360.
- Marek, L., Krok, M., and Jernvall, J. (1990). Fetal-specific 20 β -hydroxysteroid dehydrogenase is an enzyme for the short-chain, non-metabolizable alcohol dehydrogenase type. *FEBS Lett.* 266, 61-64.
- Morver, T., and Edelman, I. S. (1976). Dihydrocortisol, a potent mineralocorticoid. *J. Steroid Biochem.* 8, 1-7.
- Naseon, H. L. (1950). Isolation of adrenal cortical hormones from urine: 17 β -hydroxycorticosterone and 17 β -hydroxy-11-dehydrocorticosterone. *J. Biol. Chem.* 182, 131-149.
- Masters, J. N., Finch, C. E., and Sapolsky, R. M. (1989). Glucocorticoid enhancement of hippocampal neurons does not involve deoxyribonucleic acid cleavage. *Endocrinology (Baltimore)* 124, 3083-3088.
- Mattings, D., Tyler, C., and Bilton, E. (1970). Plasma 11-hydroxycorticoid levels after cortisone sodium. *Br. Med. J.* 3, 486-501.
- McDonald, I. H., Thum, K. A., and Evans, J. (1988). Glucocorticoids in the blood plasma of the platypus *Ornithorhynchus anatinus*. *J. Endocrinol.* 116, 407-416.
- McEwen, B. S., Jambolin, L. T., Rainbow, T. C., and De Nicola, A. F. (1986). Aldosterone effects on salt appetite in adrenalectomized rats. *Neuroendocrinology* 43, 38-43.
- McGinnis, J. F., and de Vellis, J. (1981). Cell surface modulation of gene expression in brain cells by down regulation of glucocorticoid receptors. *Proc. Natl. Acad. Sci. U.S.A.* 78, 1298-1292.
- Meigs, R. A., and Engel, J. L. (1961). The metabolism of adrenocortical steroids by human tissues. *Endocrinology (Baltimore)* 69, 152-162.
- Mercer, W. H., and Krozowski, Z. S. (1992). Localization of an 11 β hydroxysteroid dehydrogenase activity to the distal nephron. Evidence for the existence of two species of dehydrogenase in the rat kidney. *Endocrinology (Baltimore)* 130, 640-643.
- Meyer, W. J., and Nichols, N. H. (1981). Mineralocorticoid binding in cultured smooth muscle cells and fibroblasts from rat aorta. *J. Steroid Biochem.* 14, 1167-1168.
- Michaud, N. J., and Burton, A. F. (1977). Maternal fetal relationships in corticosteroid metabolism. *Biol. Neonate* 32, 132-137.
- Miller, L. L., and Axelrod, J. R. (1953). Cortisone metabolism in the perfused normal and experimental cirrhotic rat liver. *Metab. Clin. Exp.* 3, 439-448.
- Mittels, H., Vagstad, A., and Goodman, A. D. (1967). A syndrome resembling primary aldosteronism but without mineralocorticoid excess. *Clin. Res.* 15, 482.
- Mitchell, H. F., Storon-Forté, M., Hems, D. L., and Jaffe, R. B. (1981). Cortisol production and metabolism in the late gestation rhesus monkey fetus. *Endocrinology (Baltimore)* 109, 916-924.
- Mitchell, B. F., Storon-Forté, M., and Jaffe, R. B. (1982). Cortisol-cortisone interrelationships in the late gestation rhesus monkey fetus in vitro. *Endocrinology (Baltimore)* 111, 1837-1842.
- Miyake, S., Kimbata, R., and Hatanaka, T. (1973). Metabolism and conjugation of cortisol by various dog tissues in vitro. *J. Steroid Biochem.* 4, 567-576.
- Mogilkevich, M., and Heynaud, J. P. (1980). Evidence for a specific mineralocorticoid receptor in rat pituitary and brain. *J. Steroid Biochem.* 12, 309-314.
- Molteni, M. J., Seckl, J. R., Mondor, C., Agnew, A. K., White, P. C., and Edwards, C. R. W. (1990a). 11 β -hydroxysteroid dehydrogenase in brain: mRNA expression and bioactivity in rat cerebellum. *J. Neuroendocrinol.* 2, 853-858.
- Molteni, M. J., Seckl, J. R., and Edwards, C. R. W. (1990b). 11 β -hydroxysteroid dehydro-

- genase bioactivity and messenger RNA expression in rat forebrain: localization in hypothalamus, hippocampus and cortex. *Endocrinology (Baltimore)* 127, 1450-1456.
- Molinar, M. P., Edwards, C. R. W., and Seckl, J. R. (1992). Ontogeny of 11 β -hydroxysteroid dehydrogenase in rat brain and kidney. *Endocrinology (Baltimore)* 130, 400-409.
- Molina, R., Fietta, X., Herranz, M., Fraile, M., Velasco, A., Zanon, O., Martinez-Obaya, M. J., and Ballate, A. M. (1990). Biochemistry of cyst fluid in fibrocystic disease of the breast. *Ann. N.Y. Acad. Sci.* 689, 29-42.
- Monder, C. (1991a). Heterogeneity of 11 β -hydroxysteroid dehydrogenase in rat tissues. *J. Steroid Biochem.* 40, 633-636.
- Monder, C. (1991b). Corticosteroids, receptors, and the organ-specific functions of 11 β -hydroxysteroid dehydrogenase. *FASEB J.* 5, 3047-3064.
- Monder, C. (1991c). Corticosteroids, kidneys, sweet taste and dirty drugs. *Mol. Cell. Endocrinol.* 78, C95-C98.
- Monder, C., and Hradlow, H. L. (1990). Corticoid action: Exploration at the frontier of corticosteroid metabolism. *Recent Prog. Horm. Res.* 36, 305-400.
- Monder, C., and Lakshmi, V. (1989a). Evidence for kinetically distinct forms of corticosteroid 11 β -dehydrogenase in rat liver microsomes. *J. Steroid Biochem.* 32, 77-83.
- Monder, C., and Lakshmi, V. (1989b). Corticosteroid 11 β -hydroxysteroid dehydrogenase activities in vertebrate liver. *Steroids* 52, 515-528.
- Monder, C., and Lakshmi, V. (1990). Corticosteroid 11 β -dehydrogenase of rat tissues: Immunological studies. *Endocrinology (Baltimore)* 126, 2435-2443.
- Monder, C., and Shackleton, C. H. L. (1984). 11 β -hydroxysteroid dehydrogenase: Fact or fancy? *Steroids* 44, 383-417.
- Monder, C., and White, A. (1983). Purification and properties of a sheep liver 21-hydroxysteroid dehydrogenase. *J. Biol. Chem.* 258, 767-774.
- Monder, C., and White, A. (1986). The 21-hydroxysteroid dehydrogenase of liver: A NADP⁺ dehydrogenase and two NAD⁺ dehydrogenases. *J. Biol. Chem.* 261, 71.
- Monder, C., Shackleton, C. H. L., Hradlow, H. L., New, M. L., Stoner, E., Jahan, F., and Lakshmi, V. (1988). The syndrome of apparent mineralocorticoid excess: Its association with 11 β -dehydrogenase and 6 β -reductase deficiency and some consequences for corticosteroid metabolism. *J. Clin. Endocrinol. Metab.* 63, 560.
- Monder, C., Stewart, P. M., Lakshmi, V., Valentino, R., Burt, D., and Edwards, C. R. W. (1989). Licorice inhibits corticosteroid 11 β -dehydrogenase of rat kidney and liver: In vivo and in vitro studies. *Endocrinology (Baltimore)* 125, 1046-1053.
- Monder, C., Lakshmi, V., and Miroff, Y. (1991). Kinetic studies on rat liver 11 β -hydroxysteroid dehydrogenase. *Biochim. Biophys. Acta* 1115, 23-29.
- Monder, C., Sakai, R. R., Blanchard, R. J., Blanchard, D. G., Lakshmi, V., Miroff, Y., Phillips, D. M., and Hradlow, H. L. (1993). The mediation of reticular function by 11 β -hydroxysteroid dehydrogenase. In *Stress and Reproduction* (J. W. Punder and J. H. Doublich, eds.), Raven Press, New York (in press).
- Moura, A. M., and Worcel, M. (1984). Direct action of aldosterone on transmembrane ²²Na efflux from arterial smooth muscle. *Hypertension* 9, 425-430.
- Munch, A., and Laing, K. (1977). Glucocorticoid receptors and mechanisms of action. In *Receptors and Mechanisms of Action of Steroid Hormones* (J. P. Magalhães, ed.), pp. 311-397. Dekker, New York.
- Murphy, B. E. P. (1977a). Chloride membrane as an extra-adrenal source of fetal cortisol in human amniotic fluid. *Nature (London)* 266, 179-181.


- Murphy, B. E. P. (1977b). Conversion of cortisol to cortisone by the human uterus and its reversal in pregnancy. *J. Clin. Endocrinol. Metab.* 44, 1214-1217.
- Murphy, B. E. P. (1978). Cortisol production and inactivation by the human lung during gestation and infancy. *J. Clin. Endocrinol. Metab.* 47, 243-248.
- Murphy, B. E. P. (1979a). The influence of serum proteins on the metabolism of cortisol by the human placenta. *J. Steroid Biochem.* 10, 387-392.
- Murphy, B. E. P. (1979b). Cortisol and cortisone in human fetal development. *J. Steroid Biochem.* 11, 609-613.
- Murphy, B. E. P. (1981). Ontogeny of cortisol-cortisone interconversion in human tissues: A role for cortisone in human fetal development. *J. Steroid Biochem.* 14, 811-817.
- Murphy, B. E. P. (1982). The absorption by the human fetus of intra-amniotically injected cortisol. *J. Steroid Biochem.* 16, 416-417.
- Murphy, B. E. P. (1991). Steroids and depression. *J. Steroid Biochem. Mol. Biol.* 38, 637-659.
- Murphy, B. E. P., and Branchaud, C. T. L. (1983). Fetal metabolism of cortisol. *Curr. Top. Exp. Endocrinol.* 6, 197-220.
- Murphy, B. E. P., and Bieschoux, R. C. (1972). Steroid levels in the human fetus: Cortisol and cortisone. *J. Clin. Endocrinol. Metab.* 35, 678-683.
- Murphy, B. E. P., and Veldy, D. (1981). Specificity of human 11 β -hydroxysteroid dehydrogenase. *J. Steroid Biochem.* 14, 807-809.
- Murphy, B. E. P., and Veldy, D. L. (1982). Radioenzyme assay for some substrates and inhibitors of human placental 11 β -hydroxysteroid dehydrogenase. *J. Immunology* 129, 17-30.
- Murphy, B. E. P., Clark, S. J., Donald, I. R., Flinkey, M., and Veldy, D. (1974). Conversion of maternal cortisol to cortisone during placental transfer to the human fetus. *Am. J. Obstet. Gynecol.* 118, 638-641.
- Naray-Felcs-Tóth, A., and Felcs-Tóth, G. (1990). Glucocorticoid receptors mediate mineralocorticoid-like effects in cultured collecting duct cells. *Am. J. Physiol.* 259, F672-F678.
- Naray-Felcs-Tóth, A., Wallington, C. D., and Felcs-Tóth, G. (1991). 11 β -hydroxysteroid dehydrogenase activity in the renal target cells of aldosterone. *Endocrinology (Baltimore)* 129, 17-21.
- Nelson, D. L., Semmel, L. T., Willardson, D. G., and Tyler, F. H. (1981). The levels of 17-hydroxycorticosteroids in peripheral blood of human subjects. *J. Clin. Endocrinol.* 11, 1021-1029.
- New, M. L., and Levine, L. S. (1977). An unidentified ACTH-stimulable adrenal steroid in childhood hypertension. In *Juvenile Hypertension* (M. L. New and L. S. Levine, eds.), pp. 143-163. Raven Press, New York.
- New, M. L., Levine, L. S., Biglieri, E. O., Pareira, J., and Ulick, S. (1977). Evidence for an unidentified steroid in a child with apparent mineralocorticoid hypertension. *J. Clin. Endocrinol. Metab.* 44, 924-933.
- New, M. L., Oberfield, S. E., Carey, R., Grigg, F., Ulick, S., and Levine, L. S. (1982). A genetic defect in cortisol metabolism as the basis for the syndrome of apparent mineralocorticoid excess. *Semin. Symp.* 60, 86-101.
- Nguyen-Thong-Thuan, R., and Burton, A. F. (1971). The uptake and metabolism of 3H-corticosterone and fluorimetrically determined corticosterone in tissues of several mouse strains. *Biol. Membr.* 16, 78-84.
- Nichols, T. E., and Kim, Y. A. (1976). The metabolism of 3H cortisone- and 3H-cortisol by the isolated perfused rat and guinea pig lungs. *Steroids* 25, 387-402.

- Nichols, T. E., and Lugg, M. A. (1982). The physiological significance of 11 β -hydroxysteroid dehydrogenase in the rat lung. *J. Steroid Biochem.* 17, 113-118.
- Nichols, N. R., Olsson, C. A., and Punder, J. W. (1983). Steroid effects on protein synthesis in cultured smooth muscle cells from rat aorta. *Endocrinology (Baltimore)* 113, 1096-1101.
- Nichols, N. R., McNally, M., Campbell, J. H., and Punder, J. W. (1984). Overlapping but not identical protein synthetic domains in cardiovascular cells in response to glucocorticoid hormones. *J. Lipids* 2, 663-669.
- Nichols, N. R., Nguyen, H. H., and Meyer, W. J. (1985). Physical separation of corticoid receptors with Type I and Type II specificities. *J. Steroid Biochem.* 22, 677-682.
- Noma, J., Ileyelli, N., and Sekiba, K. (1991). Automated direct HPLC assay for estradiol, estrone, and corticosterone in serum and amniotic fluid. *J. Chromatogr.* 669, 35-41.
- Oberfeld, S. E., Levine, L. S., Carey, R. M., Greig, F., Ullick, S., and New, M. L. (1983). Metabolic and blood pressure responses to hydrocortisone in the syndrome of apparent mineralocorticoid excess. *J. Clin. Endocrinol. Metab.* 58, 332-339.
- Ojima, M., Satoh, K., Gombuch, T., Itoh, N., Kim, S., Fukuchi, S., and Miyachi, Y. (1990). The inhibitory effects of glycyrrhizin and glycyrrhetic acid on the metabolism of corticoid and prednisolone. *Nippon Koshonji Gakai Zasshi* 69, 584-596.
- Olson, R. E., Thayer, S. A., and Kopp, L. J. (1944). The glycoconjugate activity of certain crystalline steroids of the adrenal cortex when administered singly and with corticoid extract to fasted, normal, and adrenalectomized rats. *Endocrinology (Baltimore)* 36, 464-472.
- Ong, J., Kerr, D. I. B., Copper, H. R., and Johnston, G. A. (1990). Cortisone: A potent GABA_A antagonist in the guinea pig isolated ileum. *J. Pharm. Pharmacol.* 42, 662-664.
- Oono, K., Drave, E. L., and Threlk, R. C. (1979). Sodium, extracellular fluid volume and cardiac output changes in the genesis of mineralocorticoid hypertension in the intact dog. *Hypertension* 1, 331-336.
- Oshaki, P. A. (1960). Steroid 11 β -dehydrogenase in human placenta. *Nature (London)* 187, 777.
- Pal, S. B. (1967). Pattern of the excretion of urinary corticoid, tetrahydrocortisone, allopregnenolone, and tetrahydrocortisol in normal human individuals and in patients with rheumatic disorders. *Clin. Chim. Acta* 16, 67-67.
- Pasqualini, J. R., Nguyen, B. J., Uhlrich, F., Winkler, N., and Jicafany, E. (1970). Cortisol and cortisone metabolism in the human fetal-placental unit at mid-gestation. *J. Steroid Biochem.* 1, 209-219.
- Pasqualini, J. R., Costa-Novoa, S. C., Ito, Y., and Nguyen, B. J. (1970). Hepatocortisol-cortisone conversion in the fetal tissue and subcellular fractions of fetal and adult guinea pig liver. *J. Steroid Biochem.* 1, 341-347.
- Pepo, G. J. (1979). The production and secretion of corticoid by baboon neonates. *Steroids* 33, 261-260.
- Pepo, G. J., and Albrecht, E. D. (1984a). Comparison of corticoid-cortisone interconversion *in vitro* by the human and baboon placenta. *Steroids* 44, 229-239.
- Pepo, G. J., and Albrecht, E. D. (1984b). Transplacental metabolism of corticoid and cortisone during mid- and late gestation in the baboon. *Endocrinology (Baltimore)* 116, 1946-1951.
- Pepo, G. J., and Albrecht, E. D. (1985a). The effects of cortisone on the interconversion of corticoid and cortisone in the baboon. *J. Steroid Biochem.* 23, 276-278.
- Pepo, G. J., and Albrecht, E. D. (1985b). Transplacental corticosteroid metabolism during baboon pregnancy. In "Research in Perinatal Medicine. IV. Perinatal Endocrinology" (E. D. Albrecht and G. D. Pepo, eds.), p. 201. Perinatology Press, New York.
- Pepo, G. J., and Albrecht, E. D. (1987). Fetal regulation of transplacental corticoid-cortisone metabolism in the baboon. *Endocrinology (Baltimore)* 120, 2528-2533.
- Pepo, G. J., and Townley, J. D. (1976). The metabolism of corticoid by term baboon neonates (*Phoca populi*). *Endocrinology (Baltimore)* 99, 456-459.
- Pepo, G. J., Waddell, B. J., Stahl, S. J., and Albrecht, E. D. (1988). The regulation of transplacental corticoid-cortisone metabolism by estrogen in pregnant baboons. *Endocrinology (Baltimore)* 122, 78-83.
- Perschel, F. H., Dohler, H., and Hertzler, K. (1991). Bile acids and their amides inhibit 11 β -hydroxysteroid dehydrogenase obtained from rat kidney. *Pluggers Arch.* 418, 536-543.
- Peterson, N. A., Chalkoff, L. L., and Jones, C. (1965). The *in vitro* conversion of cortisone by subcellular brain fractions of young and adult rats. *J. Neurochem.* 12, 273-276.
- Peterson, R. E., and Pierce, C. E. (1960). The metabolism of corticosterone in man. *J. Clin. Invest.* 39, 741-757.
- Peterson, R. E., Wengert, J. R., Guerra, S. L., Brodie, B. B., and Bunim, J. J. (1969). The physiological disposition and metabolic fate of hydrocortisone in man. *J. Clin. Invest.* 34, 1779.
- Phillips, G., and Higgins, B. A. (1965). A new defect in the peripheral conversion of corticoid to corticoid. *J. Steroid Biochem.* 22, 435-436.
- Phillips, D. M., Takahashi, Y., and Mondor, C. (1989). Corticosteroid 11 β -dehydrogenase in rat testis. *Endocrinology (Baltimore)* 125, 209-216.
- Pinder, R. M., Brogan, R. N., Sawyer, P. R., Speight, T. M., Spencer, R., and Avery, Q. S. (1976). Carbenoxolone: A review of its pharmacological properties and therapeutic efficacy in peptic ulcer disease. *Drugs* 11, 245-307.
- Porter, G. A. (1970). Synergistic effect of carbenoxolone sodium on aldosterone-enhanced active sodium transport in toad skin. In "A Symposium on Carbenoxolone Sodium" (J. H. Baron and F. M. Sullivan, eds.), pp. 33-47. Butterworth, London.
- Quirk, S. J., Gannell, J. E., and Punder, J. W. (1983). Aldosterone binding sites in pregnant and lactating rat mammary glands. *Endocrinology (Baltimore)* 113, 1812-1817.
- Quirk, S. J., Slatery, J., and Punder, J. W. (1990a). 11 β -hydroxysteroid dehydrogenase activity in the mammary gland. *J. Steroid Biochem.* 38, 623-626.
- Quirk, S. J., Slatery, J. A., and Punder, J. W. (1990b). Epithelial and adipose cells isolated from mammary glands of pregnant and lactating rats differ in 11 β -hydroxysteroid dehydrogenase activity. *J. Steroid Biochem. Mol. Biol.* 37, 639-644.
- Ratson, J. K., Lyons, J. M., Mehlhorn, R. J., and Keith, A. D. (1971). Temperature-induced phase changes in mitochondrial membranes detected by spin labeling. *J. Biol. Chem.* 246, 4036-4040.
- Rajin, S. (1972). Reconstitution of biological membranes. *Biochim. Biophys. Acta* 256, 241-266.
- Ramirez, J. C., Levine, L. S., Gunzler, P., Zancanato, G., Ramirez, L. C., Raub, W., Roeder, A., Undrow, H. L., and New, M. L. (1979). A syndrome of mineralocorticoid excess associated with defects in the peripheral metabolism of corticoid. *J. Clin. Endocrinol. Metab.* 49, 767-769.

- Reevers, F. (1948). Behaviour of the uvea ventriculi in uvea duodeni met. saucis II-quantitative. *Ned. Tijdschr. Geneesk.* 92, 2968-2971.
- Reich, H., Nelson, D. H., and Zafaroni, A. (1960). Isolation of 17-hydroxycorticosterone from blood obtained from adrenal veins of dogs. *J. Biol. Chem.* 187, 411-417.
- Reichstein, T., and Shoppe, C. W. (1943). The hormones of the adrenal cortex. *Vitam. Horm. (N.Y.)* 1, 346-413.
- Reul, J. M. H. M., and de Kloet, E. R. (1985). Two receptor systems for corticosterone in rat brain: Microdistribution and differential occupation. *Endocrinology (Baltimore)* 117, 2505-2511.
- Ropoff, J. M., and Stewart, G. N. (1972). The influence of adrenal extracts on the survival period of adrenalectomized dogs. *Science* 68, 327-328.
- Romano, L. P., Rodriguez, R. M., Seelye, J. M., and Tincin, O. (1967). Identification of tetrahydrocortisol and tetrahydrocorticosterone in the urine of normal and eclampsic men. *J. Clin. Endocrinol. Metab.* 17, 777-785.
- Roseblum, J. M., Yamada, T., Callard, G. V. (1985). Validation of radioimmunoassay systems for the measurement of 11-keto- and 11 β -hydroxy-testosterone in teleost blood. *Comp. Biochem. Physiol. B* 82D, 659-665.
- Rousseau, G. Q., Baxter, J. D., and Tomkins, G. M. (1972). Glucocorticoid receptors: Relations between steroid binding and biological effects. *J. Mol. Biol.* 67, 99-116.
- Rousseau, R., Baxter, J. D., Punder, J. W., Edelman, I. S., and Tomkins, G. M. (1972). Glucocorticoid and mineralocorticoid receptors for aldosterone. *J. Steroid Biochem.* 3, 219-227.
- Rundel, S. E., Punder, J. W., Lakhani, V., and Monder, C. (1989a). The intrarenal localization of mineralocorticoid receptors and 11 β -dehydrogenase: Immunocytochemical studies. *Endocrinology (Baltimore)* 125, 1700-1704.
- Rundel, S. E., Smith, I. A., Stockman, D., and Punder, J. W. (1989b). Immunocytochemical demonstration of mineralocorticoid receptors in rat and human kidney. *J. Steroid Biochem.* 33, 1215-1242.
- Sakai, R. R., Lakhani, V., Monder, C., Punder, J. W., Krasnowski, Z., and McEwen, B. S. (1980). Colocalization of 11 β -hydroxysteroid dehydrogenase and type I receptor immunoreactivity in the rat brain. *Soc. Neurosci. Abstr.* 19, Part 2, 1309.
- Sakal, R. R., and Mehl, A. Z. (1980). Corticosteroids and their role in the extrarenal electrolyte secreting organs of nonmammalian vertebrates. In "Steroids and Their Mechanism of Action in Nonmammalian Vertebrates" (G. Delio and J. Wrabetz, eds.), pp. 33-49. Raven Press, New York.
- Sander, T., Mehl, A. Z., and Faucher, A. O. (1977). Corticosteroid binding macromolecules in the salt activated nasal gland of the domestic duck *Anas platyrhynchos*. *Gen. Comp. Endocrinol.* 35, 348-359.
- Sander, T., Mehl, A. Z., and Dittmann, J. A. (1983). Further studies on the corticosteroid-receptor system of the nasal gland of the domestic duck (*Anas platyrhynchos*). *Gen. J. Biochem. Cell Biol.* 61, 731-743.
- Sang, G. W., Lorenzo, B., and Reidenberg, M. M. (1991). Inhibitory effects of gossypol on corticosteroid 11 β -hydroxysteroid dehydrogenase from guinea pig kidney: A possible mechanism for causing hypokalemia. *J. Steroid Biochem. Mol. Biol.* 39, 169-176.
- Sapolsky, R. M., and Pauline, W. A. (1986). Glucocorticoid potentiate ischemic injury in neurons: Therapeutic implications. *Science* 229, 1381-1387.
- Savage, D. C. L., Forsyth, C. G., McGafferty, E., and Cameron, J. (1976). The excretion of

- individual adrenocortical steroids during normal childhood and adolescence. *Acta Endocrinol. (Copenhagen)* 79, 551-567.
- Saward, K., Burelein, S., Rosenkrantz, H., and Dorfman, R. J. (1983). The metabolism of adrenosterone in vivo. *J. Biol. Chem.* 258, 717-725.
- Schultz, W., Kobayashi, N., Siebs, H., and Illersholzer, K. (1987). 11 β -hydroxysteroid dehydrogenase (11 β -HSD)—its function in renal corticosteroid metabolism. In "Molecular Nephrology: Biochemical Aspects of Kidney Function" (Z. Koyasovic and W. Guder, eds.), pp. 361-387. de Gruyter, Berlin.
- Schultz, W., U., and Speth, M. (1980). Investigation on the possible involvement of phospholipids in the glucose-6-phosphate transport system of rat liver microsomal glucose-6-phosphatase. *Eur. J. Biochem.* 109, 505-514.
- Sack, J. R., Kelly, P. A. T., and Sharkey, J. (1981). Glycinecortic acid, an inhibitor of 11 β -hydroxysteroid dehydrogenase, alters local cerebral glucose utilization in vivo. *J. Steroid Biochem. Mol. Biol.* 39, 777-779.
- Barnett, J. A., Brown, R. D., Inland, D. P., Yarbro, L. R., Watson, J. T., Slason, P. E., Hallfield, H. W., and Liddle, C. W. (1976). Evidence for a new mineralocorticoid in patients with low-renin essential hypertension. *Circ. Res.* 36, Suppl. 1, 2-9.
- Shackleton, C. H. L., Honour, J. W., Dillon, M. J., Chantler, G., and Jones, R. W. A. (1980). Hypertension in a 4 year old child: GC/MS evidence for deficient hepatic metabolism of steroids. *J. Clin. Endocrinol. Metab.* 60, 786-792.
- Shackleton, C. H. L., Rodriguez, J., Arceaga, E., Lopez, J. M., and Winter, J. S. D. (1985). Congenital 11 β -hydroxysteroid dehydrogenase deficiency associated with juvenile hypertension: Corticosteroid metabolite profiles of four patients and their families. *Clin. Endocrinol. (Oxford)* 22, 701-712.
- Sheppard, K., and Punder, J. W. (1987a). Mineralocorticoid specificity of renal type I receptors. *Am. J. Physiol.* 252, E224-E229.
- Sheppard, K., and Punder, J. W. (1987b). Type I receptors in parotid, colon, and pulmonary are aldosterone selective in vitro. *Am. J. Physiol.* 253, E467-E471.
- Shollon, L. J., Work, E. K., Jr., and MacGregor, J. (1985). Metabolism of corticoid 4- α -C and corticoid 4- β -C by rat brain homogenates. *Mol. Cell. Biochem.* 124, 1122-1127.
- Silker, W. J., Althaus, Z. R., Rowland, J. M., Hill, D. E., and Hendricks, A. O. (1982). Comparison of the transplacental pharmacokinetics of corticoid and triamcinolone acetonide in the rhesus monkey. *J. Pharmacol. Exp. Ther.* 223, 308-314.
- Slorber, R. B., Velliquette, G., and Abrams, O. M. (1989). Selective loss of hippocampal granule cells in the mature rat brain after adrenalectomy. *Science* 245, 635-638.
- Smith, R. T. (1978). The role of pulmonary corticosteroid 11-reductase activity in lung maturation in the fetal rat. *Endocrinol.* 102, 12-14.
- Smith, R. T., Torday, J. S., and Giroud, C. J. P. (1973). The growth promoting effect of corticoid on human fetal lung cells. *Steroids* 22, 518-524.
- Smith, R. T., Tanenwell, K. A., Minshall, D., Bogue, W. N., and Vrethen, E. (1982). Influence of corticosteroids on glycogen content and steroid 11-reductase activity in lung and liver of the fetal and newborn rat. *Biol. Neonate* 42, 201-207.
- Smith, R. E., and Punder, J. W. (1991). Renal 11 β -hydroxysteroid dehydrogenase activity: Effects of age, sex, and altered hormonal status. *J. Steroid Biochem. Mol. Biol.* 39, 265-267.
- Sources, G. W., and Morris, D. J. (1990). The effects of 11-dehydrocorticosterone (A) on the Na⁺ retaining actions of aldosterone. *Program, 72nd Annu. Meet. Endocr. Soc. Abstr.* 486, p. 146.
- Sources, G. W., and Morris, D. J. (1991). 11-dehydrocorticosterone (compound A) in the presence of calyculin is a more potent sodium reabsorber than its parent steroid

- CALC. MONDER AND VERHOEF C. WHITE:**
- corticosterone compound III. *Program 73rd Annu. Meet. Endocr. Soc. Abstr.* 604, p. 108.
- Sowell, J. G., Hagen, A. A., and Troop, R. C. (1971). Metabolism of corticosterone- ^{14}C by rat cortical ketone reductase. *Steroid* 18, 289-301.
- Lung, U. G., Mann, M. L., and Thewissen, K. B. (1961). Physiological effects of cortisone and ACTH in man. *J. Biol. Chem.* 236, 316-326.
- Stegler, M., and Reichman, T. (1957). Dehydrocorticosterone (21-oxyprogesterone) and 5 α -pregnen-20-one. *Biochim. Biophys. Acta* 20, 1164-1179.
- Seiten, M. II., and Burnett, F. E. (1967). Some properties of variously activated microsome glucose-6-phosphatase, inorganic phosphatases and inorganic pyrophosphatase. Shift in pH optimum. *Biochim. Biophys. Acta* 139, 138-147.
- Gilbert, P. M., and Edwards, C. W. (1960). Specificity of the mineralocorticoid receptor. Critical role of 11 β -hydroxysteroid dehydrogenase. *JEMF* 1, 225-230.
- Stewart, P. M., and Edwards, C. W. (1961). The cortisol-cortisone shuttle and hypotension. *J. Steroid Biochem. Mol. Biol.* 40, 601-609.
- Stewart, P. M., Volante, A. M., Valentini, R., Butti, D., Shackleton, C. H. L., and Edelman, J. S. (1967). Mineralocorticoid activity of hydrocortisone: 11 β -hydroxysteroid dehydrogenase deficiency comes off age. *Lancet* i, 821-823.
- Stewart, P. M., Corrie, J. E. T., Shackleton, C. H. L., and Edwards, C. W. (1968). Syndrome of apparent mineralocorticoid excess. A defect in the cortisol-cortisone shuttle. *J. Clin. Invest.* 41, 340-349.
- Sheppard, M. C. (1991). Localization of renal 11 β -dehydrogenase by in situ hybridization. *Anatomical Record* 229, 2129-2136.
- Shahar, K., Takada, K., and Kalaykji, M. (1966). P450 μ -dependent conversion of corticosterone to cortisone, and 19-hydroxyandrostenedione to 19-oxoandrosterone. *Biochemistry Res. Commun.* 26, 369-375.
- Sweet, M. I., and Bryson, M. J. (1960). Role of phosphorylating nucleosides in the metabolism of cortisol by peripheral muscle. *Biochim. Biophys. Acta* 44, 217-223.
- Hughes, W. W., and Hemmington, J. W. (1944). The role of the adrenal cortex in physiological processes. *Physiol. Rev.* 24, 89-127.
- Takeda, R., Miyamoto, T., Suma, H., Matsuda, Y., and Ikeda, M. (1987). Glycyrrhetic acid as mineralocorticoid agonist. *J. Steroid Biochem.* 27, 845-849.
- Takemoto, D. J., Abel, J. L., Jr., and Allen, J. C. (1976). The metabolism of corticosterone in the salt gland of the duck, *Anas platyrhynchos*. *Gen. Comp. Endocrinol.* 26, 226-232.
- Tamura, Y., Nishikawa, T., Yamashita, K., Yamanaka, M., and Kumagai, A. (1979). Effects of glycerethinic acid and its derivatives on delta-4,5 alpha and 6 beta reductases in liver. *Astringin-Jorsch*, 29, 647-649.
- Tanford, C., and Reynolds, J. A. (1976). Characterization of membrane proteins in detergent solutions. *Biochim. Biophys. Acta* 467, 133-170.
- Tamplin, G. N., Agnew, A. K., Mondor, C., New, M. I., and White, P. C. (1991). The human gene for 11 β -hydroxysteroid dehydrogenase: Structure, tissue distribution and chromosomal localization. *J. Biol. Chem.* 266, 16553-16558.
- Tajima, A. K., Worthington, D. J., and Smith, R. T. (1977). Human adrenoleukodystrophy: Corticosteroid 11-oxidoreductase activity. *J. Clin. Endocrinol. Metab.* 46, 721-725.
- Pashine, V., Terui, M., Sob, H., Mizunuma, H., Kobayashi, H., and Murumori, F. (1999).
- Identical properties of aldosterone and corticosterone binders and their presence in fetal brain and kidney. *J. Neurosci.* (Tokyo) 19, 446-454.
- Taylor, N. P., Bartlett, W. A., Dawson, D. J., and Kinoshita, H. A. (1984). Corticosterone deficiency: Evidence for a new inborn error of metabolism of adrenal steroids. *J. Endocrinol. Suppl.* 102 Abstr. 50.
- Trudgel, B., Mehta, A. D. II., Butti, D., McIntyre, M. A., Pratt, L., and Edwards, C. W. (1990). Identification of hydrocortisone activity in skin by glycyrrhetic acid. *Lancet* ii, 1060-1063.
- Temple, D. L., and Libowitz, S. F. (1989). PVN steroid implants: Effect on feeding patterns and macronutrient selection. *Brain Res. Bull.* 23, 553-560.
- Thorn, C. W. (1944). Clinical use of extracts from adrenal cortex. *JAMA*, J. Am. Med. Assoc. 126, 10-15.
- Tomlin, G. M., and Leachman, K. J. (1965). Enzymatic reduction of cortisone. *J. Am. Chem. Soc.* 76, 3100-3101.
- Thorby, J. S., Smith, B. T., and Giroud, C. J. P. (1976). The rabbit fetal lung as a glucocorticoid target tissue. *Endocrinology* (Baltimore) 99, 1463-1467.
- Torday, J. S. (1980). Glucocorticoid dependence of fetal lung maturation in vitro. *Endocrinology* (Baltimore) 107, 839-844.
- Torday, J. S., Olson, E. B., Jr., and Piret, N. L. (1976). Production of cortisol from cortisol by the isolated perfused fetal rabbit lung. *Steroid* 27, 689-690.
- Tuchene, J. C., O'Hara, J. F., and Keipour, M. (1963). Cortisol from human nerve endings. *J. Clin. Invest.* 32, 1276-1279.
- Science 141, 1276-1279.
- Tuckey, A., Ausley, A., Hodgson, A., Luton, J. P., and Galan, P. (1984). 11-Hydroxy-11-ketosteroide equilibrium. A source of misinterpretation in steroid synthesis: evidence through the effects of trifenylamine on 11-hydroxysteroid dehydrogenase in sheep and human adrenals in vitro. *J. Steroid Biochem.* 20, 783-786.
- Turner, B. (1979). Steroid metabolism in fish. Identification of steroid metabolites of hydrolyzable conjugates of cortisol in the bile of trout *Salmo gairdnerii*. *Gen. Comp. Endocrinol.* 38, 196-206.
- Uye, I. M., and Burton, A. F. (1980). Variations in the pattern of metabolism of corticosterone in fetal mouse tissues. *J. Lipid Sci.* 26, 35-39.
- Ulrich, S., Ramirez, L. C., and New, M. I. (1971). An abnormality in steroid reductive cleavage in fetal mouse tissues. *J. Clin. Endocrinol. Metab.* 44, 799-802.
- Ulrich, S., Ramirez, L. C., and New, M. I. (1971). A syndrome of apparent mineralocorticoid excess associated with a hypernatremic hyponatremia. *N. Engl. J. Med.* 285, 121-122.
- Ulrich, S., Levine, L. S., Gunter, P., Zencaro, G., Ramirez, L. C., Rush, W., Foster, A., and Davidson, H. L. (1979). A syndrome of apparent mineralocorticoid excess associated with defects in the peripheral metabolism of cortisol. *J. Clin. Endocrinol. Metab.* 49, 757-764.
- Ulrich, S., Chou, C. K., Rao, K. N., Edassery, J., and Montero, F. (1989). A new form of the syndrome of apparent mineralocorticoid excess. *J. Steroid Biochem.* 32, 209-212.
- Ulick, S., Teddo, R., and Montero, F. (1990). Pathogenesis of the type 2 variant of the syndrome of apparent mineralocorticoid excess. *J. Clin. Endocrinol. Metab.* 70, 200-206.
- Ullmann, A., Menard, J., and Corvol, P. (1976). Binding of glycyrrhetic acid to kidney mineralocorticoid receptors. *Endocrinology* (Baltimore) 97, 46-51.
- Vinger, F., Guimille, R., and Schibler, R. W. (1978). Seasonal variation in adrenal 11 β -hydroxysteroid dehydrogenase in the meadow vole. *Gen. Comp. Endocrinol.* 36, 111-118.
- Van den Berg, D. T. W. M., deKloet, R. H., Van Dijken, H. II., and Dolong, W. (1990). Differential central effects of mineralocorticoids and glucocorticoids agonists and antagonists on blood pressure. *Endocrinology* (Baltimore) 126, 118-124.



| | | | | | | | |
|---------------|---|-------------------------------------|-----------|-----------|------------------|----------|-------|
| PubMed | Nucleotide | Protein | Genome | Structure | PopSet | Taxonomy | OMIM |
| Search PubMed | <input checked="" type="checkbox"/> for | | | | | Go | Clear |
| Links | Preview/Index | History | Clipboard | | | | |
| Display | Abstract | <input checked="" type="checkbox"/> | Save Text | Order | Add to Clipboard | | |

Entrez
PubMed☐ 1: *Endocr Res* 1996 Nov;22(4):793-801[Related Articles, Books, LinkOut](#)PubMed
Services

Endogenous 11 beta-hydroxysteroid dehydrogenase inhibitors and their role in glucocorticoid Na⁺ retention and hypertension.

Morris DJ, Souness GW

Department of Pathology and Laboratory Medicine, Miriam Hospital, Lifespan and Brown University School of Medicine, Providence, RI 02906, USA.


Related
Resources

11 beta-hydroxysteroid dehydrogenase (11 beta-HSD) metabolizes active glucocorticoids to their inactive 11-dehydro products and protects renal mineralocorticoid receptors from the high circulating levels of endogenous glucocorticoids. 11 beta-HSD has been suggested to be important not only in the control of renal sodium retention but also blood pressure. We had previously shown that 11 alpha- and 11 beta-hydroxyprogesterone (11 alpha- and 11 beta-OHP) were (I) potent inhibitors of 11 beta-HSD (Isoforms 1 and 2) activity in vitro, (ii) able to confer mineralocorticoid (MC) activity upon corticosterone (B) in vivo and (iii) hypertensinogenic when chronically infused into Sprague-Dawley (SD) rats. In addition we also showed that 3 alpha,5B-tetrahydroprogesterone (3 alpha,5B-THP) and chenodeoxycholic acid (CDCA) were potent inhibitors of 11 beta-HSD1 activity but not 11 beta-HSD2 activity, however, these substances were still able to confer MC activity upon B in the adrenalectomized rat. To assess the possible blood pressure modulating effects of 3 alpha,5B-THP and CDCA we have now infused these substances into intact SD rats continuously for 14 days. Both 3 alpha,5B-THP and CDCA caused a significant elevation in blood pressure within seven days, an effect that persisted throughout the 14-day infusion. These results show that both 3 alpha,5B-THP and CDCA are hypertensinogenic in the rat and that the inhibition of either 11 beta-HSD2 or 11 beta-HSD1 activity by endogenous progesterone metabolites and CDCA may be involved in the pathology of hypertension.

PMID: 8969942, UI: 97124806

| | | | | | |
|---------|----------|-------------------------------------|-----------|-------|------------------|
| Display | Abstract | <input checked="" type="checkbox"/> | Save Text | Order | Add to Clipboard |
|---------|----------|-------------------------------------|-----------|-------|------------------|

[Write to the Help Desk](#)



Entrez PubMed Nucleotide Protein Genome Structure OMIM PMC Journals Books

Search PubMed for

Limits Preview/Index History Clipboard Details

About Entrez

Citation Text

Text Version

Entrez PubMed

Overview
Help | FAQ
Tutorial
New/Noteworthy
E-Utilities

PubMed Services

Journals Database
MeSH Database
Single Citation Matcher
Batch Citation Matcher
Clinical Queries
LinkOut
Cubby

Related Resources

Order Documents
NLM Gateway
TOXNET
Consumer Health
Clinical Alerts
ClinicalTrials.gov
PubMed Central

Privacy Policy

☒ 1: J Clin Endocrinol Metab. 1995 Nov;80(11):3155-9.

[Related Articles, Links](#)

Carbenoxolone increases hepatic insulin sensitivity in man: a novel role for 11-oxosteroid reductase in enhancing glucocorticoid receptor activation.

Walker BR, Connacher AA, Lindsay RM, Webb DJ, Edwards CR.

University of Edinburgh, Department of Medicine, Western General Hospital, Scotland.

In the kidney, conversion of cortisol to cortisone by the enzyme 11 beta-hydroxysteroid dehydrogenase protects mineralocorticoid receptors from cortisol. In the liver, a different isoform of the enzyme favors 11 beta-reductase conversion of cortisone to cortisol. We have tested the hypothesis that hepatic 11 beta-reductase enhances glucocorticoid receptor activation in the liver by inhibiting the enzyme with carbenoxolone and observing effects on insulin sensitivity. Seven healthy males took part in a double blind randomized cross-over study in which oral carbenoxolone (100 mg every 8 h) or placebo was administered for 7 days. Euglycemic hyperinsulinemic clamp studies were then performed, including measurement of forearm glucose uptake. Carbenoxolone increased whole body insulin sensitivity (M values for dextrose infusion rates, 41.1 +/- 2.4 mumol/kg.min for placebo vs. 44.6 +/- 2.3 for carbenoxolone; P < 0.03), but had no effect on forearm insulin sensitivity. We infer that carbenoxolone, by inhibiting hepatic 11 beta-reductase and reducing intrahepatic cortisol concentration, increases hepatic insulin sensitivity and decreases glucose production. Thus, plasma cortisone provides an inactive pool that can be converted to active glucocorticoids at sites where 11 beta-reductase is expressed, abnormal hepatic 11 beta-reductase activity might be important in syndromes of insulin resistance, and manipulation of hepatic 11 beta-reductase may be useful in treating insulin resistance.

Publication Types:

- Clinical Trial
- Randomized Controlled Trial

MeSH Terms:






- 11-beta-Hydroxysteroid Dehydrogenases
- Adult
- Blood Glucose/metabolism
- Carbenoxolone/pharmacology*
- Cross-Over Studies
- Double-Blind Method
- Forearm/blood supply

- Human
- Hydroxysteroid Dehydrogenases/physiology*
- Insulin/blood
- Insulin/physiology*
- Liver/drug effects*
- Liver/physiology*
- Male
- Receptors, Glucocorticoid/physiology*
- Support, Non-U.S. Gov't

Substances:

- Blood Glucose
- Receptors, Glucocorticoid
- Insulin
- Carbenoxolone
- Hydroxysteroid Dehydrogenases
- 11-beta-Hydroxysteroid Dehydrogenases

PMID: 7593419 [PubMed - indexed for MEDLINE]

| | | | | | | |
|---|----------|--|--|---|------|---|
|  Display | Citation |  Show: 20 |  Sort |  Send to | Text |  |
|---|----------|--|--|---|------|---|

[Write to the Help Desk](#)
[NCBI](#) | [NLM](#) | [NIH](#)
[Department of Health & Human Services](#)
[Freedom of Information Act](#) | [Disclaimer](#)

May 12 2004 06:43:50

| | | | | | | | |
|---------|---------------|------------------------------|-----------|-----------|------------------|----------|------|
| NCBI | | National Library of Medicine | | PubMed | | | |
| PubMed | Nucleotide | Protein | Genome | Structure | PopSet | Taxonomy | OMIM |
| Search | PubMed | for | | Go | Clear | | |
| Limits | Preview/Index | History | Clipboard | | | | |
| Display | Abstract | Save | Text | Order | Add to Clipboard | | |

Entrez
PubMed☐ 1: *Endocrinology* 1993 Jun;132(6):2287-92

Related Articles, Books

PubMed
Services**Licorice inhibits 11 beta-hydroxysteroid dehydrogenase messenger ribonucleic acid levels and potentiates glucocorticoid hormone action.****Whorwood CB, Sheppard MC, Stewart PM**

Department of Medicine, University of Birmingham, Queen Elizabeth Hospital, Edgbaston, United Kingdom.

Related
Resources

11 beta-Hydroxysteroid dehydrogenase (11 beta HSD) is responsible for the interconversion of cortisol to cortisone [corticosterone (B) to 11-dehydrocorticosterone in rodents] and confers ligand specificity to the mineralocorticoid receptor. Inhibition of 11 beta HSD by licorice derivatives [glycyrrhizic and glycyrrhetic (GE) acids] results in cortisol/B and not aldosterone acting as a potent mineralocorticoid. 11 beta HSD is ubiquitously expressed and, by converting active glucocorticoid to inactive metabolites, may be an important prereceptor regulator of ligand access to the glucocorticoid receptor (GR). To investigate this further, we have studied the effect of 11 beta HSD inhibition by licorice derivatives on PRL gene expression (a known glucocorticoid target gene) in rat pituitary GH3 cells. Glycyrrhizic acid administration to rats in vivo (75 mg/kg.day for 5 days) resulted in inhibition of 11 beta HSD activity, as previously reported, but also a significant reduction in steady state 11 beta HSD mRNA levels in both predominantly mineralocorticoid (kidney and distal colon) and glucocorticoid (liver and pituitary) target tissues. In vitro, 11 beta HSD mRNA and activity were present in rat pituitary GH3 cells (81% conversion of B to 11-dehydrocorticosterone/4 x 10⁶ cells after 24-h incubation) and inhibited by GE in a dose-dependent fashion. While B or GE alone (10⁻⁸-10⁻⁵ M) had little or no effect on PRL mRNA levels or immunoassayable PRL, combinations of GE plus B resulted in marked inhibition of PRL mRNA levels and secretion, to such an extent that a concentration of 10⁻⁶ M B with 10⁻⁶ M GE was more potent than equimolar concentration of the synthetic GR agonist RU 28362. This inhibitory effect on PRL mRNA levels was blocked by a 10-fold excess of the GR antagonist RU 38486, but not by a 10-fold excess of the mineralocorticoid receptor antagonist RU 26752, confirming that this potentiation of glucocorticoid hormone action was operating through the GR and not the mineralocorticoid receptor. In addition to its established role as a competitive inhibitor of 11 beta HSD, licorice results in pretranslational inhibition of 11 beta HSD both in vitro and in vivo. 11 beta HSD is clearly an important mechanism in regulating tissue

levels of active glucocorticoid and, hence, ligand supply to the GR.

PMID: 8504732, UI: 93279206



[Write to the Help Desk](#)
[NCBI](#) | [NLM](#) | [NIH](#)
[Department of Health & Human Services](#)
[Freedom of Information Act](#) | [Disclaimer](#)

The Journal of
**Steroid
Biochemistry &
Molecular
Biology**



PERGAMON



Pergamon

0960-0760(94)E0010-I

J. Steroid Biochem. Molec. Biol. Vol. 49, No. 1, pp. 81-85, 1994
 Copyright © 1994 Elsevier Science Ltd
 Printed in Great Britain. All rights reserved
 0960-0760/94 \$7.00 + 0.00

Inhibition of 11β -Hydroxysteroid Dehydrogenase Obtained from Guinea Pig Kidney by Furosemide, Naringenin and Some Other Compounds

Yin Di Zhang,* Beverly Lorenzo and Marcus M. Reidenberg†

Departments of Pharmacology and Medicine, Cornell University Medical College, 1300 York Avenue, New York, NY 10021, U.S.A.

Inhibition of 11β -hydroxysteroid dehydrogenase (11β -OHSD) can cause excess mineralocorticoid effects and hypokalemia. Several substances causing hypokalemia (glycyrrhizic acid in licorice and gossypol) inhibit this enzyme. We tested other compounds for activity to inhibit 11β -OHSD in guinea pig kidney cortex microsomes with NADP as cofactor and cortisol as substrate. Furosemide was an inhibitor while bumetanide was not, indicating a mechanism for the increased K^+ excretion caused by furosemide compared with bumetanide. Naringenin (found in grapefruit juice), ethacrynic acid, and chenodeoxycholic acid had inhibitor IC_{50} values similar to glycyrrhizic acid. We conclude that various compounds can inhibit this enzyme and may play a role in K^+ metabolism and adrenocorticosteroid action.

J. Steroid Biochem. Molec. Biol., Vol. 49, No. 1, pp. 81-85, 1994

INTRODUCTION

The syndrome of apparent mineralocorticoid excess, first described by Ulick, Ramirez and New in 1977 [1], has led to much research on the enzyme 11β -hydroxysteroid dehydrogenase (11β -OHSD). Deficient activity of this enzyme in children leads to their inability to oxidize cortisol to inactive cortisone, providing high cortisol levels in the kidney which activate renal mineralocorticoid receptors and cause hypertension and hypokalemia. Subsequently, the mechanism of licorice-induced hypermineralocorticoidism was shown to be the inhibition of 11β -OHSD by the active principle of licorice, glycyrrhizic acid. Since then, much research has been done to explore the role that this enzyme plays in regulating the interactions of cortisol with mineralocorticoid and glucocorticoid receptors [2-6].

Gossypol, a polyphenolic constituent of cotton seed, has been studied in China as a potential oral contraceptive for men because it suppresses sperm motility

and formation without affecting testosterone levels [7]. Some Chinese men who received gossypol developed hypokalemia although the cause remained obscure [7]. This is particularly remarkable since idiopathic hypokalemia, often associated with hyperthyroidism, occurs widely in China; in addition, normal Chinese men have serum potassium levels lower than men in four other countries, with 9% having values below 3.5 mmol/l [8].

In studies investigating how gossypol causes hypokalemia, we found that gossypol inhibited 11β -OHSD activity in guinea pig [9] and human [10] renal cortical microsomes. We also found that certain bioflavonoids inhibit rat liver 11β -OHSD [10]. Others have reported inhibition of the rat kidney enzyme by bile acids [11] and by steroidal and triterpenoid compounds [12], and inhibition of the rat liver enzyme by some substances in human urine [13]. We therefore decided to test a variety of compounds for their possible enzyme inhibiting effect, choosing drugs that can cause hypokalemia or sodium retention as a side effect, flavonoids from grapefruit juice that inhibit the oxidation of dehydropyridine calcium channel blocking drugs [15-17] or sterols in vegetable oils at concentrations of 100-500 mg/dl [14].

*Present address: Nanjing Medical College, Add, 140, Han Zhong Road, Nanjing, Jiangsu, China

†Correspondence to M. M. Reidenberg, Department of Pharmacology.

Received 8 July 1993; accepted 4 Jan. 1994.

MATERIALS AND METHODS

Chemicals and solutions

Sitosterol was a gift from Eli Lilly and Co. (Indianapolis, IN). Campesterol was purchased from Aldrich Chemical Co. (Milwaukee, WI). All other chemicals were purchased from Sigma Co. (St Louis, MO).

Most sterols, furosemide, ethacrynic acid, naringin and naringenin were dissolved in ethanol and diluted with methanol. Cholic acid, chenodeoxycholic acid, bumetanide, hydrochlorothiazide and spironolactone were dissolved in methanol. Phenylbutazone and indomethacin were dissolved in distilled water (pH 9). Glycyrrhizic acid was dissolved in distilled water. Corticosterone and hydrocortisone were dissolved in methanol (144 $\mu\text{mol/l}$) and kept at -4°C .

Enzyme preparation and measurement of 11β -OHSD activity

Kidney cortex was obtained from long-haired male Hartley guinea pigs. Tissue was homogenized in Krebs-Henseleit buffer as described previously [9], except for the use of a Tekmar Tissuemizer (Cincinnati, OH). Microsomes were prepared as described previously [9], except that they were diluted to a concentration of 1.25 mg protein/ml prior to storage at -70°C .

The enzyme activity in guinea pig kidney cortex microsomes was determined by measuring the rate of conversion of cortisol to cortisone. Five minutes before incubation, 2 μl of concentrated Triton DF-18 was added to each milliliter of the microsome suspension. The assay mixture contained 500 μl Krebs-Henseleit buffer (pH 7.2), 50 μl 5 mmol/l NADP, 40 μl of 144 $\mu\text{mol/l}$ phosphate-sucrose buffer, 20–50 μl (25–63.5 μg) of microsome suspension in 0.01 M phosphate-sucrose buffer and various concentrations of each compound studied. This mixture was incubated in duplicate or triplicate. The total volume was 700 μl . Methanol concentration was kept at $<10\%$. Control studies showed that this concentration did not inhibit the reaction. After 1 h of incubation at 37°C , the reaction was terminated by the addition of 3 ml methylene chloride and 20 μl 144 $\mu\text{mol/l}$ corticosterone solution as the internal standard for assay of cortisone and cortisol.

The enzyme inhibition constant for furosemide was determined by adding furosemide in various amounts to achieve concentrations from 3.9 to 62 $\mu\text{mol/l}$ in the incubation mixture and cortisol concentrations of 4, 8, and 16 $\mu\text{mol/l}$. The constants were obtained from a Dixon plot and a kinetic program (Chou J, Chou T-C: Michaelis-Menton analysis with microcomputers, Disk No. 1, Elsevier-Biosoft, 1989, Cambridge, England).

A modification of the HPLC method of Sang [9] was used to measure cortisol, cortisone and corticosterone in the microsomal incubation mixture. The steroids were extracted into methylene chloride by vortexing for

1 min, then centrifuged at 750 g for 15 min. The aqueous layer was removed by aspiration. 300 μl of 0.1 NaOH was added to the organic phase followed by vortexing for 30 s. The mixture was centrifuged and the aqueous layer removed. The organic phase was washed with 500 μl of milli-Q water (Millipore Corp., Bedford, MA). The 1.5 ml organic phase was transferred to clean glass tubes and dried by evaporation in a 45 – 50°C water bath. The residue was dissolved into 200 μl of methanol and 5 μl of this solution was injected into the HPLC apparatus. A standard curve for cortisol and cortisone was determined in duplicate in each enzyme experiment by using the same amount of microsome suspension after boiling to inactivate the enzyme. Standard curves were plotted as the ratio of peak height of cortisone (or cortisol) divided by the peak height of the internal standard vs steroid concentration. All unknown concentrations of cortisol and cortisone were determined from the standard curves from each experiment. The drug concentrations that inhibited the enzyme by 50% (IC_{50}) were estimated from at least 3 different concentrations of each compound evaluated by a dose-response program (Chou and Chou: Dose-effect analysis with microcomputers, Disk No. 2, Elsevier-Biosoft, 1989, Cambridge, England).

The HPLC apparatus used for quantitating the steroids consisted of a Waters Model 6000 A solvent delivery system, U6K injector, model 680 automated gradient controller, Waters 486 tunable absorbance detector and a BBC chart recorder (Model SE 120). The mobile phase contained methanol-water (30:70, v/v) at a flow rate of 1.0 ml/min. The Waters stainless steel Novapak C_{18} column (3.9 \times 150 mm, 4 μ) was kept at room temperature. The retention times for cortisone, cortisol and corticosterone were 6.5, 7.0 and 9.0 min, respectively.

RESULTS

The efficacy of the compounds tested to inhibit the NADP-utilizing form of 11β -OHSD from guinea pig renal cortex with cortisol as substrate is shown in Tables 1 and 2. Furosemide was the most potent inhibitor tested, with glycyrrhizic acid, naringenin, ethacrynic acid and chenodeoxycholic acid having potencies similar to each other but an order of magnitude less potent than furosemide. Data for glycyrrhizic acid, naringenin and naringin are shown in Fig. 1. The correlation coefficient (r value) for the computed values agreeing with the measured values for the potent inhibitors was 0.99 for furosemide, glycyrrhizic acid, and naringenin, 0.96 for ethacrynic acid and 0.86 for chenodeoxycholic acid. It was above 0.95 for all of the other compounds tested except for phenylbutazone which was 0.86.

The observations of enzyme inhibition by furosemide at varying concentrations of cortisol is

Inhibition of 11β -OHSD

83

Table 1. Inhibition of 11β -OHSD by various compounds

| Compound | IC ₅₀ (μ mol/l) | Concentrations tested (μ mol/l) |
|-----------------------|---------------------------------|--------------------------------------|
| Furosemide | 59 | 12, 50, 100, 200, 500, 1000 |
| Glycyrrhizic acid | 254 | 132, 246, 529 |
| Naringenin | 336 | 12, 25, 50, 100, 1000, 2000, 5000 |
| Ethacrynic acid | 452 | 50, 100, 200, 400, 2000 |
| Chenodeoxycholic acid | 513 | 200, 400, 600, 800 |
| Phenylbutazone | 1358 | 167, 667, 1344 |
| Sitosterol | 1395 | 500, 1000, 1500 |
| Stigmasterol | 1968 | 500, 1000, 1500 |
| Naringin | 2373 | 582, 1163, 1744 |
| Cholic acid | 3529 | 1250, 2500, 3750, 5000 |

Campesterol inhibited 33% at the highest concentration tested of 1000 μ mol/l. Since a second higher point could not be measured because of limited solubility of the compound, an IC₅₀ was not calculated.

shown as a double reciprocal plot in Fig. 2. Most of the lines converge near the ordinate. A Dixon plot indicated that the inhibition by furosemide is competitive. The enzyme kinetic constants were: $K_m = 8 \mu$ mol/l and $V_{max} = 30$ nmol/ μ g microsomal protein/h. The K_i for furosemide was 7.7μ mol/l nearly the same as the K_m for cortisol.

DISCUSSION

We have tested a number of compounds for their ability to inhibit the NADP-utilizing form of 11β -OHSD from guinea pig renal cortex with cortisol as substrate. We found that furosemide is a much more potent inhibitor than glycyrrhizic acid, and that naringenin, ethacrynic acid and chenodeoxycholic acid inhibit with a potency almost equal to that of glycyrrhizic acid.

The compounds selected for study were chosen for a variety of reasons: the diuretics because they cause potassium loss with spironolactone as a control since it does not; glycyrrhizic acid and the bile salts, as reference compounds, since data about these compounds have been published and therefore they can be used in this study to evaluate relative potency of the other compounds studied; naringin and naringenin because they are active compounds in grapefruit juice that inhibit a particular pathway of drug oxidation (cytochrome P_{450} 3A4) and we were curious to see if they also inhibited this oxidation pathway (11β -OHSD); the sterols since they are present in vegetable oils and have a

structure suggesting that they might inhibit 11β -OHSD; and the cyclooxygenase inhibitors because they inhibit prostaglandin formation and cause salt retention.

The K_m of our enzyme preparation for cortisol (8μ mol/l) is similar to that of rat for corticosterone (2μ mol/l) found by Monder *et al.* [18]. Working with purified enzyme from rat liver (gift from Dr C. Monder), we have found an IC₅₀ of 12 nmol/l for glycyrrhizic acid [10], similar to the dissociation constant of the enzyme-inhibitor complex of 8 nmol/l reported by Monder *et al.* [18]. In a previous study from our laboratory, glycyrrhizic acid had an IC₅₀ of 1994 μ mol/l for guinea pig renal cortex microsomes with corticosterone as the substrate without Triton in the incubation mixture [9] compared with 254 μ mol/l in the present study using Triton and cortisol as the substrate. Buhler *et al.* [12] working with rat kidney microsomes and corticosterone at 0.1 μ mol/l, found an IC₅₀ of 4 μ M; in our study of guinea pig microsomes with a substrate concentration of 23 μ mol/l we found an IC₅₀ of 254 μ M. Perschel *et al.* [11] working with rat kidney microsomes found cholic acid to inhibit this

Table 2. Compounds that failed to inhibit 11β -OHSD

| Compound | Maximum concentration tested (μ mol/l) |
|---------------------|---|
| Bumetanide | 2000 |
| Hydrochlorothiazide | 8000 |
| Indomethacin | 1100 |
| Spironolactone | 2000 |

The maximum concentration tested was limited by the solubility of the compound.




Fig. 1. Inhibition of 11β -OHSD by glycyrrhizic acid from licorice and flavonoids from grapefruit juice.




Fig. 2. Double reciprocal plot of $1/v$ for 11β -OHSD with varying concentrations of furosemide in incubation mixture. K_m for cortisol is $8 \mu\text{mol/l}$. V_{max} is $30 \text{ nmol}/\mu\text{g}$ microsomal protein/h. K_i for furosemide is $7.7 \mu\text{mol/l}$.

enzyme at 1/27th the potency of chenodeoxycholic acid. We found it to be 1/7th the potency in our system.

We studied the NADP-requiring form of the enzyme that is present in most tissues rather than the NAD-requiring form that is present in the distal nephron [19–21]. Whether inhibition of the NAD-requiring enzyme is different from that of the NADP-requiring enzyme is not known. Since the mechanism of inhibition of glycyrrhetic acid [18], gossypol [9] and furosemide (this study) is competitive, one might speculate that competitive inhibition of the NAD-requiring form of the enzyme by these compounds might also occur.

The fact that furosemide is an inhibitor of the enzyme while bumetanide is not may explain why furosemide causes more potassium excretion per unit sodium excretion than bumetanide [22–24]. It is excreted by patients with heart failure at a rate of $15\text{--}30 \mu\text{g}/\text{min}$ [25]. Assuming a $1 \text{ ml}/\text{min}$ urine flow, the furosemide concentration would be $76 \mu\text{M}$, compared with its IC_{50} of $59 \mu\text{M}$ in this study.

Three flavonoids: the sugar conjugates of naringenin, quercetin and kaempferol, along with some others are found in grapefruit juice [26]. These are hydrolyzed in the intestine to the aglycons which are absorbed. We found that naringenin inhibited the enzyme in this study, and previously that the flavonoids morin and quercetin were weak inhibitors [10]. The importance, if any, of these dietary constituents as *in vivo* inhibitors of this enzyme remains to be determined.

REFERENCES

1. Ulick S., Ramirez L. C. and New M. I.: An abnormality in steroid reductive metabolism in hypertensive syndrome. *J. Clin. Endocr. Metab.* 44 (1977) 799–802.
2. Monder C.: Corticosteroids, receptors, and the organ-specific functions of 11β -hydroxysteroid dehydrogenase. *FASEB J.* 5 (1991) 3047–3054.
3. Funder J. W., Pearce P. T., Myles K. and Roy L. P.: Apparent mineralocorticoid excess, pseudohypoaldosteronism, and urinary electrolyte excretion: toward a redefinition of mineralocorticoid action. *FASEB J.* 4 (1990) 3234–3238.
4. Monder C. and White P. C.: 11β beta-hydroxysteroid dehydrogenase. *Vit. Horm.* 47 (1993) 187–271.
5. Edwards C. R. W., Walker B. R., Benediktsson R. and Seckl J. R.: Congenital and acquired syndromes of apparent mineralocorticoid excess. *J. Steroid Biochem. Molec. Biol.* 45 (1993) 1–5.
6. Edwards C. R. W.: Lessons from licorice. *New Engl. J. Med.* 325 (1991) 1242–1243.
7. Qian S. Z. and Wang Z. G.: Gossypol: a potential antifertility agent for males. *A. Rev. Pharmac. Toxic.* 24 (1984) 329–360.
8. Reidenberg M. M., Gu Z.-P., Lorenzo B., Coutinho E., Athayde C., Frick J., Alvarez F., Brache V. and Emuveyan E. E.: Differences in serum potassium concentrations in normal men in different geographic locations. *Clin. Chem.* 39 (1993) 72–75.
9. Sang G. W., Lorenzo B. and Reidenberg M. M.: Inhibitory effects of gossypol on corticosteroid 11β -hydroxysteroid dehydrogenase from guinea pig kidney: a possible mechanism for causing hypokalemia. *J. Steroid Biochem. Molec. Biol.* 39 (1991) 169–176.
10. Song D., Lorenzo B. and Reidenberg M. M.: Inhibition of 11β -hydroxysteroid dehydrogenase by gossypol and bioflavonoids. *J. Lab. Clin. Med.* 120 (1992) 792–797.
11. Perschel F. H., Buhler H. and Hierholzer K.: Bile acids and their amides inhibit 11β -hydroxysteroid dehydrogenase obtained from rat kidney. *Pflüger Archs* 418 (1991) 538–543.
12. Buhler H., Perschel F. H. and Hierholzer K.: Inhibition of rat renal 11β -hydroxysteroid dehydrogenase by steroidal compounds and triterpenoids; structure/function relationship. *Biochim. Biophys. Acta* 1075 (1991) 206–212.
13. Morris D. J., Semafuko W. E. B., Latif S. A., Vogel B., Grimes C. A. and Sheff M. F.: Detection of glycyrrhetic acid-like factors (GALFs) in human urine. *Hypertension* 20 (1992) 356–360.

14. Wehrauch J. L. and Gardner J. M.: Sterol content of foods of plant origin. *J. Am. Diet. Assoc.* 73 (1978) 39-47.
15. Bailey D. G., Spence J. D., Munoz C. and Arnold J. M. O.: Interaction of citrus juices with felodipine and nifedipine. *Lancet* 337 (1991) 268-269.
16. Miniscalco A., Lundahl J., Regardh C. G., Edgar B. and Eriksson U. G.: Inhibition of dihydropyridine metabolism in rat and human liver microsomes by flavonoids found in grapefruit in juice. *J. Pharmac.* 261 (1992) 1195-1199.
17. Soons P. A., Vogels B. A. P. M., Roosemalen M. C. M., Schoemaker H. C., Uchida E., Edgar B., Lundahl J., Cohen A. F. and Breimer D. D.: Grapefruit juice and cimetidine inhibit stereoselective metabolism of nitrendipine in humans. *Clin. Pharmac. Ther.* 50 (1991) 394-403.
18. Monder C., Stewart P. M., Lakshmi V., Valentino R., Burt D. and Edwards C. R. W.: Licorice inhibits corticosteroid 11β -dehydrogenase of rat kidney and liver: *in vivo* and *in vitro* studies. *Endocrinology* 125 (1989) 1046-1053.
19. Mercer W. R. and Krozowski Z. S.: Localization of an 11β hydroxysteroid dehydrogenase activity to the distal nephron. Evidence for the existence of two species of dehydrogenase in the rat kidney. *Endocrinology* 130 (1992) 540-543.
20. Walker B. R., Campbell J. C., Williams B. C. and Edwards C. R. W.: Tissue-specific distribution of NAD^+ -dependent isoform of 11β -hydroxysteroid dehydrogenase. *Endocrinology* 131 (1992) 970-972.
21. Monder C.: The forms and functions of 11β -hydroxysteroid dehydrogenase. *J. Steroid Biochem. Molec. Biol.* 45 (1993) 161-165.
22. Asbury M. J., Gatenby P. B. B., O'Sullivan S. and Bourke E.: Bumetanide: potent new "loop" diuretic. *Br. Med. J.* 1 (1972) 211-213.
23. Ramsay L. E., McInnes G. T., Hettiarachchi J., Shelton J. and Scott P.: Bumetanide and frusemide: a comparison of dose-response curves in healthy men. *Br. J. Clin. Pharmac.* 5 (1978) 243-247.
24. Brater D. C., Fox B. S. and Chennavasin P.: Electrolyte excretion, patterns, intravenous and oral doses of bumetanide compared to furosemide. *J. Clin. Pharmac.* 21 (1981) 599-603.
25. Brater D. C., Day B., Burdette A. and Anderson S.: Bumetanide and furosemide in heart failure. *Kidney Int.* 26 (1984) 183-189.
26. Kuhnau J.: The flavonoids. A class of semi-essential food components: Their role in human nutrition. *Wld Rev. Nutr. Diet.* 24 (1976) 117-191.